

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

School of Podiatric Medicine Publications and Presentations

School of Podiatric Medicine

---

8-4-2023

## Glycemic Markers and Insulin-related Parameters in the Progression of Diabetes and Lower Extremities Complications: A Pilot Study in Mexican Americans from Starr County

Peter Elvin

*The University of Texas Rio Grande Valley, peter.elvin@utrgv.edu*

Garrett Edwards

*The University of Texas Rio Grande Valley*

Walid Fakhouri

*University of Texas Health Science Center at Houston*

Craig Hanis

*University of Texas Health Science Center at Houston*

Cláudia Cristina Biguetti

*The University of Texas Rio Grande Valley, claudia.biguetti@utrgv.edu*

Follow this and additional works at: [https://scholarworks.utrgv.edu/sopm\\_pub](https://scholarworks.utrgv.edu/sopm_pub)



Part of the [Podiatry Commons](#)

---

### Recommended Citation

Elvin, P., Edwards, G., Fakhouri, W., Hanis, C., & Biguetti, C. (2023, August 4). Glycemic Markers and Insulin-related Parameters in the Progression of Diabetes and Lower Extremities Complications: A Pilot Study in Mexican Americans from Starr County. 2023 School of Podiatric Medicine Research Day, Harlingen, TX. [https://scholarworks.utrgv.edu/sopm\\_pub/5](https://scholarworks.utrgv.edu/sopm_pub/5)

This Conference Proceeding is brought to you for free and open access by the School of Podiatric Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in School of Podiatric Medicine Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

# Glycemic Markers and Insulin-related Parameters in the Progression of Diabetes and Lower Extremities Complications: A Pilot Study in Mexican Americans from Starr County

Peter Elvin<sup>1\*</sup>, Garrett Edwards<sup>1\*</sup>, Walid Fakhouri<sup>2</sup>, Craig Hanis<sup>3</sup>, Claudia Bigueti<sup>1</sup>

1. Department of Surgery and Biomechanics, School of Podiatric Medicine, University of Texas Rio Grande Valley

2. Department of Diagnostic and Biomedical Sciences, School of Dentistry at Houston, The University of Texas Health Science Center

3. Department of Epidemiology, Human Genetics and Environmental Sciences, School of Public Health, The University of Texas Health Science Center at Houston



Regenerative Medicine Research Laboratory

## INTRODUCTION/BACKGROUND

- Type 2 Diabetes (T2D) is a chronic disease involving alteration in glucose metabolism and uptake, often resulting in an uncontrolled hyperglycemic state
- Metabolic alterations can lead to neurologic and vascular compromise, allowing progression into lower extremity complications, like diabetic foot ulcers
- **Research Gap:** Guidelines linking glycemic metabolic parameters to disease progression remains limited. The extent to which markers can predict the progression of diabetes to lower limb complications has yet to be fully explored




**Figure 2.** Diabetic foot ulcer on plantar aspect of hallux. (Photo Courtesy of Dr. Luis Venegas)



**Figure 1.** 15-25% of diabetic patients may develop foot ulcers (1)

## POPULATION AND COLLECTION DATA



**Figure 3.** A) Descriptive analysis for a sample of 48 diabetic individuals (D), 48 non-diabetic individuals (ND), and 24 diabetic foot ulcer individuals (DFU). B) Age of the participants by time of data collection, C) Body Mass Index (BMI) D) Distribution of male and female. Values collected by the time of data collection. IRBs for this study HSC-SPH-06-225 and UTRGV-IRB22-0273

## HYPOTHESIS AND AIM

- **AIM:** To evaluate metabolic parameters in a Mexican American cohort of diabetics with foot ulcers (DFU), diabetic controls (D), and Non-diabetics (ND) from Starr County.

**HYPOTHESIS:** Disease development and progression to DFU is associated with imbalance in glycemic markers and insulin-related parameters from pre-diabetic to diabetic patients with DFUs.

### MARKERS MEASURED

- Fasting Glucose
- Use of Diabetic Medications
- HbA1C
- Metformin Use
- Fasting Insulin
- C-Peptide Levels

### STATISTICAL ANALYSIS


- Analysis of variance (ANOVA) with the post hoc Tukey's test and Kruskal-Wallis with Dunn's for significance was used for multiple comparisons.

- Chi-squared or Fisher's exact test for categorical variables.



**Figure 4.** Depiction of glycemic markers that were statistically significant and insignificant. Green checks indicate significance and red X's indicate insignificance.

## RESULTS



## CONCLUSION/FUTURE WORK

★ DFU individuals had elevated FG and HbA1c levels, as well as reduced use of metformin compared to D controls.

- Larger sample study would be required to validate our findings to establish if HbA1c, metformin use, and fasting glucose levels can be used as clinical predictors of diabetes progression into foot ulceration.

## REFERENCES

1. Yazdanpanah L, et al. (2018). Incidence and Risk Factors of Diabetic Foot Ulcer: A Population-Based Diabetic Foot Cohort (ADFC Study)-Two-Year Follow-Up Study. International journal of endocrinology, 2018, 7631659.

## ACKNOWLEDGMENT

Laboratory of Regenerative Medicine is supported by SOPM –UTRGV, UT System. Thanks to NIH/NCATS - UL1 Mechanism for the funding. Thank you Dr. Luis Venegas for diabetic foot ulcer photos. Thank you Dr. Hagner Andrade for your support in the poster preparation.

## CONTACT INFORMATION

**Advisor:** claudia.bigueti@utrgv.edu

**Student:** peter.elvin01@utrgv.edu