

2021

A Comparison of the Prevalence of Cirrhosis in a Hispanic and Non-Hispanic Population Based on Body Mass Index

Alexa Perlick

The University of Texas Rio Grande Valley, alexa.perlick01@utrgv.edu

Abaigeal Thompson

The University of Texas Rio Grande Valley, abaigeal.thompson01@utrgv.edu

Colton Wayne

The University of Texas Rio Grande Valley, colton.wayne01@utrgv.edu

Jose C. Maldonado

The University of Texas Rio Grande Valley, jose.campomaldonado@utrgv.edu

Angel Rendon

The University of Texas Rio Grande Valley, angel.rendon01@utrgv.edu

Follow this and additional works at: <https://scholarworks.utrgv.edu/som9331>



Part of the [Digestive System Diseases Commons](#)

Recommended Citation

Perlick, Alexa; Thompson, Abaigeal; Wayne, Colton; Maldonado, Jose C.; and Rendon, Angel, "A Comparison of the Prevalence of Cirrhosis in a Hispanic and Non-Hispanic Population Based on Body Mass Index" (2021). *MEDI 9331 Scholarly Activities Clinical Years*. 15.

<https://scholarworks.utrgv.edu/som9331/15>

This Article is brought to you for free and open access by the School of Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in MEDI 9331 Scholarly Activities Clinical Years by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Alexa Perlick
9331 Research Elective Report

Retrospective Study on Liver Cirrhosis in the Valley and Two Case Reports on COVID-19 and Palliative Care in the Setting of Traumatic Brain Injury

9331 Submission #1

Retrospective Study 1

Part 1.1 Abstract

Title: *A Comparison of the Prevalence of Cirrhosis in a Hispanic and Non-Hispanic Population Based on Body Mass Index*

Authors: Alexa Perlick, BS¹, Abbie Thompson, BS¹, Colton Wayne, BS¹, Angel Rendón, BS¹, and Jose Campo Maldonado¹, MD

Affiliations: University of Texas Rio Grande Valley School of Medicine¹

ABSTRACT

BACKGROUND:

The obesity epidemic is a growing public health concern. In addition to the already known complications and comorbidities associated with obesity, data suggest that obesity is an independent risk factor for the development of liver disease.^{1,2} However, there is a paucity of data regarding the clinical correlation of obesity and cirrhosis in a predominantly Hispanic population. Therefore, the aim of this study is to investigate the prevalence of cirrhosis stratified by BMI in a predominately Hispanic population.

RESULTS:

DISCUSSION:

Previous study by Jiao and colleagues observed a 4-fold higher prevalence of cirrhosis in Hispanics versus the general US population in South Texas.³ Therefore, these researchers also expect an increase in prevalence of cirrhosis in Hispanics compared to the US population. Previous studies have found an association between cirrhosis and body mass index.^{4,5} However, there is a lack in data in a predominantly Hispanic population.

REFERENCES

1. Berzigotti A, Garcia-Tsao G, Bosch J, et al. Obesity is an independent risk factor for clinical decompensation in patients with cirrhosis. *Hepatology*. 2011;54(2):555-561. doi:10.1002/hep.24418
2. Naveau S, Chaput JC, Fallik D, et al. Risk factors of fibrosis in alcohol-induced liver disease. *Hepatology*. 2002;35(3):635-638. doi:10.1053/jhep.2002.31782
3. Jiao J, Watt GP, Lee M, et al. Cirrhosis and advanced fibrosis in Hispanics in Texas: The dominant contribution of central obesity. *PLoS One*. 2016;11(3):1-14. doi:10.1371/journal.pone.0150978
4. Liu B, Balkwill A, Reeves G, Beral V. Body mass index and risk of liver cirrhosis in

middle aged UK women: Prospective study. *BMJ*. 2010;340(7747):633.
doi:10.1136/bmj.c912

5. Manco M, Bedogni G, Marcellini M, et al. Waist circumference correlates with liver fibrosis in children with non-alcoholic steatohepatitis. *Gut*. 2008.
doi:10.1136/gut.2007.142919

MANUSCRIPT

INTRODUCTION

The obesity epidemic is a growing public health concern. According to the CDC, the overall prevalence of obesity in the United States was 39.8% in 2015-2016. ¹ The prevalence of obesity is even higher in individuals of Hispanic origin (47.0%).¹ In addition to the already known complications and comorbidities associated with obesity, data suggest that obesity is an independent risk factor for the development of liver disease. ^{2,3} Obesity has been implicated as a risk factor for alcoholic-like inflammation, septal fibrosis, and progression to liver cirrhosis. ^{2,4,5} The adipose tissue in obesity is proinflammatory, proangiogenic, and profibrogenic in hepatic tissue. ^{6,7} A study from the United Kingdom found that an estimated 17% of cases of liver cirrhosis are attributable to excess body weight. ⁸ Moreover, each 5 kg/m² increase in body mass index (BMI) was associated with about 30% higher risk of hepatic mortality. ⁹

Studies have found ethnic differences in the incidence of cirrhosis and rate of disease progression. ^{10,11} Specifically, one study found a 4-fold higher prevalence of cirrhosis in Cameron County, Texas (0.94%) than the national prevalence (0.27%). ¹¹ According to the United States Census Bureau, 89.9% of the population of Cameron County is Hispanic. Furthermore, in the United States, chronic liver disease and cirrhosis are the 7th leading cause of death in Hispanics, 11th in Non-Hispanics white, and 14th in Non-Hispanic blacks. ¹² These ethnic variations are likely due to a complex interplay between biological, social and behavioral factors.

There is a paucity of data regarding the clinical correlation of obesity and cirrhosis in a predominantly Hispanic population. Additionally, data is limited on the contribution of BMI to the incidence of liver cirrhosis in Hispanics in South Texas. Therefore, the aim of this study is to investigate the prevalence of cirrhosis stratified by BMI in a predominately Hispanic population.

MATERIALS AND METHODS

This retrospective single-centered study was approved by the University of Texas Rio Grande Valley institutional ethics and review board. Since this is a retrospective study, informed consent was waived. We retrospectively reviewed patient electronic medical records admitted to Valley Baptist Medical Center in Harlingen, Texas with cirrhosis between October 2015 to November 2020 age > 18. In-hospital admission variables were collected through review of electronic medical records. These variables include basic demographic information, vital signs, social history, past medical history, and laboratory values. Additional data collected during hospital stay were also collected, such as abdominal imaging exams (e.g., ultrasound, CT of liver) and pathology results of liver biopsies.

RESULTS

Demographics and presenting features of the patients on admission are illustrated in Table 1.

FIGURES

Table 1. Characteristics of patients.

Variable	Total (N =)	Underweight (<18.5 kg/m ²) (n =)	Normal Weight (18.5 - 24.9 kg/m ²) (n =)	Overweight (25.0 – 29.9 kg/m ²) (n =)	Obese (≥ 30.0 kg/m ²) (n =)	P value
Female						
Male						
Age, years						
Hispanic						
Non-Hispanic						
Hypertension						
Hyperlipidemia						
Liver disease						
AST						
ALT						
Alkaline Phosphatase						
GGT						
Creatinine						
GFR						
Fasting plasma glucose						
Triglycerides						
LDL						
HDL						
HCV antibody						
HBV surface antigen						
Fasting insulin						
Platelets						
Total bilirubin						

DISCUSSION

The main objective of the current study is to identify the prevalence of cirrhosis stratified by BMI in a predominately Hispanic population.

Previous study by Jiao and colleagues observed a 4-fold higher prevalence of cirrhosis in Hispanics versus the general US population in South Texas.¹¹ Therefore, these researchers also expect an increase in prevalence of cirrhosis in Hispanics compared to the US population.

Previous studies have found an association between cirrhosis and body mass index, there is a lack in data in a predominantly Hispanic population.^{8,13}

REFERENCES

1. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of Obesity Among Adults and Youth: United States, 2015-2016. *NCHS Data Brief*. 2017;(288):1-8.
2. Berzigotti A, Garcia-Tsao G, Bosch J, et al. Obesity is an independent risk factor for clinical decompensation in patients with cirrhosis. *Hepatology*. 2011;54(2):555-561. doi:10.1002/hep.24418
3. Naveau S, Chaput JC, Fallik D, et al. Risk factors of fibrosis in alcohol-induced liver disease. *Hepatology*. 2002;35(3):635-638. doi:10.1053/jhep.2002.31782
4. Ratziu V, Giral P, Charlotte F, et al. Liver fibrosis in overweight patients. *Gastroenterology*. 2000;118(6):1117-1123. doi:10.1016/S0016-5085(00)70364-7
5. Ratziu V, Bonyhay L, Di Martino V, et al. Survival, liver failure, and hepatocellular carcinoma in obesity-related cryptogenic cirrhosis. *Hepatology*. 2002;35(6):1485-1493. doi:10.1053/jhep.2002.33324
6. Fernández M, Semela D, Bruix J, Colle I, Pinzani M, Bosch J. Angiogenesis in liver disease. *J Hepatol*. 2009;50(3):604-620. doi:10.1016/j.jhep.2008.12.011
7. Mejias M, Garci-Pras E, Tiani C, Miquel R, Bosch J, Fernandez M. Beneficial effects of sorafenib on splanchnic, intrahepatic, and portocollateral circulations in portal hypertensive and cirrhotic rats. *Hepatology*. 2009;49(4):1245-1256. doi:10.1002/hep.22758
8. Liu B, Balkwill A, Reeves G, Beral V. Body mass index and risk of liver cirrhosis in middle aged UK women: Prospective study. *BMJ*. 2010;340(7747):633. doi:10.1136/bmj.c912
9. MacMahon S, Baigent C, Duffy S, et al. Body-mass index and cause-specific mortality in 900 000 adults: Collaborative analyses of 57 prospective studies. *Lancet*. 2009;373(9669):1083-1096. doi:10.1016/S0140-6736(09)60318-4
10. Sajja KC, Mohan DP, Rockey DC. Age and ethnicity in cirrhosis. *J Investig Med*. 2014;62(7):920-926. doi:10.1097/JIM.000000000000106
11. Jiao J, Watt GP, Lee M, et al. Cirrhosis and advanced fibrosis in Hispanics in Texas: The dominant contribution of central obesity. *PLoS One*. 2016;11(3):1-14. doi:10.1371/journal.pone.0150978
12. Heron M. Heron 2017.pdf. *Natl Vital Stat Reports*. 2019;68(6):1-77.
13. Manco M, Bedogni G, Marcellini M, et al. Waist circumference correlates with liver fibrosis in children with non-alcoholic steatohepatitis. *Gut*. 2008. doi:10.1136/gut.2007.142919