

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

Mathematical and Statistical Sciences Faculty  
Publications and Presentations

College of Sciences

---

1-16-1995

## A completely integrable system and parametric representation of solutions of the Wadati-Konno-Ichikawa hierarchy

Zhijun Qiao

The University of Texas Rio Grande Valley, zhijun.qiao@utrgv.edu

Follow this and additional works at: [https://scholarworks.utrgv.edu/mss\\_fac](https://scholarworks.utrgv.edu/mss_fac)



Part of the [Mathematics Commons](#)

---

### Recommended Citation

Qiao, Zhijun, "A completely integrable system and parametric representation of solutions of the Wadati-Konno-Ichikawa hierarchy" (1995). *Mathematical and Statistical Sciences Faculty Publications and Presentations*. 17.

[https://scholarworks.utrgv.edu/mss\\_fac/17](https://scholarworks.utrgv.edu/mss_fac/17)

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

# A completely integrable system and parametric representation of solutions of the Wadati–Konno–Ichikawa hierarchy

Zhijun Qiao<sup>a1</sup>

*Institute of Mathematics, Fudan University, Shanghai 200433, People's Republic of China*

(Received 16 November 1994; accepted for publication 16 January 1995)

A finite-dimensional involutive system is presented, and the Wadati–Konno–Ichikawa (WKI) hierarchy of nonlinear evolution equations and their commutator representations are discussed in this article. By this finite-dimensional involutive system, it is proven that under the so-called Bargmann constraint between the potentials and the eigenfunctions, the eigenvalue problem (called the WKI eigenvalue problem) studied by Wadati, Konno, and Ichikawa [*J. Phys. Soc. Jpn.* **47**, 1698 (1979)] is nonlinearized as a completely integrable Hamiltonian system in the Liouville sense. Moreover, the parametric representation of the solution of each equation in the WKI hierarchy is obtained by making use of the solution of two compatible systems. © 1995 American Institute of Physics.

## I. INTRODUCTION

Since the beautiful “nonlinearization theory” came into use,<sup>1,2</sup> quite a few completely integrable Hamiltonian systems in the Liouville sense have been successively found in recent years.<sup>3–10</sup> This theory is used to not only search for new finite-dimensional involutive systems in explicit form, but also provide the representations of solutions for the soliton hierarchy. This article is devoted to the investigation of the Wadati–Konno–Ichikawa (WKI) hierarchy by the use of the “nonlinearization theory,” and is a continuation of a previous work.<sup>10</sup> It is well-known that the hierarchy of nonlinear evolution equations (called the WKI hierarchy below) related to the eigenvalue problem (called the WKI eigenvalue problem below) presented by Wadati, Konno, and Ichikawa were obtained about eleven years ago.<sup>11</sup> But discussions on the representations of solutions of the WKI hierarchy were very few. In this article, we first give an involutive system of functions, which guarantees integrability of the nonlinearized WKI eigenvalue problem under the so-called Bargmann constraint. Then, giving the WKI hierarchy of nonlinear evolution equations by the spectral gradient method<sup>12</sup> and their commutator representations, and using the solution of two compatible equations, we present the parametric representation of the solution for each equation in the WKI hierarchy.

## II. A FINITE-DIMENSIONAL INVOLUTIVE SYSTEM $\{E_k\}_{k=1}^N$ AND HAMILTONIAN SYSTEMS $(F_m)$

Denote the standard Poisson bracket of two functions  $F$  and  $G$  on the symplectic manifold  $(R^{2N}, dp \wedge dq)$  by<sup>13</sup>

$$(F, G) = \sum_{j=1}^N \left( \frac{\partial F}{\partial q_j} \frac{\partial G}{\partial p_j} - \frac{\partial F}{\partial p_j} \frac{\partial G}{\partial q_j} \right) = \left\langle \frac{\partial F}{\partial q}, \frac{\partial G}{\partial p} \right\rangle - \left\langle \frac{\partial F}{\partial p}, \frac{\partial G}{\partial q} \right\rangle,$$

which is skew symmetric, bilinear, and satisfies the Jacobi identity and the Leibnitz rule:  $(EF, H) = E(F, H) + F(E, H)$ , where  $\langle \cdot, \cdot \rangle$  stands for the standard inner product in  $R^{2N}$ .

<sup>a1</sup>Permanent address: Department of Mathematics, Liaoning University, Shenyang 110036, P. R. China.

Let  $\lambda_1, \dots, \lambda_N$  be  $N$  different constants and  $\lambda_1 < \dots < \lambda_N$ . Write

$$\Gamma_k = \sum_{\substack{j=1 \\ j \neq k}}^N \frac{\lambda_k \lambda_j B_{kj}^2}{\lambda_k - \lambda_j}, \quad B_{kj} = p_k q_j - q_k p_j. \quad (1)$$

Then through a lengthy calculation, we obtain the following proposition.

**Proposition 1:** A system of functions  $E_1, \dots, E_N$  defined as follows consists of a new finite-dimensional involutive system:

$$E_k = -\langle \Lambda p, q \rangle p_k q_k + \sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle} p_k q_k + \frac{1}{2} \Gamma_k, \quad k = 1, 2, \dots, N, \quad (2)$$

where  $q = (q_1, \dots, q_N)^T$ ,  $p = (p_1, \dots, p_N)^T$ ,  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_N)$ , and  $\langle \xi, \eta \rangle = \sum_{j=1}^N \xi_j \eta_j$ .

Define a bilinear function  $Q_z(\xi, \eta)$  on  $R^N$  as follows:<sup>14</sup>

$$Q_z(\xi, \eta) = \langle (z - \Lambda)^{-1} \xi, \eta \rangle = \sum_{k=1}^N (z - \lambda_k)^{-1} \xi_k \eta_k. \quad (3)$$

Then the generated functions of the involutive system  $\{E_k\}$  are

$$-\langle \Lambda p, q \rangle Q_z(p, q) + \sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle} Q_z(p, q) + \frac{1}{2} \begin{vmatrix} Q_z(\Lambda q, q) & Q_z(\Lambda q, p) \\ Q_z(\Lambda p, q) & Q_z(\Lambda p, p) \end{vmatrix} = \sum_{k=1}^N \frac{E_k}{z - \lambda_k}. \quad (4)$$

**Proposition 2:** Let  $F_m = \sum_{k=1}^N \lambda_k^m E_k$ ,  $m = 0, 1, 2, \dots$ . Then we have

$$F_0 = -\langle \Lambda p, q \rangle \langle p, q \rangle + \sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle} \langle p, q \rangle, \quad (5)$$

$$F_m = -\langle \Lambda p, q \rangle \langle \Lambda^m p, q \rangle + \sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle} \langle \Lambda^m p, q \rangle + \frac{1}{2} \sum_{j=0}^{m-1} \begin{vmatrix} \langle \Lambda^{j+1} q, q \rangle & \langle \Lambda^{j+1} q, p \rangle \\ \langle \Lambda^{m-j} p, q \rangle & \langle \Lambda^{m-j} p, p \rangle \end{vmatrix} \quad (6)$$

and  $(F_m, F_l) = 0$ ,  $\forall m, l \in Z^+$ .

*Proof:* Obviously,  $(F_m, F_l) = 0$ . When  $|z| > \max\{|\lambda_1|, \dots, |\lambda_N|\}$ , we get

$$(z - \lambda_k)^{-1} = \sum_{m=0}^{\infty} z^{-m-1} \lambda_k^m, \quad Q_z(\xi, \eta) = \sum_{m=0}^{\infty} \langle \Lambda^m \xi, \eta \rangle z^{-m-1}.$$

Hence, Eqs. (5) and (6) are deduced by virtue of substituting the Laurent expansion of  $Q_z(\xi, \eta)$  and the power series of  $(z - \lambda_k)^{-1}$  into both sides of Eq. (5), and sorting terms with the power of  $z$ .

By virtue of this proposition, we promptly have

**Theorem 1:** The Hamiltonian systems

$$(F_m): \quad q_{t_m} = \frac{\partial F_m}{\partial p}, \quad p_{t_m} = -\frac{\partial F_m}{\partial q}, \quad m = 0, 1, 2, \dots, \quad (7)$$

are completely integrable in the Liouville sense.

### III. THE WKI HIERARCHY AND COMMUTATOR REPRESENTATIONS

In this section, we first give the WKI hierarchy of nonlinear evolution equations by the spectral gradient  $\nabla\lambda$  of eigenvalue  $\lambda$  with respect to the potentials  $u$  and  $v$ , then present the commutator representation of each equation in the WKI hierarchy.

Consider the WKI eigenvalue problem<sup>15</sup>

$$y_x = My, \quad M = \begin{pmatrix} -i\lambda & \lambda u \\ \lambda v & i\lambda \end{pmatrix}, \quad i^2 = -1, \tag{8}$$

i.e.,

$$Ly = \lambda y, \quad L = L(u, v) = \frac{1}{1-uv} \begin{pmatrix} i & -u \\ -v & -i \end{pmatrix} \partial, \quad \partial = \frac{\partial}{\partial x}. \tag{9}$$

Let  $y = (y_1, y_2)^T$  satisfy Eq. (9) and  $\nabla\lambda = (\lambda y_2^2, -\lambda y_1^2)^T$ . Then  $\nabla\lambda$  satisfies

$$K\nabla\lambda = \lambda \cdot J\nabla\lambda, \tag{10}$$

where  $K$  and  $J$  are two skew-symmetric operators ( $\partial = \partial/\partial x, \partial\partial^{-1} = \partial^{-1}\partial = 1$ )

$$K = \frac{1}{2i} \begin{pmatrix} -\frac{1}{2} \partial^2 \frac{u}{w} \partial^{-1} \frac{u}{w} \partial^2 & \partial^3 + \frac{1}{2} \partial^2 \frac{u}{w} \partial^{-1} \frac{v}{w} \partial^2 \\ \partial^3 + \frac{1}{2} \partial^2 \frac{v}{w} \partial^{-1} \frac{u}{w} \partial^2 & -\frac{1}{2} \partial^2 \frac{v}{w} \partial^{-1} \frac{v}{w} \partial^2 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & -\partial^2 \\ \partial^2 & 0 \end{pmatrix}, \quad w = \sqrt{1-uv}, \tag{11}$$

which are called the pair of Lenard's operators of Eq. (8).

Now, recursively define the Lenard's gradient sequence  $\{G_m\}_{m=-1}^\infty$  as follows:

$$\begin{cases} KG_{m-1} = JG_m, & (m=0, 1, 2, \dots) \\ G_{-1} = (1, 1)^T \in \ker J, & G_0 = \begin{pmatrix} v/w & u/w \end{pmatrix}^T, \end{cases} \tag{12}$$

which exactly generate the WKI hierarchy of the soliton equation

$$(u, v)_t^T = J(J^{-1}K)^m G_0, \quad m=0, 1, 2, \dots \tag{13}$$

Particularly, as  $v = -1, (u, v)_t^T = KG_0 = JG_1$  can be reduced to the well-known Harry-Dyn equation  $s_t = -(1/\sqrt{s})_{xxx}$  with  $s = 1 + u$ . The WKI hierarchy of soliton equations (13) possess the following commutator (or Lax) representations:

$$L_t = [W_m, L], \quad m=0, 1, 2, \dots,$$

$$W_m = \sum_{j=0}^m \left\{ \frac{1}{2i} \begin{pmatrix} 0 & (G_{j-1}^{(2)})_x - iuA(G_{j-1})_x \\ (G_{j-1}^{(1)})_x + ivA(G_{j-1})_x & 0 \end{pmatrix} L^{m+1-j} \right.$$

$$\left. + iA(G_{j-1}) \begin{pmatrix} -i & u \\ v & i \end{pmatrix} L^{m+2-j} \right\}, \tag{14}$$

with  $A(G_{j-1}) = (1/2iw)\partial^{-1}((u/w)G_{j-1}^{(1)} - (v/w)G_{j-1}^{(2)})_x, w = \sqrt{1-uv}$ , where  $G_{j-1} = (G_{j-1}^{(1)}, G_{j-1}^{(2)})^T (j=0, 1, 2, \dots, m)$  are determined by Eq. (12),  $L$  is defined by Eq. (9).

*Remark:* In Eq. (14), the operators  $W_m$  ( $m=0,1,2,\dots$ ) are a set of isospectral Lax operators<sup>16</sup> of the WKI spectral problem (9), which can compose an operator Lie algebra. These results are going to be reported in another article.

**IV. INTEGRABILITY OF THE NONLINEARIZED WKI SPECTRAL PROBLEM (8)**

Let  $\lambda_1, \dots, \lambda_N$  be  $N$  different eigenvalues of Eq. (8), and  $y = (q_j, p_j)^T$  be the eigenfunctions corresponding to  $\lambda_j$  ( $j = 1, 2, \dots, N$ ). The so-called Bargmann constraint (Ref. 3):  $G_0 = \sum_{j=1}^N \nabla \lambda_j$  reads

$$u = \frac{i\langle \Lambda q, q \rangle}{\sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle}}, \quad v = \frac{-i\langle \Lambda p, p \rangle}{\sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle}}. \tag{15}$$

*Proposition 3:* Under the Bargmann constraint (15), the nonlinearized system of the WKI eigenvalue problem (9) can be expressed as the Hamiltonian structure

$$\begin{cases} q_x = -i\Lambda q + \frac{i\langle \Lambda q, q \rangle}{\sqrt{1 + \langle \Lambda q, q \rangle \langle \Lambda p, p \rangle}} \Lambda p = \frac{\partial H}{\partial p} \\ p_x = i\Lambda p - \frac{i\langle \Lambda p, p \rangle}{\sqrt{1 + \langle \Lambda q, q \rangle \langle \Lambda p, p \rangle}} \Lambda q = -\frac{\partial H}{\partial q}, \end{cases} \tag{16}$$

with the Hamiltonian function

$$H = -i\langle \Lambda p, q \rangle + i\sqrt{1 + \langle \Lambda q, q \rangle \langle \Lambda p, p \rangle}. \tag{17}$$

*Proof:* Obvious.

*Proposition 4:* The Hamiltonian function  $H$  and  $F_m$  are involutive, i.e.,

$$(H, F_m) = 0, \quad \forall m \in Z^+. \tag{18}$$

*Proof:* Equation (18) is obtained through substituting Eqs. (17) and (6) into the Poisson bracket

$$(H, F_m) = \left\langle \frac{\partial H}{\partial q}, \frac{\partial F_m}{\partial p} \right\rangle - \left\langle \frac{\partial H}{\partial p}, \frac{\partial F_m}{\partial q} \right\rangle$$

and carefully calculating it.

According to this proposition, we get

**Theorem 2:** The Hamiltonian system (16) is completely integrable in the Liouville sense.

**V. PARAMETRIC REPRESENTATIONS OF SOLUTIONS**

$(H, F_m) = 0$  implies that the Hamiltonian systems ( $H$ ) and ( $F_m$ ) are compatible, and thus their corresponding phase flow  $g_0^x g_m^{t_m}$  commute (see Ref. 13). Define

$$\begin{pmatrix} q(x, t_m) \\ p(x, t_m) \end{pmatrix} = g_0^x g_m^{t_m} \begin{pmatrix} q(0, 0) \\ p(0, 0) \end{pmatrix},$$

which is an analytic function of  $(x, t_m)$ .

**Theorem 3:** Let  $(q(x, t_m), p(x, t_m))^T$  be a solution of the compatible system ( $H$ ) and ( $F_m$ ). Then the nonlinear WKI evolution equations

$$4i \begin{pmatrix} u \\ v \end{pmatrix}_{t_{m+2}} = J(J^{-1}K)^m G_0, \quad m=0,1,2,\dots, \quad (19)$$

possess the parametric representations of solutions as follows:

$$\begin{aligned} u(x, t_m) &= \frac{i \langle \Lambda q(x, t_m), q(x, t_m) \rangle}{\sqrt{1 + \langle \Lambda q(x, t_m), q(x, t_m) \rangle \langle \Lambda p(x, t_m), p(x, t_m) \rangle}}, \\ v(x, t_m) &= \frac{-i \langle \Lambda p(x, t_m), p(x, t_m) \rangle}{\sqrt{1 + \langle \Lambda q(x, t_m), q(x, t_m) \rangle \langle \Lambda p(x, t_m), p(x, t_m) \rangle}}, \end{aligned} \quad (20)$$

where  $J, K$  are expressed by Eq. (11).

*Proof:* Let  $Q = \sqrt{1 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle}$ . Then, from Eq. (20) we have

$$\begin{pmatrix} u \\ v \end{pmatrix}_{t_{m+2}} = iQ^{-3} \langle \Lambda q, q_{t_{m+2}} \rangle \begin{pmatrix} 2 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle \\ \langle \Lambda p, p \rangle^2 \end{pmatrix} - iQ^{-3} \langle \Lambda p, p_{t_{m+2}} \rangle \begin{pmatrix} \langle \Lambda q, q \rangle^2 \\ 2 + \langle \Lambda p, p \rangle \langle \Lambda q, q \rangle \end{pmatrix}. \quad (21)$$

In addition, letting the operator  $(J^{-1}K)^m$  operate upon the equality  $G_0 = \sum_{j=1}^N \nabla \lambda_j$ , and noticing Eqs. (10) and (12), we get

$$(J^{-1}K)^m G_0 = \sum_{j=1}^N \lambda_j^m \nabla \lambda_j. \quad (22)$$

After Eqs. (6) and (7) are substituted into Eq. (21), through a series of careful calculations, Eq. (21) reads as

$$4i \begin{pmatrix} u \\ v \end{pmatrix}_{t_{m+2}} = \begin{pmatrix} 0 & -\partial^2 \\ \partial^2 & 0 \end{pmatrix} \begin{pmatrix} \langle \Lambda^{m+1} p, p \rangle \\ -\langle \Lambda^{m+1} q, q \rangle \end{pmatrix} = J \sum_{j=1}^N \lambda_j^m \nabla \lambda_j = J(J^{-1}K)^m G_0,$$

where Eq. (22) is used.

*Remark:* The C. Neumann constraint of the WKI hierarchy and its corresponding integrability, have already been discussed in another article.<sup>17</sup>

## ACKNOWLEDGMENTS

The author would like to express his sincerest thanks to Professor Gu Chao hao and Professor Hu Hesheng for their enthusiastic encouragement and guidance. The author is also grateful to Associate Professor Ma Wenxiu, Dr. Zhou Ruguang, and Dr. Lu Baoqun for helpful discussions.

This work has been supported by The Doctoral Program Foundation of Institution High Education and Natural Science Foundation of Shanghai, China. This work has also been supported by the Shanghai Science and Technology "Morning Star" Planning for the Project.

<sup>1</sup>C. Cao, *Sci. China A* **33**, 528 (1990).

<sup>2</sup>C. Cao, *Chin. Q. J. Math.* **3**, 90 (1988).

<sup>3</sup>C. Cao and X. Geng, *Research Reports in Physics: Nonlinear Physics*, edited by C. Gu *et al.* (Springer-Verlag, Berlin, 1990), pp. 68–78.

<sup>4</sup>C. Cao and X. Geng, *J. Phys. A* **23**, 4117 (1990).

<sup>5</sup>C. Cao and X. Geng, *J. Math. Phys.* **32**, 2323 (1991).

<sup>6</sup>M. Antonowicz and S. Rauch-Wojciechowski, *Phys. Lett. A* **147**, 455 (1990).

<sup>7</sup>Y. Zeng and Y. Li, *J. Math. Phys.* **30**, 1679 (1990).

<sup>8</sup>X. Geng, *Physica A* **180**, 241 (1992).

- <sup>9</sup>X. Geng, Phys. Lett. A **162**, 375 (1992).  
<sup>10</sup>Z. Qiao, Phys. Lett. A **172**, 224 (1993).  
<sup>11</sup>M. Boiti, F. Pempinelli, and G. Tu, Prog. Theor. Phys. **69**, 48 (1983).  
<sup>12</sup>B. Fuchssteiner, "Applications of spectral gradient methods to nonlinear evolution equations," preprint (1979); A. S. Fokas and R. L. Anderson, J. Math. Phys. **23**, 1066 (1982).  
<sup>13</sup>V. I. Arnol'd, *Mathematical Methods of Classical Mechanics* (Springer-Verlag, Berlin, 1978).  
<sup>14</sup>C. Cao, Henan Sci. **5**, 1 (1987).  
<sup>15</sup>M. Wadati, K. Konno, and Y. H. Ichikawa, J. Phys. Soc. Jpn. **47**, 1698 (1979).  
<sup>16</sup>W. Ma, J. Phys. A **25**, 5329 (1992).  
<sup>17</sup>Z. Qiao, J. Math. Res. Exp. **13**, 377 (1993).