

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Mathematical and Statistical Sciences Faculty
Publications and Presentations

College of Sciences

2-2019

Level 17 Ramanujan-sato Series

Timothy Huber

The University of Texas Rio Grande Valley

Daniel Schultz

Dongxi Ye

Follow this and additional works at: https://scholarworks.utrgv.edu/mss_fac



Part of the [Mathematics Commons](#)

Recommended Citation

Huber, Timothy; Schultz, Daniel; and Ye, Dongxi, "Level 17 Ramanujan-sato Series" (2019). *Mathematical and Statistical Sciences Faculty Publications and Presentations*. 32.

https://scholarworks.utrgv.edu/mss_fac/32

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

LEVEL 17 RAMANUJAN-SATO SERIES

TIM HUBER, DANIEL SCHULTZ, AND DONGXI YE

ABSTRACT. Two level 17 modular functions

$$r = q^2 \prod_{n=1}^{\infty} (1 - q^n)^{\left(\frac{n}{17}\right)}, \quad s = q^2 \prod_{n=1}^{\infty} \frac{(1 - q^{17n})^3}{(1 - q^n)^3},$$

are used to construct a new class of Ramanujan-Sato series for $1/\pi$. The expansions are induced by modular identities similar to those level of 5 and 13 appearing in Ramanujan's Notebooks. A complete list of rational and quadratic series corresponding to singular values of the parameters is derived.

1. INTRODUCTION

Let τ be a complex number with positive imaginary part and set $q = e^{2\pi i\tau}$. Define

$$r(\tau) = q^2 \prod_{n=1}^{\infty} (1 - q^n)^{\left(\frac{n}{17}\right)}, \quad s(\tau) = q^2 \prod_{n=1}^{\infty} \frac{(1 - q^{17n})^3}{(1 - q^n)^3}.$$

In this paper, we derive level 17 Ramanujan-Sato expansions for $1/\pi$ of the form

$$q \frac{d}{dq} \log s = \sum_{n=0}^{\infty} A_n \left(\frac{r(r^2s + 8rs - r - s)}{8r^3s - 3r^2s + r - s} \right)^n, \quad \frac{1}{s} = r + \frac{1}{r} - 2\sqrt{\frac{4}{r} - 4r - 15}, \quad (1.1)$$

where A_n is defined recursively. These relations are analogous to those at level 13 and 5 [5, 12],

$$q \frac{d}{dq} \log \mathcal{S} = \sum_{n=0}^{\infty} \mathcal{A}(n) \left(\frac{\mathcal{R}(1 - 3\mathcal{R} - \mathcal{R}^2)}{(1 + \mathcal{R}^2)^2} \right)^n, \quad \frac{1}{\mathcal{S}} = \frac{1}{\mathcal{R}} - 3 - \mathcal{R}, \quad (1.2)$$

$$q \frac{d}{dq} \log S = \sum_{n=0}^{\infty} a(n) \left(\frac{R^5(1 - 11R^5 - R^{10})}{(1 + R^{10})^2} \right)^n, \quad \frac{1}{S} = \frac{1}{R^5} - 11 - R^5. \quad (1.3)$$

Here $a(n)$, $\mathcal{A}(n)$ are recursively defined sequences induced from differential equations and

$$\mathcal{R}(\tau) = q \prod_{n=1}^{\infty} (1 - q^n)^{\left(\frac{n}{13}\right)}, \quad R(\tau) = q^{1/5} \prod_{n=1}^{\infty} (1 - q^n)^{\left(\frac{n}{5}\right)}. \quad (1.4)$$

Identity (1.3) and explicit evaluations for $R(\tau)$ were used to formulate expansions for $1/\pi$ including

$$\frac{1}{\pi} = \frac{1705}{81\sqrt{47}} \sum_{n=0}^{\infty} a(n) \left(n + \frac{71}{682} \right) \left(\frac{-1}{15228} \right)^n, \quad a(n) = \binom{2n}{n} \sum_{j=0}^n \binom{n}{j}^2 \binom{n+j}{j}. \quad (1.5)$$

This expression is a generalization of 17 such formulas stated by Ramanujan [3, 18]. In each formula, the algebraic constants come from explicit evaluations for a modular function. The sequence A_k are coefficients in a series solution to a differential equation satisfied by a relevant modular form. Generalizing such relations to higher levels requires finding differential equations for modular parameters and relevant identities. The series [5, 9, 10, 12] have common construction for primes $p - 1 \mid 24$, where $X_0(p)$ has genus zero. More work remains to unify constructions for other levels.

The purpose of this paper is to construct level 17 Ramanujan-Sato series as a prototype for levels such that $X_0(N)$ has positive genus. Central to the construction is the fact that r and s generate the field of functions invariant under action by an index two subgroup of $\Gamma_0(17)$. These constructions and singular value evaluations yield new Ramanujan-Sato expansions, including the rational series

$$\frac{1}{\pi} = \frac{1}{\sqrt{11}} \sum_{k=0}^{\infty} A_k \frac{307 + 748k}{(-21)^{k+2}}. \quad (1.6)$$

Following [5, 6], an expansion for $1/\pi$ is said to be rational or quadratic if C/π can be expressed as a series of algebraic numbers of degree 1 or 2, respectively, for some algebraic number C . We derive a complete list of series of rational and quadratic series from singular values of parameters in (1.1).

In the next section, we give an overview of results at levels 5 and 13 that motivate the approach of the paper. Section 3 includes an analogous construction of level 17 modular functions. This construction is used to motivate the differential equation satisfied by

$$z(\tau) = \theta_q \log s, \quad \theta_q := q \frac{d}{dq}, \quad (1.7)$$

with coefficients in the field $\mathbb{C}(x)$, where

$$x(\tau) = \frac{r(r^2s + 8rs - r - s)}{8r^3s - 3r^2s + r - s}. \quad (1.8)$$

We conclude with Section 4 in which singular values are derived for x and used to construct a new class of series approximations for $1/\pi$ of level 17. We derive a complete list of values of $x(\tau)$ with $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$ within the radius of convergence for z as a powers series in x and therefore provide a complete list of linear and quadratic Ramanujan-Sato series corresponding to $x(\tau)$.

2. LEVEL 5 AND 13 SERIES

The product $R(\tau)$, defined by (1.4), is the Rogers-Ramanujan continued fraction [19]. Together, $R(\tau)$ and $S(\tau)$, defined by

$$R(\tau) = \frac{q^{1/5}}{1 + \frac{q}{q + \frac{q^2}{1 + \frac{q^3}{1 + \dots}}}}, \quad S(\tau) = q \prod_{n=1}^{\infty} \frac{(1 - q^{5n})^6}{(1 - q^n)^6}, \quad (2.1)$$

generate the field of functions invariant under the congruence subgroup

$$\Gamma_0^2(d) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid c \equiv 0 \pmod{p} \text{ and } \chi(d) = 1 \right\}.$$

This motivates Ramanujan's reciprocal identity [17], [1, p. 267]

$$\frac{1}{R^5} - 11 - R^5 = \frac{1}{S}. \quad (2.2)$$

Equation (1.3) expresses the logarithmic derivative $Z = \theta_q \log S(\tau)$ in terms of a series solution to a third order linear differential equation [5] satisfied by S and the $\Gamma_0(5)$ invariant function $T = T(\tau)$

$$(16T^2 + 44T - 1)Z_{TTT} + (48T^2 + 66T)Z_{TT} + (44T^2 + 34T)Z_T + (12T^2 + 6T)Z = 0, \quad (2.3)$$

$$Z_T = T \frac{d}{dT} Z, \quad T = \frac{R^5(1 - 11R^5 - R^{10})}{(1 + R^{10})^2}. \quad (2.4)$$

The form of the equation may be anticipated from a general theorem [22] (c.f. [24]).

Theorem 2.1. *Let Γ be subgroup of $SL_2(\mathbb{R})$ commensurable with $SL_2(\mathbb{Z})$. If $t(\tau)$ is a nonconstant meromorphic modular function and $F(\tau)$ is a meromorphic modular form of weight k with respect to Γ , then $F, \tau F, \dots, \tau^k F$, as functions of t , are linearly independent solutions to a $(k+1)$ st order differential linear equation with coefficients that are algebraic functions of t . The coefficients are polynomials when $\Gamma \setminus \mathfrak{H}$ has genus zero and t generates the field of modular functions on Γ .*

Therefore, from (2.3),

$$Z = \sum_{k=0}^{\infty} a(n)T^n, \quad |T| < \frac{5\sqrt{5}-11}{8}, \quad (2.5)$$

where $a(n)$ is recursively determined from (2.3), and expressible in closed form [5] in terms of the summand appearing in (1.5). The final ingredient needed for Ramanujan-Sato series at level 5 are explicit evaluations for the Rogers-Ramanujan continued fraction within the radius of convergence of the power series. Such singular values for $R(\tau)$ were given by Ramanujan in his first letter to Hardy [2] and can be derived from modular equations satisfied by $T(\tau)$ and $T(n\tau)$. We provide a general approach in Section 4.

To formulate the analogous construction at level 13, define $\mathcal{R} = \mathcal{R}(\tau)$ by (1.4) and

$$\mathcal{S}(\tau) = q \prod_{n=1}^{\infty} \frac{(1-q^{13n})^2}{(1-q^n)^2}, \quad \mathcal{T}(\tau) = \frac{\mathcal{R}(1-3\mathcal{R}-\mathcal{R}^2)}{(1+\mathcal{R}^2)^2}. \quad (2.6)$$

A third order linear differential equation [12] is satisfied by the Eisenstein series $\mathcal{Z}(\tau) = \theta_q \log \mathcal{S}$ with coefficients that are polynomials in the $\Gamma_0(13)$ invariant function $\mathcal{T}(\tau)$. For both the level 5 and 13 cases, the weight zero functions T and \mathcal{T} may be uniformly presented as the quotient of a weight 4 cusp form and the square of a weight 2 Eisenstein series

$$\mathcal{T} = \frac{\mathcal{U}\mathcal{V}}{\mathcal{Z}^2}, \quad T = \frac{UV}{Z^2}, \quad (2.7)$$

where $\mathcal{U}(\tau) = \theta_q \log \mathcal{R}$, $U(\tau) = \theta_q \log R$,

$$\mathcal{V}(\tau) = \sum_{n=1}^{\infty} \binom{n}{13} \frac{q^n}{(1-q^n)^2}, \quad V(\tau) = \sum_{n=1}^{\infty} \binom{n}{5} \frac{q^n}{(1-q^n)^2}. \quad (2.8)$$

Both levels require singular values for \mathcal{T}, T [11]. Explicit evaluations for \mathcal{Z} and Z follow from

$$\mathcal{W} = \frac{\theta_q \log \mathcal{T}}{\mathcal{Z}} = \sqrt{1-12\mathcal{T}-16\mathcal{T}^2}, \quad W = \frac{\theta_q \log T}{Z} = \sqrt{1-44T+16T^2}. \quad (2.9)$$

The pairs $(\mathcal{T}, \mathcal{W})$, (T, W) , respectively, generate the field of invariant functions for $\Gamma_0(13)$, $\Gamma_0(5)$, and r and s generate invariant function fields for the congruence subgroup $\Gamma_0(17)$.

Proposition 2.2. *Let $A_0(\Gamma)$ denote the field of functions invariant under Γ and denote by χ the real quadratic character modulo p . Then*

- (1) $A_0(\Gamma_0(5)+) = \mathbb{C}(T)$ and $A_0(\Gamma_0(5)) = \mathbb{C}(T, W)$.
- (2) $A_0(\Gamma_0(13)+) = \mathbb{C}(\mathcal{T})$ and $A_0(\Gamma_0(13)) = \mathbb{C}(\mathcal{T}, \mathcal{W})$.
- (3) For prime $p \equiv 1 \pmod{4}$, $A_0(\Gamma_0^2(p)) = \mathbb{C}(R_p, S_p)$, where

$$R_p = q^{\ell_p} \prod_{n=1}^{\infty} (1-q^n)^{\chi(n)}, \quad S_p = q^{a_p} \prod_{n=1}^{\infty} \frac{(1-q^{np})^{b_p}}{(1-q^n)^{b_p}}, \quad (2.10)$$

$$\ell_p = \sum_{n=1}^{\frac{p-1}{2}} \frac{n(n-p)}{2p} \chi(n), \quad \frac{p-1}{24} = \frac{a_p}{b_p}, \quad \gcd(a_p, b_p) = 1. \quad (2.11)$$

A proof of the first two parts of Proposition 2.2 may be given along the lines of the proof of Proposition 3.2. The third part of the Proposition is a main result of [15]. The results of [15] explain Ramanujan's level 5 reciprocal relation (2.2) and his level 13 reciprocal relation [1, Equation (8.4)]

$$\frac{1}{\mathcal{R}} - 3 - \mathcal{R} = \frac{1}{\mathcal{S}}. \quad (2.12)$$

For our present work at level 17, we apply a new identity proven in [15]

$$r + \frac{1}{r} - 2\sqrt{\frac{4}{r} - 4r - 15} = \frac{1}{s}. \quad (2.13)$$

Our next task is to construct functions analogous to T and W in terms of r, s and Eisenstein series.

3. FUNCTIONS INVARIANT UNDER $\Gamma_0(17)$ AND A DIFFERENTIAL EQUATION

In this Section we prove an analogue to Proposition 2.2 and derive a second order linear differential equation for z defined by (1.7) with coefficients in $\mathbb{C}(x)$, where x is defined by (1.8). In order to construct functions that are invariant under $\Gamma_0(17)$, we introduce sums of eight Eisenstein series considered in [14]. Set

$$\mathcal{E}_1(\tau) := \frac{1}{8} \sum_{\chi(-1)=-1} E_{\chi,k}(\tau), \quad E_{\chi,k}(\tau) = 1 + \frac{2}{L(1-k, \chi)} \sum_{n=1}^{\infty} \chi(n) \frac{n^{k-1} q^n}{1-q^n}, \quad (3.1)$$

where the sum in (3.1) is over the odd primitive Dirichlet characters modulo 17 and $L(1-k, \chi)$ is the analytic continuation of the associated Dirichlet L -series and $\chi(-1) = (-1)^k$. For $a \in (\mathbb{Z}/17\mathbb{Z})^*$, apply the diamond operator [13] to define, for $1 \leq k \leq 8$,

$$\langle a \rangle \mathcal{E}_1(\tau) = \frac{1}{8} \sum_{\chi(-1)=-1} \chi(a) E_{\chi,1}(\tau), \quad \mathcal{E}_k(\tau) = \pm(3)^{k-1} \mathcal{E}_1(\tau). \quad (3.2)$$

The sign in Equation (3.2) is chosen so that the first coefficient in the q -series expansion is 1. The parameters $\mathcal{E}_k(\tau)$ have the product representations [14, Theorems 3.1-3.5]

$$\begin{aligned} \mathcal{E}_1(\tau) &= \left(\begin{matrix} q^8, q^9, q^{17}, q^{17} \\ q^2, q^3, q^{14}, q^{15} \end{matrix} ; q^{17} \right)_{\infty}, & \mathcal{E}_2(\tau) &= q \left(\begin{matrix} q^3, q^{14}, q^{17}, q^{17} \\ q, q^5, q^{12}, q^{16} \end{matrix} ; q^{17} \right)_{\infty}, \\ \mathcal{E}_3(\tau) &= q^3 \left(\begin{matrix} q, q^{16}, q^{17}, q^{17} \\ q^4, q^6, q^{11}, q^{13} \end{matrix} ; q^{17} \right)_{\infty}, & \mathcal{E}_4(\tau) &= q \left(\begin{matrix} q^6, q^{11}, q^{17}, q^{17} \\ q^2, q^7, q^{10}, q^{15} \end{matrix} ; q^{17} \right)_{\infty}, \\ \mathcal{E}_5(\tau) &= q^3 \left(\begin{matrix} q^2, q^{15}, q^{17}, q^{17} \\ q^5, q^8, q^9, q^{12} \end{matrix} ; q^{17} \right)_{\infty}, & \mathcal{E}_6(\tau) &= q \left(\begin{matrix} q^5, q^{12}, q^{17}, q^{17} \\ q^3, q^4, q^{13}, q^{14} \end{matrix} ; q^{17} \right)_{\infty}, \\ \mathcal{E}_7(\tau) &= q \left(\begin{matrix} q^4, q^{13}, q^{17}, q^{17} \\ q, q^7, q^{10}, q^{16} \end{matrix} ; q^{17} \right)_{\infty}, & \mathcal{E}_8(\tau) &= q^2 \left(\begin{matrix} q^7, q^{10}, q^{17}, q^{17} \\ q^6, q^8, q^9, q^{11} \end{matrix} ; q^{17} \right)_{\infty}, \end{aligned} \quad (3.3)$$

$$\left(\begin{matrix} a_1, \dots, a_m \\ b_1, \dots, b_n \end{matrix} ; z \right)_{\infty} = \prod_{n=1}^{\infty} \frac{(a_1; z)_{\infty} \cdots (a_m; z)_{\infty}}{(b_1; z)_{\infty} \cdots (b_n; z)_{\infty}}, \quad (a; z)_{\infty} = \prod_{n=0}^{\infty} (1 - az^n).$$

A function Ω is now introduced as a level 17 analogue to the level 5 cusp form UV . Define

$$\Omega(\tau) = \mathcal{E}_1 \mathcal{E}_2 - \mathcal{E}_2 \mathcal{E}_3 + \mathcal{E}_3 \mathcal{E}_4 - \mathcal{E}_4 \mathcal{E}_5 + \mathcal{E}_5 \mathcal{E}_6 - \mathcal{E}_6 \mathcal{E}_7 - \mathcal{E}_7 \mathcal{E}_8 - \mathcal{E}_8 \mathcal{E}_1. \quad (3.4)$$

Proposition 3.1 demonstrates that the weight two parameters z and Ω , respectively, play a role at level seventeen analogous to that played by the parameters Z and UV at level 5.

Proposition 3.1. *Let $z = z(\tau)$ be defined by (1.7). Then*

- (1) *The Eisenstein space of weight two $E_2(\Gamma_0(17))$ is generated by z .*

- (2) The space of cusp forms of weight two $S_2(\Gamma_0(17))$ is generated by Ω .
- (3) Both z and Ω change sign under $|_{W_{17,2}}$, where $f|_{W_{17,k}}(\tau) = 17^{-k/2}\tau^{-k}f(-1/17\tau)$.
- (4) Both z and Ω have zeros at the elliptic points ρ_{\pm} , and in the case of Ω , the zeros are simple.

Proof. From (3.2) and the definition of the \mathcal{E}_i , if arithmetic is performed modulo 8 on the subscripts,

$$\langle 3 \rangle \mathcal{E}_k = \epsilon_k \mathcal{E}_{k+1}, \quad \epsilon_1, \dots, \epsilon_8 = +1, -1, +1, -1, +1, -1, -1, -1. \quad (3.5)$$

This, coupled with the transformation formula for Eisenstein series,

$$\mathcal{E}_k \left(\frac{a\tau + b}{c\tau + d} \right) = (c\tau + d) \cdot \langle a \rangle \mathcal{E}_k(\tau), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(17) \quad (3.6)$$

implies that Ω and z are modular forms of weight two with respect to $\Gamma_0(17)$. From their q -expansions, we deduce that Ω and z are linearly independent over \mathbb{C} . Therefore, from dimension formulas for the respective vector spaces [13], we see that these parameters generate the vector space of weight two forms for $\Gamma_0(17)$. Thus, we obtain the first two claims of Proposition 3.1. The third claim follows from the fact that W_{17} normalizes $\Gamma_0(17)$.

As fundamental domain for $\mathbb{H}/\Gamma_0(17)$ we take $\bigcup_{k=-8}^8 F_k(D) \cup D$, where D is the usual fundamental domain for the full modular group and $F_k(\tau) = \frac{-1}{\tau+k}$. The two elliptic points of order 2 are $\rho_{\pm} = F_{\pm 4}(i)$. Since $\mathbb{H}/\Gamma_0(17)$ has two elliptic points of order 2 and two cusps, the valence formula for a weight k modular form f on $\Gamma_0(17)$ reads as

$$\text{ord}_{\infty} f + \text{ord}_0 f + \frac{\text{ord}_{\rho_+} f}{2} + \frac{\text{ord}_{\rho_-} f}{2} + \sum_{\tau \in \mathbb{H} - \{\rho_{\pm}\}} \text{ord}_{\tau} f = \frac{k}{12} \cdot 18$$

From the q -expansion and the fact that Ω changes sign under $|_{W_{17,2}}$, we know that the two cusps are zeros of Ω , so the valence formula for $f = \Omega$ reads as

$$1 + 1 + \frac{\text{ord}_{\rho_+} f}{2} + \frac{\text{ord}_{\rho_-} f}{2} + \sum_{\tau \in \mathbb{H} - \{\rho_{\pm}\}} \text{ord}_{\tau} f = 3.$$

Since Ω changes sign under $|_{W_{17,2}}$ and the fixed point of W_{17} is not a zero (as one may check numerically), the zeros must come in pairs. Accordingly, the two other zeros must be the two elliptic points, and these are simple zeros. A similar argument gives the result for z . \square

The cusp form and Eisenstein series from Proposition 3.1 can now be used in the construction of a $\Gamma_0(17)$ invariant function of the same form as T, \mathcal{T} given by (2.7). Although the representation for $x(\tau)$ given here appears to differ from that given in the introduction, we ultimately demonstrate agreement of the two representations in Proposition 3.3. The parameters $x(\tau)$ and $w(\tau)$, defined in Proposition 3.2, play roles analogous to corresponding parameters T and W in Proposition 2.2.

Proposition 3.2. *If the Fricke involution is denoted $W_{17} = W_{17,0}$ and x and w are defined by*

$$x(\tau) = \frac{\Omega}{z}, \quad w(\tau) = \frac{2}{z} \theta_q \log x, \quad (3.7)$$

- (1) x is invariant under $\Gamma_0(17)$ as well as W_{17} ; and
- (2) x has two simple zeros on $\mathbb{H}/\Gamma_0(17)$ at the two cusps.
- (3) The field of functions invariant under $\Gamma_0(17)$ and W_{17} is $A_0(\langle \Gamma_0(17) \rangle) = \mathbb{C}(x)$.
- (4) The field of functions invariant under $\Gamma_0(17)$ is given by $A_0(\Gamma_0(17)) = \mathbb{C}(x, w)$.
- (5) The relation $w^2 = -127x^4 - 48x^3 - 66x^2 - 16x + 1$ holds.

Proof. The first two assertions follow directly from Proposition 3.1. The third assertion is then a direct consequence of the first two. For the fourth assertion, the functions $x(\tau)$ and $w(\tau)$ are invariant under $\Gamma_0(17)$, so it suffices to show that they generate the whole field. Since x has order 2, we have $[A_0(\Gamma_0(17)) : \mathbb{C}(x)] = 2$. Since $w \notin \mathbb{C}(x)$ because it changes sign under W_{17} , we must have $[A_0(\Gamma_0(17)) : \mathbb{C}(x, w)] = 1$, that is, the second assertion holds. For the final assertion, the function w^2 is fixed under W_{17} and has the same set of poles as x , hence it is a polynomial in x . We bound the degree of this polynomial by 4 and find its coefficients by comparing q -expansions. \square

The parameter x is expressible as the rational function of r and s appearing in the Introduction and in terms of the McKay-Thompson series 17A [8, Table 4A].

Proposition 3.3. *Define $\eta(\tau) = q^{1/24}(q; q)_\infty$, and let x be defined as in Proposition 3.2. Then*

$$x = \frac{r(r^2s + 8rs - r - s)}{8r^3s - 3r^2s + r - s}, \quad (3.8)$$

$$\frac{1-x}{2x} = \frac{1}{4\eta(\tau)^2\eta(17\tau)^2} \left(\sum_{m,n=-\infty}^{\infty} (e^{\pi im} - e^{\pi in}) q^{\frac{1}{4}n^2 + \frac{17}{4}m^2} \right)^2. \quad (3.9)$$

Proof. From the product representation for r and those for the Eisenstein sums \mathcal{E}_i , from (3.3)

$$r = \frac{\mathcal{E}_1\mathcal{E}_3\mathcal{E}_5\mathcal{E}_7}{\mathcal{E}_2\mathcal{E}_4\mathcal{E}_6\mathcal{E}_8}. \quad (3.10)$$

Therefore, r is the quotient of weight four modular forms for $\Gamma_1(17)$, and $x = \Omega/z$ is the quotient of weight two modular forms for $\Gamma_1(17)$. Hence, the quadratic relation between x and r ,

$$\frac{4}{r} - 4r - 15 = \frac{(xr - 1)^2(4r - 1)^2}{(x + r)^2} \quad (3.11)$$

may be transcribed as a relation between modular forms of weight 20 for $\Gamma_1(17)$ and proved from the Sturm bound by verifying the q -expansion to order $481 = 1 + 20 \cdot 288/12$. Then

$$\sqrt{\frac{4}{r} - 4r - 15} = \frac{(xr - 1)(4r - 1)}{(x + r)}, \quad (3.12)$$

where the branch of the square root is determined using the definition of x and r . Therefore,

$$x = \frac{4r - 1 + r\beta(r)}{4r^2 - r - \beta(r)}, \quad \beta(r) = \sqrt{\frac{4}{r} - 4r - 15}. \quad (3.13)$$

The first equation of (3.13) is seen to be equivalent to (3.8) by applying (2.13). Equation (3.9) may be derived from respective q -expansions since each side is a Hauptmodul for $\Gamma_0(17)+$. \square

It follows from the first part of Proposition 3.2 and Theorem 2.1 that z satisfies a third order linear homogeneous differential equation with coefficients in $\mathbb{C}(x)$. In order to formulate the differential equation, we state the following preliminary nonlinear differential equation in terms of the differential operator $\theta_q := q \frac{d}{dq}$. This is written even more succinctly as $f_q := \theta_q f$.

Lemma 3.4.

$$\frac{2zz_{qq} - 3z_q^2}{3z^4} = \frac{x(127x^5 - 222x^4 + 126x^3 + 4x^2 + 27x + 2)}{4(x-1)^2}$$

Proof. Let $f(\tau)$ denote the function on the left hand side of the proposed equality. If z satisfies the functional equation

$$z\left(\frac{a\tau + b}{c\tau + d}\right) = \epsilon \frac{(c\tau + d)^2}{ad - bc} z(\tau)$$

one can compute that

$$\frac{2zz_{qq} - 3z_q^2}{3z^4} \left(\frac{a\tau + b}{c\tau + d}\right) = \frac{1}{\epsilon^2} \frac{2zz_{qq} - 3z_q^2}{3z^4}(\tau).$$

By Proposition 3.1, we have $\epsilon = 1$ for elements of $\Gamma_0(17)$ and $\epsilon = -1$ for W_{17} . Thus we see that $f(\tau)$ is invariant under $\Gamma_0(17)$ and W_{17} in weight 0. According to Theorem 4.4, we see that x does not have a pole at the two elliptic points, i.e. $x(\rho_{\pm}) = 1$. This means that the two zeros of z at these elliptic points are both simple. Hence, z has two other simple zeros p_1 and $p_2 = W_{17}(p_1)$, which are also the poles of x , modulo $\Gamma_0(17)$, as observed in the proof of Proposition 3.1. Since all of the poles of z are simple, we can take the expansion

$$z(\tau) = c(\tau - r) + \dots$$

at the zeros $r = \rho_+, \rho_-, p_1, p_2$, where c is non-zero. Each of these zeros contributes a quadruple pole to $f(\tau)$ since

$$\frac{2zz_{qq} - 3z_q^2}{3z^4}(\tau) = \frac{3}{(2\pi c)^2(\tau - r)^4} + \dots$$

In the fundamental domain of $\mathbb{H}/\Gamma_0(17)$, the translate $F_4(D)$ is adjacent to itself. Thus $x(\tau)$ must identify the two halves of the corresponding side of $F_4(D)$ (the side that contains $F_4(i)$). Likewise for F_{-4} . Therefore, at the elliptic point ρ_{\pm} , the function $x(\tau)$ is locally a holomorphic function of $((\tau - \rho_{\pm})/(\tau - \rho_{\pm}^*))^2$ so that

$$x(\tau) = 1 + c_{\pm}(\tau - \rho_{\pm})^2 + \dots$$

We see now that $(x - 1)^2 f$ has poles only at p_1 and p_2 , each of order six. It is therefore a polynomial of degree six in x , and we can compute that

$$4(x - 1)^2 f - x(127x^5 - 222x^4 + 126x^3 + 4x^2 + 27x + 2) = O(q^7).$$

The left hand side has poles of order 6 at p_1, p_2 and zeros at least order 7 at 0 and ∞ . This contradicts the valence formula unless the left hand side is constant. \square

We now give the third order linear differential equation for z with rational coefficients in x . The concise formulation of the differential equation in (3.14) is motivated by the general form of such differential equations from [22, 24].

Theorem 3.5. *With respect to the function x , the form $f = z$ satisfies the differential equation.*

$$\begin{aligned} 0 = & 3x(254x^6 - 714x^5 + 681x^4 - 250x^3 - 6x^2 - 28x - 1)f \\ & + x(x - 1)(1397x^5 - 2482x^4 + 1094x^3 - 28x^2 + 197x + 14)f_x \\ & + 6x(x - 1)^3(127x^3 + 36x^2 + 33x + 4)f_{xx} \\ & + (x - 1)^3(127x^4 + 48x^3 + 66x^2 + 16x - 1)f_{xxx}. \end{aligned}$$

Proof. The differential equation satisfied by $f = z$ is given as

$$\det \begin{pmatrix} f & f_x & f_{xx} & f_{xxx} \\ (z) & (z)_x & (z)_{xx} & (z)_{xxx} \\ (z \log q) & (z \log q)_x & (z \log q)_{xx} & (z \log q)_{xxx} \\ (z \log^2 q) & (z \log^2 q)_x & (z \log^2 q)_{xx} & (z \log^2 q)_{xxx} \end{pmatrix} = 0. \quad (3.14)$$

When expanding this determinant, we make the following substitutions:

- (1) For the differential with respect to x , use the definition (3.7) in the form

$$\theta_x = x \frac{\partial}{\partial x} = \frac{2}{wz} \theta_q.$$

- (2) When the first derivative x_q appears, use the definition (3.7) in the form

$$x_q = \frac{1}{2} x w z.$$

- (3) When the first derivative w_q appears, use the relation between w and x to obtain

$$w_q = -x(127x^3 + 36x^2 + 33x + 4)z.$$

- (4) When the second derivative z_{qq} appears, use Lemma 3.4 in the form

$$z_{qq} = \frac{3z_q^2}{2z} + \frac{3x(127x^5 - 222x^4 + 126x^3 + 4x^2 + 27x + 2)}{8(x-1)^2} z^3$$

When these substitutions are made in (3.14), the claimed differential equation results after clearing denominators by multiplying by $(1-x)^3 w^5/16$ and using Proposition 3.2 (5). \square

The linear differential equation in Theorem 3.5 induces a series expansion for z in terms of x with coefficients A_n .

Corollary 3.6.

$$z = \sum_{n=0}^{\infty} A_n x^n \quad |x| < 0.05122\dots, \quad (3.15)$$

where $A_0 = 2$, $A_{-1, \dots, -6} = 0$ and

$$\begin{aligned} 0 = & (n+1)^3 A_{n+1} + (-19n^3 - 24n^2 - 14n - 3) A_n \\ & - 3(5n^3 + 27n^2 - 8n + 4) A_{n-1} + (101n^3 - 300n^2 + 213n - 52) A_{n-2} \\ & - 3(55n^3 - 267n^2 + 491n - 305) A_{n-3} + 3(n-3)(101n^2 - 297n + 253) A_{n-4} \\ & - 9(n-4)(n-3)(37n - 66) A_{n-5} + 127(n-5)(n-4)(n-3) A_{n-6}. \end{aligned}$$

The radius of convergence is the positive root of $127x^4 + 48x^3 + 66x^2 + 16x - 1$.

To make use of the series appearing in Corollary 3.6, we require explicit evaluations for the $x(\tau)$ within the domain of validity. In the next section, we prove that the number of singular values is finite and compile a complete list of quadratic evaluations and expansions.

4. SINGULAR VALUES AND SERIES FOR $1/\pi$

In this Section, singular values for $x(\tau)$ are derived and used to formulate Ramanujan-Sato expansions. The work culminates in a proof that there are precisely 11 singular values for $x(\tau)$ of degree at most two over \mathbb{Q} within the radius of convergence of Corollary 3.6. The series given by (1.6) is the only such expansion with a rational singular value for $x(\tau)$. The main challenge in proving the expansions lies in rigorously determining exact evaluations for $x(\tau)$ for given τ and deriving constants appearing in the Ramanujan-Sato series. To do this, we formulate modular equations for $x(\tau)$ and provide an explicit relation between the modular equations and constants appearing in the series.

We demonstrate in the proof of Theorem 4.7 that the following table is a complete list of singular values for $x(\tau)$ in a fundamental domain for $\Gamma_0(17)$ with $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$ within the radius of convergence of Corollary 3.6. Each value τ is listed by the coefficients (a, b, c) of its minimal polynomial, and the values are ordered by discriminant.

$b^2 - 4ac$	$\tau(a, b, c)$	$x(\tau)$	
-1411	(17, 17, 25)	$(-1025 - 252\sqrt{17})^{-1}$	
-1003	(17, 17, 19)	$(-345 - 84\sqrt{17})^{-1}$	
-595	(17, -17, 13)	$(-90 - 21\sqrt{17})^{-1}$	
-427	(17, -27, 17)	$(30 + 33i\sqrt{7})^{-1}$	
-427	(17, -41, 31)	$(30 - 33i\sqrt{7})^{-1}$	
-408	(17, -34, 23)	$(55 + 24\sqrt{2})^{-1}$	(4.1)
-408	(34, -68, 37)	$(55 - 24\sqrt{2})^{-1}$	
-340	(17, -34, 22)	$(29 + 4\sqrt{85})^{-1}$	
-323	(17, -17, 9)	$(-22 - 7\sqrt{17})^{-1}$	
-187	(17, -17, 7)	$-1/21$	
-136	(17, -34, 19)	$(12 + 3\sqrt{17})^{-1}$	

TABLE 1. Complete list of singular values of $x(\tau)$ of degree at most 2 within the radius of convergence of Corollary 3.6, ordered by discriminant.

By Proposition 3.3, finding singular values for $x(\tau)$ is equivalent to finding singular values for the normalized Thompson series 17A. The fundamental results needed for such evaluations are presented in [7]. Our work below is a detailed rendition of the general presentation in [7] tailored to the modular function $x(\tau)$. We begin with the set of matrices

$$\Delta_n^*(17) = \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathbb{Z}^{2 \times 2} \mid \gcd(\alpha, \beta, \gamma, \delta) = 1 \text{ and } \alpha\delta - \beta\gamma = n \text{ and } \gamma \equiv 0 \pmod{17} \right\}.$$

Lemma 4.1. *If $\gcd(n, 17) = 1$, then $\Delta_n^*(17)$ has the coset decomposition*

$$\Delta_n^*(17) = \bigcup_{\substack{\alpha\delta=n \\ 0 \leq \beta < \delta \\ \gcd(\alpha, \beta, \delta)=1}} \Gamma_0(17) \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix},$$

and the double coset representation

$$\Delta_n^*(17) = \Gamma_0(17) \begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} \Gamma_0(17).$$

Proof. Any $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Delta_n^*$ can be converted to an upper triangular matrix by multiplying on the left by a matrix of the form

$$\left(\begin{array}{cc} * & * \\ \frac{\gamma}{\gcd(\alpha, \gamma)} & \frac{-\alpha}{\gcd(\alpha, \gamma)} \end{array} \right) \in \Gamma_0(17).$$

It is then easy to see that the claimed representatives are distinct modulo $\Gamma_0(17)$. This proves the decomposition formula. Next, by performing elementary row and column operations on the matrix $m \in \Delta_n^*(17)$, we find matrices $\gamma_1, \gamma_2 \in \Gamma(1)$ such that $m = \gamma_1 \begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} \gamma_2$. Since

$$\begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} = \begin{pmatrix} a & b \\ nc & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} \begin{pmatrix} a & -bn \\ -c & d \end{pmatrix},$$

we may find appropriate a, b, c , and d such that $\gamma'_1 = \gamma_1 \begin{pmatrix} a & b \\ nc & d \end{pmatrix} \in \Gamma_0(17)$. This results in an equality of the form $m = \gamma'_1 \begin{pmatrix} 1 & 0 \\ 0 & n \end{pmatrix} \gamma'_2$ where $\gamma'_1 \in \Gamma_0(17)$, which forces $\gamma'_2 \in \Gamma_0(17)$ as well. This establishes the double coset representation. \square

We now establish modular equations central to our explicit evaluations for $x(\tau)$.

Proposition 4.2. *For any integer $n \geq 2$ with $\gcd(n, 17) = 1$, there is a polynomial $\Psi_n(X, Y)$ of degree $\psi(n) = n \prod_{\substack{q|n \\ q \text{ prime}}} (1 + \frac{1}{q})$ in X and Y such that:*

- (1) $\Psi_n(X, Y)$ is irreducible and has degree $\psi(n)$ in X and Y .
- (2) $\Psi_n(X, Y)$ is symmetric in X and Y .
- (3) The roots of $\Psi_n(x(\tau), Y) = 0$ are precisely the numbers $Y = x((\alpha\tau + \beta)/\delta)$ for integers α, β and δ such that $\alpha\delta = n$, $0 \leq \beta < \delta$, and $\gcd(\alpha, \beta, \delta) = 1$.

Proof. The polynomial Ψ_n satisfies

$$(XY)^{-\psi(n)} \Psi_n(X, Y) = \prod_{\substack{\alpha\delta=n \\ 0 \leq \beta < \delta \\ (\alpha, \beta, \delta)=1}} \left(Y^{-1} - x \left(\frac{\alpha\tau + \beta}{\delta} \right)^{-1} \right), \quad (4.2)$$

where the coefficients of Y^{-k} on the right hand side should be expressed as polynomials in $1/X$ for $X = x(\tau)$ as demonstrated in the proof of Corollary 4.3. This relies on the fact that $\Gamma_0(17)$ and W_{17} permute the set of functions $x((\alpha\tau + \beta)/\delta)$ where $\alpha\delta = n$, $0 \leq \beta < \delta$, and $\gcd(\alpha, \beta, \delta) = 1$. The double coset representation in Lemma 4.1 shows that every orbit contains $x(\tau/n)$, and hence the action of $\Gamma_0(17)$ on the roots must be transitive. Since

$$\begin{pmatrix} 0 & -1 \\ 17 & 0 \end{pmatrix}^{-1} \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 17 & 0 \end{pmatrix} = \begin{pmatrix} \delta & 0 \\ -17\beta & \alpha \end{pmatrix} \in \Delta_n^*(17),$$

it is clear that W_{17} permutes these functions as well by the decomposition in Lemma 4.1. The coefficient of $X^{\psi(n)}Y^{\psi(n)}$ in $\Psi_n(X, Y)$ is the constant term of the product on the right hand side of (4.2), which is clearly non-zero because the function $x(\tau)$ does not have poles at the cusps of $\mathbb{H}/\Gamma_0(17)$. Therefore, $\Psi_n(X, X)$ has the claimed degree $2\psi(n)$. The symmetry can be proven by noting that $\tau \rightarrow -1/(17n\tau)$ interchanges $x(\tau)$ and $x(n\tau)$. \square

In Corollary 4.3, modular equations $\Psi_n(X, Y) = 0$ are derived for $X = x(\tau)$ and $y = x((\alpha\tau + \beta)/\delta)$ satisfying the conditions of Proposition 4.2. The proof indicates how modular equations for larger n may be derived and involves techniques analogous to those used to deduce classical modular equations of level n satisfied by the j invariant [16, 21].

Corollary 4.3. *We have*

$$\begin{aligned} \Psi_2(X, Y) &= -9X^3Y^3 - 12X^3Y^2 + X^3Y + 2X^3 - 12X^2Y^3 \\ &\quad + 8X^2Y^2 + 10X^2Y + XY^3 + 10XY^2 - XY + 2Y^3, \\ \Psi_3(X, Y) &= 435X^4Y^4 + 231X^4Y^3 + 231X^3Y^4 + 45X^4Y^2 - 385X^3Y^3 + 45X^2Y^4 \\ &\quad - 39X^4Y - 63X^3Y^2 - 63X^2Y^3 - 39XY^4 + 4X^4 + 9X^3Y + 123X^2Y^2 + 9XY^3 \\ &\quad + 4Y^4 + 15X^2Y + 15XY^2 - XY. \end{aligned}$$

Proof. The level $n = 2$ result is representative of $n = 3$ and other cases. For $n = 2$, we have

$$\{(\alpha, \beta, \delta) \mid \gcd(\alpha, \beta, \delta) = 1, 0 \leq \beta < \delta, \alpha\delta = n\} = \{(1, 0, 2), (2, 0, 1), (1, 1, 2)\}.$$

Let $x(\tau)$ be defined as in Proposition 3.2. Then

$$x_1 = x(\tau/2), \quad x_2 = x(2\tau), \quad x_3 = x\left(\frac{\tau+1}{2}\right).$$

By (4.2),

$$\Psi_2(1/X, 1/Y) = Y^{-3} - \left(\frac{x_1x_2 + x_1x_3 + x_2x_3}{x_1x_2x_3} \right) Y^{-2} + \left(\frac{x_1 + x_2 + x_3}{x_1x_2x_3} \right) Y^{-1} - \frac{1}{x_1x_2x_3}.$$

By Theorem 3.2, we know that $1/x(\tau)$ is analytic on $X_0(17)$ except for simple poles at the cusps. Therefore, the only poles in $\mathbb{H}/\Gamma_0(17)$ of the coefficients of Y^{-k} are at points equivalent to the cusps 0 and ∞ . We can explicitly compute the q -expansion for each of the coefficients and deduce that each has a pole of order at most 3 at $q = 0$. Since each coefficient is invariant under $\Gamma_0(17)$ and W_{17} , the coefficients may be expressed as polynomials of degree at most 3 in $1/x(\tau)$. Explicitly,

$$\begin{aligned} -\frac{x_1x_2 + x_1x_3 + x_2x_3}{x_1x_2x_3} &= -\frac{2}{q^2} - 15 + O(q), \\ \frac{x_1 + x_2 + x_3}{x_1x_2x_3} &= \frac{20}{q^2} + \frac{108}{q} + 419 + O(q), \\ -\frac{1}{x_1x_2x_3} &= \frac{8}{q^3} + \frac{62}{q^2} + \frac{316}{q} + 1307 + O(q). \end{aligned}$$

Therefore, with $x = x(\tau)$, we may determine polynomials in $1/x$ such that

$$\begin{aligned} c_1(\tau) &= -\frac{1}{x_1x_2x_3} - \left(-\frac{9}{2} - 6x^{-1} + \frac{1}{2}x^{-2} + x^{-3} \right) = O(q), \\ c_2(\tau) &= \frac{x_1 + x_2 + x_3}{x_1x_2x_3} - (-6 + 4x^{-1} + 5x^{-2}) = O(q), \\ c_3(\tau) &= -\frac{x_1x_2 + x_1x_3 + x_2x_3}{x_1x_2x_3} - \left(\frac{1}{2} + 5x^{-1} - \frac{1}{2}x^{-2} \right) = O(q). \end{aligned}$$

Since the functions $c_j(\tau)$, $j = 1, 2, 3$, are analytic on the upper half plane and at the cusps of $X_0(17)$, each c_j is constant, and $c_j(i\infty) = 0$. \square

In Theorem 4.4, we prove each evaluation from Table 1. The evaluations are proven by showing that $x(\tau)$ satisfies a modular equation of degree n for some n .

Theorem 4.4. *Each evaluation for $x(\tau(a, b, c))$ from Table 1 holds, and $x(\tau(17, -8, 1)) = 1$.*

Proof. First note that

$$\tau(17, -34 - 19) = \frac{1}{17} \left(17 + i\sqrt{34} \right).$$

Therefore,

$$x(\tau) = x\left(i\sqrt{\frac{2}{17}}\right)$$

Observe that with

$$\tau = i\sqrt{\frac{2}{17}}, \quad -\frac{1}{17\tau} = \frac{\tau}{2}.$$

Since $x(\tau)$ is invariant under the Fricke involution W_{17} , we have $x(\tau) = x(\tau/2)$. That is, $X = Y$ in the degree 2 modular equation above. Setting $Y = X$ and simplifying the equation, we get

$$X^2(X - 1)(X + 1)(9X^2 + 24X - 1) = 0. \quad (4.3)$$

Now that we have proven $x(\tau)$ satisfies (4.3), we may numerically deduce $x(\tau)$ is a root of $9X^2 + 24X - 1$, and

$$x\left(i\sqrt{\frac{2}{17}}\right) = -\frac{4}{3} + \frac{1}{3}\sqrt{17}.$$

We may similarly prove $x(\tau(17, -8, 1)) = 1$ and the remaining evaluations in Table 1 from modular equations of degree n if we can determine, for each given value of $\tau = \tau(a, b, c)$, an upper triangular matrix $(\alpha, \beta; 0, \gamma)$ such that $x(\tau) = x((\alpha, \beta; 0, \gamma)\tau)$ and $\alpha\delta = n$, $0 \leq \beta < \delta$, $\gcd(\alpha, \beta, \delta) = 1$, with $\gcd(n, 17) = 1$. For each τ , Table 2 provides a $\gamma \in \Delta_n^*(17)$ such that $\gamma\tau = \tau$ or $W_{17}\tau$ and a $\Gamma_0(17)$ equivalent upper triangular matrix $(\alpha, \beta; 0, \gamma)$. \square

$\tau(a, b, c)$	Element of $\Delta_n^*(17)$	$(\alpha, \beta; 0, \gamma)$	
(17, 17, 25)	(-1, -2; 17, -25)	(1, 2; 0, 59)	
(17, 17, 19)	(1, 0; 17, 19)	(1, 0; 0, 19)	
(17, -17, 13)	(-1, 0; 17, -13)	(1, 0; 0, 13)	
(17, -27, 17)	(13, -17; 17, -14) [†]	(1, 81; 0, 107)	
(17, -41, 31)	(20, -31; 17, -21) [†]	(1, 68; 0, 107)	
(17, -34, 23)	(-1, 0; 34, -23)	(1, 0; 0, 23)	(4.4)
(34, -68, 37)	(-2, 1; 51, -37)	(1, 11; 0, 23)	
(17, -34, 22)	(-1, 1; 17, -22)	(1, 4; 0, 5)	
(17, -17, 9)	(-2, 1; 17, -18)	(1, 9; 0, 19)	
(17, -17, 7)	(-1, 0; 17, -7)	(1, 0; 0, 7)	
(17, -34, 19)	(-1, 1; 17, -19)	(1, 1; 0, 2)	
(17, -8, 1)	(3, -1; 17, -5)	(1, 1; 0, 2)	

TABLE 2. Elements of $\Delta_n^*(17)$ mapping τ to its image under W_{17} or fixing[†] τ and a corresponding $\Gamma_0(17)$ equivalent upper triangular matrix $(\alpha, \beta; 0, \gamma)$.

For values $x(\tau)$ in the domain of validity for series from Corollary 3.6, we may construct Ramanujan-Sato expansions via Theorem 4.5, a specialization of [4, Theorem 2.1].

Theorem 4.5 (Series for $1/\pi$). *Suppose there is a matrix $(a, b; c, d) \in \langle \Gamma_0(17), W_{17} \rangle$ such that*

$$\frac{a\tau + b}{c\tau + d} = \frac{\alpha\tau + \beta}{\delta}$$

for $\alpha\delta = n$ and $0 \leq \beta < \delta$. Set $X = x(\tau)$, which is determined from $\Psi_n(X, X) = 0$, and further set

$$W = w(\tau), \quad \Psi_X = \frac{\partial \Psi_p}{\partial X}(X, X), \quad \Psi_Y = \frac{\partial \Psi_p}{\partial Y}(X, X),$$

and let $\epsilon \in \mathbb{Q}$ and $\eta = \pm 1$ satisfy for all τ

$$z\left(\frac{a\tau + b}{c\tau + d}\right) = \epsilon(c\tau + d)^2 z(\tau), \quad w\left(\frac{a\tau + b}{c\tau + d}\right) = \eta w(\tau).$$

If A_k is the sequence defined in Corollary 3.6 and

$$B = -\frac{iW(\delta^2 \Psi_X(ad - bc) + \alpha^2 \eta \epsilon \Psi_Y(c\tau + d)^4)}{2\alpha^2 c \eta \epsilon \Psi_Y(c\tau + d)^3},$$

$$C = \frac{i\delta^2(bc - ad)W}{2\alpha^2 c \eta \epsilon \Psi_Y^3(c\tau + d)^3} (\Psi_X \Psi_Y (\Psi_X + \Psi_Y) (1 + \theta_X \log W) \\ + (\Psi_X^2 \Psi_{YY} - 2\Psi_X \Psi_{XY} \Psi_Y + \Psi_{XX} \Psi_Y^2) X),$$

then

$$\frac{1}{\pi} = \sum_{k=0}^{\infty} A_k (Bk + C) X^k.$$

Proof. Differentiate the relation

$$z\left(\frac{a\tau + b}{c\tau + d}\right) = \epsilon(c\tau + d)^2 z(\tau)$$

once and the relation

$$\Psi_n\left(x(\tau), x\left(\frac{\alpha\tau + \beta}{\delta}\right)\right) = 0$$

twice and then set τ to the value in the hypothesis of the theorem. \square

Corollary 4.6. *If A_k is the sequence defined in Corollary 3.6,*

$$\begin{aligned} \frac{\sqrt{11}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{307 + 748k}{(-21)^{k+2}}, \\ \frac{2\sqrt{154\sqrt{17} - 634}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{1779 - 195\sqrt{17} + 3040k}{(-22 - 7\sqrt{17})^{k+2}}, \\ \frac{214\sqrt{119} - 882\sqrt{7}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{9241 - 1047\sqrt{17} + 21280k}{(-90 - 21\sqrt{17})^{k+2}}, \\ \frac{\sqrt{1041894\sqrt{17} - 4295839}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{71065 - 15096\sqrt{17} + 50740k}{(-345 - 84\sqrt{17})^{k+2}}, \\ \frac{9\sqrt{2038550094\sqrt{17} - 8405157343}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{74004567 - 11655082\sqrt{17} + 178775028k}{(-1025 - 252\sqrt{17})^{k+2}}, \\ \frac{\sqrt{14(1267990301 \mp 85084065i\sqrt{7})}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{3370317797 \pm 95119383i\sqrt{7} + 12974719520k}{161874(30 \pm 33i\sqrt{7})^k}, \end{aligned}$$

and

$$\begin{aligned} \frac{\sqrt{9\sqrt{17} - 37}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{32 - 3\sqrt{17} + 32k}{(12 + 3\sqrt{17})^{k+2}}, \\ \frac{261\sqrt{5} - 135\sqrt{17}}{\pi} &= \sum_{k=0}^{\infty} A_k \frac{21500 - 788\sqrt{85} + 54720k}{(29 + 4\sqrt{85})^{k+2}}, \\ \frac{539\sqrt{6} \mp 735\sqrt{3}}{\pi} 2^{(1\mp 1)/2} &= \sum_{k=0}^{\infty} A_k \frac{58962 \mp 7226\sqrt{2} + 199920k}{(55 \pm 24\sqrt{2})^{k+2}}. \end{aligned}$$

Proof. The first five series may be derived by setting $\tau = \frac{1}{2} + \frac{1}{2}\sqrt{\frac{n}{17}}i$ and using $\frac{17\tau-9}{34\tau-17} = \frac{\tau+(n-1)/2}{n}$ with $\epsilon = -1/17$ and $\eta = -1$ in Theorem 4.5 for the values $n = 11, 19, 35, 59, 83$. The subsequent pair may be derived from Theorem 4.5 by setting $\tau = (\pm 7 + \sqrt{427i})/34$, using $(11\tau - 2)/(17\tau - 3) = (\tau + 69)/107$ and $(13\tau + 3)/(17\tau + 4) = (\tau + 82)/107$, respectively. The next three arise from setting $\tau = \sqrt{\frac{n}{17}}i$ and using $\frac{-1}{17\tau} = \frac{\tau}{n}$ for $n = 2, 5, 6$. The final series comes from setting $\tau = \sqrt{3/34}i$ and using $\frac{-1}{17\tau} = \frac{2\tau}{3}$. \square

Theorem 4.7. *There are precisely eleven $\Gamma_0(17)$ inequivalent algebraic τ in the upper half plane such that $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$ with $x(\tau)$ in the radius of convergence of Corollary 3.6.*

Proof. We formulate a complete list of algebraic τ such that $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$ using well known facts about the j invariant [20]. First, for algebraic τ , the only algebraic values of $j(\tau)$ occur at $\Im\tau > 0$ satisfying $a\tau^2 + b\tau + c = 0$ for $a, b, c \in \mathbb{Z}$, with $d = b^2 - 4ac < 0$. Moreover, $[\mathbb{Q}(j(\tau)) : \mathbb{Q}] = h(d)$, where $h(d)$ is the class number. Since there is a polynomial relation $P(x, j)$ between x and j of degree 2 [7, Remark 1.5.3], we have $[\mathbb{Q}(j(\tau)) : \mathbb{Q}] \leq 2[\mathbb{Q}(x(\tau)) : \mathbb{Q}]$, and so values τ with $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$ satisfy $[\mathbb{Q}(j(\tau)) : \mathbb{Q}] = h(d) \leq 4$. Therefore, the bound $|d| \leq 1555$ for $h(d) \leq 4$ from [23] implies that the following algorithm results in a complete list of algebraic $(\tau, x(\tau))$ with $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$:

For each discriminant $-1555 \leq d \leq -1$,

- (1) List all primitive reduced $\tau = \tau(a, b, c)$ of discriminant d in a fundamental domain for $PSL_2(\mathbb{Z})$. Translate these values via a set of coset representatives for $\Gamma_0(17)$ to a fundamental domain for $\Gamma_0(17)$.

(2) Factor the resultant of $P(X, Y)$ and the class polynomial

$$H_d(Y) = \prod_{\substack{(a,b,c) \text{ reduced, primitive} \\ d=b^2-4ac}} \left(Y - j\left(\frac{-b + \sqrt{d}}{2a}\right) \right).$$

The linear and quadratic factors of the resultant correspond to a complete list of $x = x(\tau)$, for τ of discriminant d , such that $[\mathbb{Q}(x(\tau)) : \mathbb{Q}] \leq 2$. Associate candidate values τ from Step 1 to x by numerically approximating $x(\tau)$. For each tentative pair, (τ, x) , prove the evaluation $x = x(\tau)$ as indicated in the proof of Theorem 4.4.

The algorithm is easy to implement. The resulting values of $\tau(a, b, c)$ with $x(\tau)$ within the radius of convergence of Corollary 3.6 are given in Table 1. \square

REFERENCES

- [1] B. C. Berndt. *Ramanujan's notebooks. Part III*. Springer-Verlag, New York, 1991.
- [2] B. C. Berndt and R. A. Rankin. *Ramanujan*, volume 9 of *History of Mathematics*. American Mathematical Society, Providence, RI, 1995. Letters and commentary.
- [3] J. M. Borwein and P. B. Borwein. Ramanujan's rational and algebraic series for $1/\pi$. *J. Indian Math. Soc. (N.S.)*, 51:147–160 (1988), 1987.
- [4] H. H. Chan, S. H. Chan, and Z.-G. Liu. Domb's numbers and Ramanujan-Sato type series for $1/\pi$. *Adv. Math.*, 186(2):396–410, 2004.
- [5] H. H. Chan and S. Cooper. Rational analogues of Ramanujan's series for $1/\pi$. *Math. Proc. Cambridge Philos. Soc.*, 153(2):361–383, 2012.
- [6] H. H. Chan, Y. Tanigawa, Y. Yang, and W. Zudilin. New analogues of Clausen's identities arising from the theory of modular forms. *Adv. Math.*, 228(2):1294–1314, 2011.
- [7] I. Chen and N. Yui. Singular values of Thompson series. In *Groups, difference sets, and the Monster (Columbus, OH, 1993)*, volume 4 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 255–326. de Gruyter, Berlin, 1996.
- [8] J. H. Conway and S. P. Norton. Monstrous moonshine. *Bull. London Math. Soc.*, 11(3):308–339, 1979.
- [9] S. Cooper. Sporadic sequences, modular forms and new series for $1/\pi$. *Ramanujan J.*, 29(1-3):163–183, 2012.
- [10] S. Cooper, J. Ge, and D. Ye. Hypergeometric transformation formulas of degrees 3, 7, 11 and 23. *J. Math. Anal. Appl.*, 421(2):1358–1376, 2015.
- [11] S. Cooper and D. Ye. Explicit evaluations of a level 13 analogue of the Rogers-Ramanujan continued fraction. *J. Number Theory*, 139:91–111, 2014.
- [12] S. Cooper and D. Ye. The Rogers-Ramanujan continued fraction and its level 13 analogue. *J. Approx. Theory*, 193:99–127, 2015.
- [13] F. Diamond and J. Shurman. *A first course in modular forms*, volume 228 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.
- [14] T. Huber, D. Lara, and E. Melendez. Balanced modular parameterizations. *arXiv:1405.6761 [math.NT]*, Preprint.
- [15] T. Huber and D. Schultz. Generalized reciprocal identities. *Proc. Amer. Math. Soc.*, To Appear.
- [16] S. Lang. *Elliptic functions*, volume 112 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1987. With an appendix by J. Tate.
- [17] S. Ramanujan. *Notebooks. Vols. 1, 2*. Tata Institute of Fundamental Research, Bombay, 1957.
- [18] S. Ramanujan. Modular equations and approximations to π [Quart. J. Math. **45** (1914), 350–372]. In *Collected papers of Srinivasa Ramanujan*, pages 23–39. AMS Chelsea Publ., Providence, RI, 2000.
- [19] L. J. Rogers. Second Memoir on the Expansion of certain Infinite Products. *Proc. London Math. Soc.*, S1-25(1):318.
- [20] T. Schneider. Arithmetische Untersuchungen elliptischer Integrale. *Math. Ann.*, 113(1):1–13, 1937.
- [21] G. Shimura. *Introduction to the arithmetic theory of automorphic functions*. Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten, Publishers, Tokyo; Princeton University Press, Princeton, N.J., 1971. Kanô Memorial Lectures, No. 1.
- [22] P. Stiller. Special values of Dirichlet series, monodromy, and the periods of automorphic forms. *Mem. Amer. Math. Soc.*, 49(299):iv+116, 1984.
- [23] M. Watkins. Class numbers of imaginary quadratic fields. *Math. Comp.*, 73(246):907–938 (electronic), 2004.
- [24] Y. Yang. On differential equations satisfied by modular forms. *Math. Z.*, 246(1-2):1–19, 2004.