

4-1-2010

An Expeditious Synthesis of N-Substituted Pyrroles via Microwave-Induced Iodine-Catalyzed Reactions Under Solventless Conditions

Debasish Bandyopadhyay

The University of Texas Rio Grande Valley, debasish.bandyopadhyay@utrgv.edu

Sanghamitra Mukherjee

Bimal K. Banik

Follow this and additional works at: https://scholarworks.utrgv.edu/chem_fac

 Part of the [Chemistry Commons](#)

Recommended Citation

Bandyopadhyay, Debasish; Mukherjee, Sanghamitra; and Banik, Bimal K., "An Expeditious Synthesis of N-Substituted Pyrroles via Microwave-Induced Iodine-Catalyzed Reactions Under Solventless Conditions" (2010). *Chemistry Faculty Publications and Presentations*. 46.

https://scholarworks.utrgv.edu/chem_fac/46

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Communication

An Expeditious Synthesis of *N*-substituted Pyrroles via Microwave-Induced Iodine-Catalyzed Reactions under Solventless Conditions

Debasish Bandyopadhyay, Sanghamitra Mukherjee and Bimal K. Banik*

Department of Chemistry, The University of Texas-Pan American, 1201 West University Drive, Edinburg, TX 78541, USA

* Author to whom correspondence should be addressed; E-Mail: banik@panam.edu; Tel.: +956-380 8741; Fax: +956-384-5006.

Received: 20 January 2010; in revised form: 1 March 2010 / Accepted: 2 April 2010 /


Published: 9 April 2010

Abstract: An expeditious synthesis of *N*-substituted pyrroles has been developed by reacting 2,5-dimethoxy tetrahydrofuran and several amines using a microwave-induced molecular iodine-catalyzed reaction under solventless conditions.

Keywords: pyrrole; molecular iodine; microwave irradiation; solventless reaction

1. Introduction


Pyrroles are an important class of organic compounds with different types of medicinal activities [1,2], consequently, many methods for the synthesis of diversely substituted pyrroles have been developed [3]. However, the most reliable method for the synthesis of pyrroles is the Paal-Knorr reaction [4,5]. Because of the biological activities of our polyaromatic compounds [6–10], we became interested in the synthesis of pyrroles bound to the amines of different structures. We describe here a simple method of synthesis of *N*-substituted pyrroles by reacting 2,5-dimethoxytetrahydrofuran (**1**) and various amines in the absence of any solvent in a microwave oven in the presence of catalytic amounts (~5 mol%) of molecular iodine. This reaction produces pyrroles in excellent yield and within a short time under (Scheme 1).

Scheme 1. Iodine-catalyzed microwave-induced synthesis of *N*-substituted pyrroles.

2. Results and Discussion





















This reaction has been tested by reacting different types of amines **2** and 2,5-dimethoxytetrahydrofuran in a microwave oven in the presence of molecular iodine as a catalyst. The reaction yields the products extremely well in the absence of any solvent. Reaction conditions and the yields of the products are shown in Table 1. Both aliphatic and aromatic amines produce pyrroles in very high yield. Importantly, the reaction produces product with multicyclic aromatic amines. A diamine (entry 9) and a heteropolyaromatic amine (entry 10) required higher temperature and longer reaction times, probably due to the diminished availability of the nitrogen (amine) lone pair.

We hypothesized that our work on iodine-catalyzed [11–16] reactions on acetal and glycosylation might prove useful for the facile synthesis of pyrroles under mild conditions. The methoxy groups in 2,5-dimethoxytetrahydrofuran (**1**) can be deprotected under mild acidic conditions and microwave irradiation. The intermediate can easily form the dialdehyde **5**. On reaction with amines and the dialdehyde, pyrroles **3** can be prepared following a nucleophilic addition and subsequent dehydration-aromatization route (Scheme 2). After irradiating a CDCl₃ solution of **1** for 5 minutes, ¹H-NMR has been taken. A downfield signal due to the –CHO group is observed. The intensity of the –CHO group becomes more predominant in the ¹H-NMR when **1** was irradiated in CDCl₃ in the presence of catalytic amounts of iodine. This suggests the facile formation of dialdehyde **5** in the reaction media in the presence of iodine and microwave irradiation.

Scheme 2. Iodine-catalyzed pyrrole synthesis: plausible mechanism of the reaction.

The presence of small amount of iodine (~5 mol %) is essential for the success of the reaction.

Table 1. Microwave-induced iodine-catalyzed synthesis of pyrroles.

Entry	Amine	MWI (power /temp /time)	Product	Yield (%)
1		300 watts 60°C 5min.		98
2		300 watts 50°C 5min.		96
3		300 watts 75°C 5min.		92
4		300 watts 50°C 5min.		95
5		300 watts 100°C 15min.		98
6		300 watts 110°C 10 min.		90
7		300 watts 120°C 20 min.		85
8		300 watts 100°C 5 min.		85
9		300 watts 130°C 40 min.		82
10		300 watts 160°C 55 min.		75

3. Experimental

3.1. General

Melting points were determined in a Fisher Scientific electrochemical Mel-Temp* manual melting point apparatus (Model 1001) equipped with a 300 °C thermometer. Elemental analyses (C, H, N)

were conducted using the Perkin-Elmer 2400 series II elemental analyzer, their results were found to be in good agreement ($\pm 0.2\%$) with the calculated values for C, H, N. FT-IR spectra were registered on a Bruker IFS 55 Equinox FTIR spectrophotometer as KBr discs. $^1\text{H-NMR}$ (300 MHz) and $^{13}\text{C-NMR}$ (75.4 MHz) spectra were obtained at room temperature with JEOL Eclipse-300 equipment using TMS as internal standard and CDCl_3 as solvent. Analytical grade chemicals (Sigma-Aldrich Corporation) were used throughout the project. Deionized water was used for the preparation of all aqueous solutions.

3.2. General procedure for the synthesis of pyrroles (3)

Amine **2** (1.0 mmol), 2,5-dimethoxytetrahydrofuran (**1**, 1.2 mmol) and iodine (5 mol%) was irradiated in a CEM automated microwave oven, as specified in Table 1. After completion of the reaction (monitored by TLC) diethyl ether (10 mL) was added to the reaction mixture which was then filtered. Pure product was isolated from the reaction mixture after evaporation of ether. Spectroscopic data for representative compounds, e.g. monoaromatic (entries 1 and 2), alicyclic (entry 5), polyaromatic (entries 6 and 7) as well as heteropolyaromatic (entry 10) are as follows:

1-Phenyl-1H-pyrrole (**3a**, entry 1). Brown sticky oil. IR: 2923, 1312, 1106, 782, 611 cm^{-1} ; $^1\text{H-NMR}$ δ (ppm): 6.39 (m, 2H, pyrrole), 7.24 (d, 2H, $J = 3.12$, pyrrole), 7.42–7.53 (m, 5H, Ar-H); $^{13}\text{C-NMR}$ δ (ppm): 109.07 (2C), 115.51 (2C), 122.96 (2C), 124.57, 128.62 (2C), 137.08. Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{N}$: C, 83.88; H, 6.34; N, 9.78. Experimental: C, 83.79; H, 6.31; N, 8.72.

1-(4-Methoxyphenyl)-1H-pyrrole (**3b**, entry 2). Black amorphous solid. Solidified from dichloromethane/hexane mixture; Mp: 89 $^\circ\text{C}$; IR: 2954, 1301, 1191, 850, 748 cm^{-1} ; $^1\text{H-NMR}$ δ (ppm): 3.87 (s, 3H, OCH_3), 6.33 (m, 2H, pyrrole), 6.67 (d, 2H, $J = 3.30$), 6.96–7.73 (m, 4H, Ar-H); $^{13}\text{C-NMR}$ δ (ppm): 55.66 (OCH_3), 109.93 (2C), 114.71 (2C), 120.15 (2C), 126.07 (2C), 132.56, 158.21. Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{NO}$: C, 76.28; H, 6.40; N, 8.09. Experimental: C, 76.20; H, 6.27; N, 7.97.

1-(Adamantan-1-yl)-1H-pyrrole (**3e**, entry 5). Yellow crystals. Crystallized from diethyl ether/hexane mixture; Mp: 71 $^\circ\text{C}$; IR: 2923, 2855, 1479, 1450, 1219, 713, 619 cm^{-1} ; $^1\text{H-NMR}$ δ (ppm): 1.77 (m, 3H), 2.12 (d, 6H, $J = 3.00$), 2.23 (m, 6H), 6.19 (t, 2H, $J = 2.19$, pyrrole), 6.91 (m, 2H, pyrrole); $^{13}\text{C-NMR}$ δ (ppm): 29.79 (3C), 36.35 (3C), 44.04 (3C), 55.00, 107.24 (2C), 116.55 (2C). Anal. Calcd for $\text{C}_{14}\text{H}_{19}\text{N}$: C, 83.53; H, 9.51; N, 6.96. Experimental: C, 83.41; H, 9.50; N, 6.89.

1-(Chrysen-6-yl)-1H-pyrrole (**3f**, entry 6). Brown crystals. Crystallized from ethyl acetate/hexane mixture; Mp: 139 $^\circ\text{C}$; IR: 2947, 2924, 1513, 1461, 1071, 812 cm^{-1} ; $^1\text{H-NMR}$ δ (ppm): 6.49 (t, 2H, $J = 2.21$, pyrrole), 7.12 (m, 2H, pyrrole), 7.70–8.72 (m, 11H, Ar-H); $^{13}\text{C-NMR}$ δ (ppm): 109.27 (2C), 120.98, 121.29 (2C), 123.23, 123.49 (2C), 123.64 (2C), 124.05, 126.45 (2C), 126.75 (2C), 126.88 (2C), 127.42, 128.64 (2C), 132.26, 137.94. Anal. Calcd for $\text{C}_{22}\text{H}_{15}\text{N}$: C, 90.08; H, 5.15; N, 4.77. Experimental: C, 89.91; H, 5.04; N, 4.76.

1-(Pyren-1-yl)-1H-pyrrole (**3g**, entry 7). Pale yellow crystals. Crystallized from diethyl ether/hexane mixture; Mp: 78 $^\circ\text{C}$; IR: 2956, 2927, 1598, 1511, 1304, 1072, 848 cm^{-1} ; $^1\text{H-NMR}$ δ (ppm): 6.53 (m, 2H, pyrrole), 7.15 (m, 2H, pyrrole), 8.01–8.19 (m, 9H, Ar-H); $^{13}\text{C-NMR}$ δ (ppm): 109.45 (2C),

122.36, 123.83 (2C), 124.89, 125.42, 126.51, 126.72, 127.19, 127.89 (2C), 128.63 (2C), 128.92, 130.62, 131.01, 131.39, 133.43, 135.97. Anal. Calcd for C₂₀H₁₃N: C, 89.86; H, 4.90; N, 5.24. Experimental: C, 89.77; H, 4.81; N, 5.17.

5-(1*H*-Pyrrol-1-yl)-1,10-phenanthroline (**3j**, entry 10). Yellowish brown amorphous solid. Solidified from methanol/hexane mixture; Mp: 192 °C; IR: 3186, 2362, 2336, 1591, 1539, 1074, 739 cm⁻¹; ¹H-NMR δ (ppm): 6.44 (broad s, 2H, pyrrole), 7.02 (broad s, 2H, pyrrole), 7.62–9.20 (m, 7H, Ar-H); ¹³C-NMR δ (ppm): 110.14 (2C), 122.79, 123.24 (2C), 123.55 (2C), 123.74 (2C), 129.04, 132.26, 136.17, 143.91, 144.09, 150.80, 150.91. Anal. Calcd for C₁₆H₁₁N₃: C, 78.35; H, 4.52; N, 17.13. Experimental: C, 78.23; H, 4.43; N, 17.02.

4. Conclusions

In summary, the present iodine-catalyzed microwave-induced method in the absence of any solvent is excellent for the preparation of *N*-substituted pyrroles (75–98% yields). Because of the simplicity of the procedure, products can be isolated very easily. The compounds reported herein will be tested against a number of cancer cells *in vitro*.

Acknowledgements

We gratefully acknowledge the funding support from National Cancer Institute (NIH/NCI-P20, Grant# 5P20CA138022-02).

References and Notes

1. Lainton, J.A.H.; Hoffman, J.W.; Martin, D.R.; Compton, D.R. 1-Alkyl-3-(1-naphthoyl)pyrroles: a new class of cannabinoid. *Tetrahedron Lett.* **1995**, *36*, 1401–1404.
2. de Leon, C.Y.; Ganem, B. A new approach to porphobilinogen and its analogs. *Tetrahedron* **1997**, *53*, 7731–7752.
3. Gilchrist, T.L. Synthesis of aromatic heterocycles. *J. Chem. Soc., Perkin Trans 1* **1998**, 615–628 and references cited therein.
4. Ruault, P.; Pilard, J-F.; Touaux, B.; Boulet, F.T.; Hamelin, J. Rapid generation of amines by microwave irradiation of ureas dispersed on clay. *Synlett* **1994**, 935–936.
5. Danks, T.N. Microwave assisted synthesis of pyrroles. *Tetrahedron Lett.* **1999**, *40*, 3957–3960.
6. Becker, F.F.; Banik, B.K. Polycyclic aromatic compounds as anticancer agents: synthesis and biological evaluation of some chrysene derivatives. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2877–2880.
7. Becker, F.F.; Mukhopadhyay, C.; Hackfeld, L.; Banik, I.; Banik, B.K. Polycyclic aromatic compounds as anticancer agents: synthesis and biological evaluation of dibenzofluorene derivatives. *Bioorg. Med. Chem.* **2000**, *8*, 2693–2699.
8. Banik, B.K.; Becker, F.F. Polycyclic aromatic compounds as anticancer agents. 4. Structure-activity relationships of chrysene and pyrene derivatives. *Bioorg. Med. Chem.* **2001**, *9*, 593–605.

9. Banik, B.K.; Becker, F.F. Synthesis, electrophilic substitution and structure-activity relationship studies of polycyclic aromatic compounds towards the development of anticancer agents. *Curr. Med. Chem.* **2001**, *8*, 1513–1533.
10. Banik, I.; Becker, F.F.; Banik, B.K. Stereoselective Synthesis of β -Lactams with Polyaromatic Imines: Entry to New and Novel Anticancer Agents. *J. Med. Chem.* **2003**, *46*, 12–15.
11. Banik, B.K.; Manhas, M.S.; Bose, A.K. Stereospecific Glycosylation via Ferrier Rearrangement for Optical Resolution. *J. Org. Chem.* **1994**, *59*, 4714–4716.
12. Banik, B.K.; Manhas, M.S.; Bose, A.K. Studies on lactams. 103. Enantiopure α -hydroxy- β -lactams via stereoselective glycosylation. *Tetrahedron Lett.* **1997**, *38*, 5077–5080.
13. Banik, B.K.; Zegrocka, O. Manhas, M.S.; Bose, A.K. Studies on lactams. 104. Enantiomerically pure β -lactams with the thienamycin side chain via glycosylation. *Heterocycles* **1997**, *27*, 173–176.
14. Banik, B.K.; Fernandez, M.; Alvarez, C. Iodine-catalyzed highly efficient Michael reaction of indoles under solvent-free condition. *Tetrahedron Lett.* **2005**, *46*, 2479–2482.
15. Banik, B.K.; Garcia, I.; Morales, F.; Aguilar, C. Novel synthesis of substituted pyrrole bound to indolinone via molecular iodine-catalyzed reaction. *Heterocycl. Commun.* **2007**, *13*, 109–112.
16. Banik, B.K.; Garza, R. Molecular iodine-catalyzed deprotection of acetals and ketals in acetone. *Chem. Edu.* **2007**, *12*, 75–76.

Sample Availability: Samples of the compounds are available from the authors.

© 2010 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).