

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

Chemistry Faculty Publications and  
Presentations

College of Sciences

---

11-1-2007

## An Overview of the Genus *Nardostachys*

Asima Chatterjee

Utpal Dutta

Debasish Bandyopadhyay

*The University of Texas Rio Grande Valley*, [debasish.bandyopadhyay@utrgv.edu](mailto:debasish.bandyopadhyay@utrgv.edu)

Anupam Nayak

Bidyut Basak

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.utrgv.edu/chem\\_fac](https://scholarworks.utrgv.edu/chem_fac)

 Part of the [Chemistry Commons](#)

---

### Recommended Citation

Chatterjee, A., Dutta, U., Bandyopadhyay, D., Nayak, A., Basak, B., Banerji, A., & Banerji, J. (2007). An Overview of the Genus *Nardostachys*. *Natural Product Communications*. <https://doi.org/10.1177/1934578X0700201124>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Chemistry Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

---

**Authors**

Asima Chatterjee, Utpal Dutta, Debasish Bandyopadhyay, Anupam Nayak, Bidyut Basak, Avijit Banerji, and Julie Banerji

An Overview of the Genus *Nardostachys*

(Late) Asima Chatterjee, Utpal Dutta, Debasish Bandyopadhyay<sup>#</sup>, Anupam Nayak, Bidyut Basak<sup>†</sup>, Avijit Banerji and Julie Banerji<sup>\*</sup>

Centre of Advanced Studies on Natural Products including Organic Synthesis, Department of Chemistry, Calcutta University, 92; A.P.C. Road, Kolkata 700009, India

<sup>#</sup>Present address: Department of Chemistry, The University of Texas-Pan American, Edinburg, TX, USA

<sup>†</sup>Present address: DCCS, Rajbandh, Durgapur, West Bengal, India

juliebanerji@yahoo.co.in

Received: May 22<sup>nd</sup>, 2007; Accepted: September 20<sup>th</sup>, 2007


*Nardostachys jatamansi*, a medicinally important herb of Nepalese origin, has been used for centuries in the Ayurvedic and Unani systems of medicine. In combination with *Marsilea minuta* it is being used as an antistress and anticonvulsant drug and also finds use in the treatment of epilepsy. Recently, it has been reported that *N. jatamansi*, which plays an important role in protecting from cerebral ischemia and liver damage, is also used for the treatment of osteoporosis and hypercalcemia. The other member of the genus *Nardostachys*, *N. chinensis*, possesses antifungal and antimalarial properties. It is also used in the treatment of skin dysfunction. A short summary of the chemical constituents of the two species along with their physical and biological properties is reported.

**Keywords:** *Nardostachys jatamansi*, *Nardostachys chinensis*, Valerianaceae, chemical structure, biological importance.


*Nardostachys jatamansi* DC. is a high altitude plant, growing in the Himalayan foothills. Traditionally, *Nardostachys*, has been recommended in the Ayurvedic system of medicine for nervous and spasmodic symptoms, viz. heart palpitation, convulsion and hysteria [1-3].

The genus *Nardostachys* belongs to the family Valerianaceae and consists of only two species [4]; *N. jatamansi* DC and *N. chinensis* Batal. In the Indian subcontinent the oil of *N. jatamansi* is used in several massage preparations. On the other hand, Gansong, a widely used analgesic herb in China, known as *N. chinensis* [5] is used for treating pain in the chest and abdomen [6]. The combination drug made from *Nardostachys jatamansi* and *Marsilea minuta* under the code name, Ayush 56, has been patented [7] by the Central Council for Research in Ayurveda and Siddha, Government of India, and purchased by National Research Development Corporation. An intensive search for the active constituents present in this genus revealed that the roots and rhizomes are potential sources of sesquiterpenes of common and lesser-known skeleta as discussed below.

(i) **Aristolane type:** Both species i.e., *N. jatamansi* and *N. chinensis*, contain several sesquiterpenes possessing the aristolane moiety (Figure 1). The compounds isolated are mentioned in Table 1.



Aristolane type sesquiterpene



Aristolane type sesquiterpene

Figure 1

Figure 2

Representatives of another type of aristolane sesquiterpene (Figure 2) isolated from this genus are shown in Table 2.

**Spectral data of Aristolane Sesquiterpenes:** The spectral data of Kanshone F (considered as a representative member of the aristolane skeleton) has been discussed [24].


**Table 1:** Aristolane sesquiterpenes (Figure 1) isolated from the genus *Nardostachys*.

No.	Compound	Mol. Formula & Physical State	Ref.	Source	Substituents					
					R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
i)	9,10-Dehydroaristolene	C <sub>15</sub> H <sub>24</sub> , b.p. 110°	[8, 9]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (oil)	H	H	H	H	—	—
ii)	9-Aristolene-1 $\alpha$ -ol (Nardostachnol)	C <sub>15</sub> H <sub>24</sub> O, m.p. 85°	[8, 10-13]	<i>N. jatamansi</i> (roots) <i>N. chinensis</i> (oil)	OH ( $\alpha$ )	H	H	H	—	—
iii)	1(10)-Dehydroaristolene (calarene)	C <sub>15</sub> H <sub>24</sub>	[8, 9, 11, 14- 17, 48]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (oil)	—	H	H	H	H	—
iv)	1,2,9,10-Tetrahydroaristolene	C <sub>15</sub> H <sub>22</sub>	[8, 12]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (oil)	—	—	H	H	—	—
v)	1(10), 8(9)-Aristoladien-2-one	C <sub>15</sub> H <sub>20</sub> O	[18, 19]	<i>N. chinensis</i> (oil)	—	=O	H	—	—	—
vi)	1(10)-Aristolene-9-one (Gansongone)	C <sub>15</sub> H <sub>22</sub> O	[13, 20]	<i>N. chinensis</i> (oil)	—	H	H	H	=O	—
vii)	1(10)-Aristolene-2-one	C <sub>15</sub> H <sub>22</sub> O	[10,14,18,19]	<i>N. jatamansi</i> (roots) <i>N. chinensis</i> (oil)	—	=O	H	H	H	—
viii)	9-Hydroxy-1(10)-Aristolene-2-one (Debilon)	C <sub>15</sub> H <sub>22</sub> O <sub>2</sub>	[21-23]	<i>N. chinensis</i> (roots & rhizomes)	—	=O	H	H	OH ( $\alpha$ )	—
ix)	Kanshone F	C <sub>20</sub> H <sub>30</sub> O <sub>3</sub>	[24]	<i>N. chinensis</i> (roots)	OCOCH <sub>2</sub> -i-pr ( $\beta$ )	H	H	=O	—	—
x)	Kanshone G	C <sub>15</sub> H <sub>24</sub> O <sub>2</sub>	[24]	<i>N. chinensis</i> (roots)	—	OH ( $\beta$ )	H	H	OH ( $\beta$ )	—


**Table 2:** Aristolane sesquiterpenes (another type) (Figure 2) isolated from these genera.

No.	Compound	Mol. Formula & Physical State	Ref.	Source	Substituents					
					R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
i)	* $\beta$ -Maaliene	C <sub>15</sub> H <sub>24</sub>	[8, 9, 17, 25-27]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (oil)	H	Me ( $\alpha$ )	H	Me	—	—
ii)	*Maaliol	C <sub>15</sub> H <sub>26</sub> O m.p. 103.5-105°	[25, 28]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (rhizomes)	H	Me ( $\alpha$ )	H	OH ( $\beta$ )	Me ( $\alpha$ )	H
iii)	Calareneol	C <sub>15</sub> H <sub>25</sub> O	[11, 29]	<i>N. jatamansi</i> (oil)	H	—	—	OH ( $\beta$ )	Me ( $\alpha$ )	Me ( $\alpha$ )

\*Later in 1967, Ruecker et al. [25] reported that these two compounds were not found in *N. chinensis*.



Kanshone F [24]  
C<sub>20</sub>H<sub>30</sub>O<sub>3</sub>  
**Figure 3**



Kanshone C [20,30]  
C<sub>15</sub>H<sub>20</sub>O<sub>3</sub>, m.p. 136-137°  
**Figure 4**

The olefinic proton at the  $\alpha$ -carbon (C-9) of the  $\alpha,\beta$ -unsaturated ketone appeared at  $\delta$  5.93 as a doublet of doublets. Two methine protons at C-1 and C-4 resonated at  $\delta$  5.55 and  $\delta$  1.88 respectively. H-6 and H-7 were discernible at  $\delta$  1.33 and  $\delta$  1.77 while the methyl protons (at C-12, C-13, C-14, C-15, C-4', and C-5') of the six-methyl groups resonated at  $\delta$  1.22, 1.26, 1.24, 1.09, 0.97, and 0.98 respectively.

The <sup>13</sup>C NMR spectrum of the compound showed the presence of an  $\alpha,\beta$ -unsaturated carbonyl (C-8) at  $\delta$  195.9 and the ester carbonyl (C-1') at  $\delta$  171.9. Two olefinic carbons (C-9 and C-10) were discernible at

$\delta$  120.8 and  $\delta$  163.0. Two methine carbons (C-6 and C-7) linked to the isopropyl group appeared at  $\delta$  39.5 and  $\delta$  35.0 respectively. One methine (C-11) and two methyl carbons (C-12 and C-13) in the isopropyl group appeared at  $\delta$  24.4, 29.7, and 16.6 respectively.

Interestingly, the abundance of aristolane derivatives [8] present in *N. chinensis* was greater than in *N. jatamansi*, except for 9-aristolene-1-ol (nardostachnol) as indicated in Table 3.

Besides the aristolane derivatives, kanshone C (1, 10-epoxy-8, 9-aristoladione), which is an epoxy derivative of aristolane (Figure 4), was isolated from *N. chinensis*.

**Table 3:** Comparative abundance of the compounds isolated from *Nardostachys chinensis* and *N. jatamansi*.

No.	Compound	<i>N. chinensis</i> (%) [8]	<i>N. jatamansi</i> (%) [8]
i)	9-Aristolene-1 $\alpha$ -ol	8.0	69.4
ii)	9,10-dehydro Aristolene	2.1	0.2
iii)	1(10)-dehydro Aristolene	8.6	2.0
iv)	$\beta$ -Maaliene	32.8	2.9
v)	1,2,9,10-Tetrahydroaristolene	13.1	2.9

(ii) **Eudesmane type:** Sesquiterpenes possessing this skeleton were mainly isolated from *N. jatamansi*. Until now, only one compound was reported from *N. chinensis* (Table 4) (Figure 5).




Figure 5




Figure 6

(iii) **Guaiane type:** Investigation by a group of Japanese workers led to the isolation of several sesquiterpenes with the guaiane skeleton (Figure 6) from *N. chinensis*, viz. nardoguaianone A to K (Tables 5 and 6). Of these, nardognainone A to D were endoperoxides. Further research revealed the presence of two more endoperoxides isolated from the roots of *N. chinensis*, viz. nardoperoxide and isonardoperoxide, together with nardoxide, which contained an epoxy linkage (Figure 8) instead of the usual endo peroxy system (Figure 7).

**Spectral data of Guaiane Sesquiterpenes:** Spectral data of nardoguaianones (E-I) have been considered as representative of the guaiane type sesquiterpenes [34]. From the spectral properties of the guaiane type compounds it was apparent that the two protons at C-3, i.e. 3 $\alpha$ -H and 3 $\beta$ -H, resonated in the region  $\delta$  2.02-2.07 and  $\delta$  2.62-2.68 respectively as a pair of doublet of doublets. The methine proton at C-4 and the olefinic proton at C-6 appeared in the region  $\delta$  2.75-2.88 (ddq) and  $\delta$  5.75-6.33 (br t).

The -OH proton at C-8 in nardoguaianones E-G was discernible in the region  $\delta$  4.44-4.52, whereas the two methylene protons at C-8 in nardoguaianones H and I appeared in the region  $\delta$  2.43-2.45 (br. dd) and  $\delta$  2.33-2.36 (ddd). Six methyl protons at C-12 and C-13 in nardoguaianones E-G appeared in the region  $\delta$  1.15-1.18 (d) and  $\delta$  1.13-1.15 (d), whereas the methyl protons in nardoguaianone H and I showed signals in the region  $\delta$  1.14-1.42 (s) and  $\delta$  1.43-1.44 (s) respectively due to the presence of the -OH group at C-11. Six more methyl protons at C-14 and C-15 in all these compounds (nardoguaianones E-I) resonated in the region  $\delta$  1.13-1.26 and  $\delta$  1.44-1.49 respectively.


Table 4: Eudesmane type sesquiterpenes (Figure 5) isolated from the genus *Nardostachys*.

No.	Compound	Mol. Formula	Source	Substituents				Ref.
				R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	
i)	Jatamol A	C <sub>15</sub> H <sub>24</sub> O	<i>N. jatamansi</i> (roots)	OH ( $\alpha$ )	=CH <sub>2</sub>	-		[31]
ii)	Jatamol B	C <sub>20</sub> H <sub>32</sub> O <sub>3</sub>	<i>N. jatamansi</i> (roots)	OCOCH <sub>2</sub> C (Me <sub>2</sub> ) OH ( $\alpha$ )	=CH <sub>2</sub>	-		[31]
iii)	$\beta$ -Eudesmol	C <sub>15</sub> H <sub>16</sub> O	<i>N. jatamansi</i> (roots)	H	=CH <sub>2</sub>	-		[32]
iv)	Eudesm-11-en-2, 4-diol	C <sub>15</sub> H <sub>26</sub> O <sub>2</sub>	<i>N. chinensis</i>	OH ( $\alpha$ )	OH ( $\alpha$ )	Me ( $\beta$ )		[33]

Table 5: Guaiane type sesquiterpenes (Figure 6) isolated from *Nardostachys chinensis*.

No.	Compound	Mol. Formula	Source	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	Ref.
i)	Nardoguaianone E	C <sub>15</sub> H <sub>22</sub> O <sub>3</sub>	<i>N. chinensis</i> (roots)	OH ( $\alpha$ )	OH ( $\alpha$ )	Me ( $\beta$ )	[34]
ii)	Nardoguaianone F	C <sub>15</sub> H <sub>22</sub> O <sub>3</sub>	<i>N. chinensis</i> (roots)	OH ( $\beta$ )	OH ( $\beta$ )	Me ( $\alpha$ )	[34]
iii)	Nardoguaianone G	C <sub>15</sub> H <sub>22</sub> O <sub>3</sub>	<i>N. chinensis</i> (roots)	OH ( $\alpha$ )	OH ( $\beta$ )	Me ( $\alpha$ )	[34]
iv)	Nardoguaianone H	C <sub>15</sub> H <sub>22</sub> O <sub>3</sub>	<i>N. chinensis</i> (roots)	H	OH ( $\alpha$ )	Me ( $\beta$ )	[34]
v)	Nardoguaianone I	C <sub>15</sub> H <sub>22</sub> O <sub>2</sub>	<i>N. chinensis</i> (roots)	H	OH ( $\beta$ )	Me ( $\alpha$ )	[34]
vi)	Nardoguaianone J	C <sub>15</sub> H <sub>22</sub> O <sub>2</sub>	<i>N. chinensis</i> (roots)	H	Me ( $\beta$ )	OH ( $\alpha$ )	[24]
vii)	Nardoguaianone K	C <sub>15</sub> H <sub>22</sub> O <sub>2</sub>	<i>N. chinensis</i> (roots)	H	Me ( $\alpha$ )	OH ( $\beta$ )	[24]

In the  $^{13}\text{C}$  NMR spectrum the  $\alpha,\beta$ -unsaturated carbonyl carbon showed chemical shifts  $\sim\delta$  209.9-210.6. The four olefinic carbons (C-1, C-5, C-6, C-7) resonated in the region  $\delta$  140.3-142.1,  $\delta$  167.1-169.9,  $\delta$  113.1-117.4 and 164.4-167.9 respectively.



Guaiane type endoperoxide sesquiterpene

Figure 7


Nardoxide [35]  
C<sub>15</sub>H<sub>22</sub>O<sub>3</sub>

Figure 8


BR 606 [37]  
C<sub>16</sub>H<sub>24</sub>O<sub>5</sub>

Figure 9

However, the only epoxy sesquiterpene, named BR-606 (Figure 9), isolated from the roots of *N. jatamansi* has been frequently used for the treatment of osteoporosis [37].

**Spectral data of guaianane type endoperoxide moiety:** Spectral data of nardoperoxide and isonardoperoxide (considered as representative members of this skeleton) have been discussed [35].

The two protons at positions 3 and 9 appeared in the range  $\delta$  2.1-2.8 and  $\delta$  1.7-2.1 respectively, while H-4 was generally observed in the region  $\delta$  2.8-3.2 (Figure 6). The two primary carbons at 3 and 9 resonated at  $\delta$  45-46 and  $\delta$  28-29 while the two tertiary carbons at 4 and 6 were observed around

$\delta$  32-37 and  $\delta$  69-73 respectively. The carbonyl carbon was observed in the low field region in the range  $\delta$  204-205. The two quaternary fused carbon atoms at C-1 and C-5 appeared in the region  $\delta$  143-145 and  $\delta$  177-179 respectively whereas the two peroxide linked carbons, i.e. C-7 and C-10, resonated in the range  $\delta$  83-84 and  $\delta$  78-80.



Nardosaldehyde [38]  
C<sub>15</sub>H<sub>22</sub>O<sub>4</sub>

Figure 10

In this context, it is important to note that the sesquiterpene containing an endoperoxy linkage, nardosaldehyde, was isolated from *N. chinensis* roots (Figure 10). This compound is different from all other guaianane type endoperoxy sesquiterpenes isolated from this species.



Guaiadiene type sesquiterpene

Figure 11

(iv) **Guaiadiene type:** Three sesquiterpenes containing a guaiadiene moiety were isolated from *N. jatamansi* (Figure 11, Table 7). Among these compounds, *nardostachysin* needs special mention. It is an ester of jatamansic acid (guaiadiene type sesquiterpene) and a bicyclic monoterpene alcohol.


Table 6: Guaiane type endoperoxide sesquiterpenes (Figure 7) obtained from *Nardostachys chinensis*.

No.	Compound	Mol. Formula & Physical State	Ref.	Source	Substituents				Peroxo linkage
					R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	
i)	Nardoperoxide	C <sub>15</sub> H <sub>22</sub> O <sub>4</sub> m.p. 129-130°	[35]	<i>N. chinensis</i> (roots)	OH ( $\alpha$ )	i-pr ( $\beta$ )	H	Me ( $\beta$ )	O-O ( $\alpha$ )
ii)	Isonardoperoxide	C <sub>15</sub> H <sub>22</sub> O <sub>4</sub>	[35]	<i>N. chinensis</i> (roots)	OH ( $\beta$ )	i-pr ( $\alpha$ )	H	Me ( $\alpha$ )	O-O ( $\beta$ )
iii)	Nardoguaianone A	C <sub>15</sub> H <sub>22</sub> O <sub>4</sub>	[36]	<i>N. chinensis</i> (roots)	OH ( $\beta$ )	i-pr ( $\beta$ )	H	Me ( $\beta$ )	O-O ( $\alpha$ )
iv)	Nardoguaianone B	C <sub>15</sub> H <sub>22</sub> O <sub>4</sub>	[36]	<i>N. chinensis</i> (roots)	OH ( $\alpha$ )	i-pr ( $\alpha$ )	H	Me ( $\alpha$ )	O-O ( $\beta$ )
v)	Nardoguaianone C	C <sub>15</sub> H <sub>23</sub> O <sub>5</sub>	[36]	<i>N. chinensis</i> (roots)	OH ( $\alpha$ )	i-pr ( $\beta$ )	OH ( $\alpha$ )	Me ( $\beta$ )	O-O ( $\alpha$ )
vi)	Nardoguaianone D	C <sub>15</sub> H <sub>23</sub> O <sub>5</sub>	[36]	<i>N. chinensis</i> (roots)	OH ( $\beta$ )	i-pr ( $\alpha$ )	OH ( $\beta$ )	Me ( $\alpha$ )	O-O ( $\beta$ )

**Table 7:** Guaiadiene type sesquiterpenes (Figure 11) isolated from the genus *Nardostachys*.

No.	Compound	Mol. Formula & Physical State	Ref.	Source	Substituents					
					R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
i)*	Jatamansic acid	C <sub>15</sub> H <sub>22</sub> O <sub>2</sub> m.p. 123°	[15, 25, 39, 45, 46]	<i>N. jatamansi</i> (oil) <i>N. chinensis</i> (rhizomes)	H	—	—	—	—	COOH
ii)	Nardol	C <sub>15</sub> H <sub>26</sub> O b.p. 120-125°	[11, 15, 47]	<i>N. jatamansi</i> (roots)	OH ( $\alpha$ )	H	H	H	=CH <sub>2</sub>	—
iii)	Nardostachysin	C <sub>25</sub> H <sub>34</sub> O <sub>6</sub> m.p. 192°	[48]	<i>N. jatamansi</i> (rhizomes)	H	—	—	—	—	COOR*

Where, R\* =




\*Though jatamansic acid was reported earlier to be present in *N. chinensis*, its presence was contradicted by Ruecker *et al.* [25].

**Spectral data of Guaiadiene Sesquiterpenes:** Spectral data of nardostachysin and jatamansic acid (considered as representative compounds bearing the guaiadiene skeleton) have been discussed [39,48].

The two protons at the 7/5 ring juncture, i.e. H-1 and H-5, were discernible in the regions  $\delta$  3.0-3.1 and  $\delta$  1.65-1.75 respectively (Figure 10). The two-methine protons of the 7-membered ring, i.e. H-8 and H-9, resonated between  $\delta$  5.7-5.8 and  $\delta$  7.1-7.2 while the two-methyl and one-methine protons in the isopropyl group appeared at  $\delta$  1.07,  $\delta$  1.05, and  $\delta$  2.43 respectively. The <sup>13</sup>C NMR spectrum showed the presence of the acid carbonyl of jatamansic acid at  $\delta$  176.4 and the ester carbonyl of nardostachysin at  $\delta$  170.1. The two carbon atoms at the 7/5-ring juncture, i.e. C-1 and C-5 were observed at  $\delta$  45-47 and  $\delta$  43-45 while the two quaternary carbons of the 7-membered ring, i.e. C-7 and C-10, appeared in the regions  $\delta$  160-161 and  $\delta$  133-135 respectively. The three CH carbons of the 7/5-ring system (i.e. C-4, C-8, and C-9) were observed in the regions  $\delta$  38-40,  $\delta$  116-119, and  $\delta$  134-137 respectively.


(v) **Valerane and elemene type:** The roots and rhizomes of *N. jatamansi* were also found to contain valeranone and elemol as secondary metabolites. It is worth mentioning that valeranone possesses the valerane moiety whereas elemol is a member of the elemene group. The structures of the compounds have been shown in Figure 12.

So far, sesquiterpenes of common skeleta have been discussed. Several sesquiterpenes of uncommon

**Figure 12**

skeleta, viz. nardosinane [13,19-21,23,25,30,42, 49-54] and nor-eremophilane [11,13,21,55,56] type were isolated from *N. chinensis*.


(vi) **Nardosinane type:** From the oil and gum of *N. chinensis* several sesquiterpenes possessing the nardosinane skeleton were isolated viz. isonardosinone, kanshone A-B, kanshone D-E, nardofuran, nardonoxide, nardosinone and nardosinonediol. An investigation of their structure revealed that isonardosinone, kanshone D-E,

**Figure 13**

nardoxide, and nardofuran possessed an epoxy bridge whereas kanshone B and nardosinone contained a peroxy bridge (Figure 13).

**Significant spectral data of the Nardosinane moiety:**

The spectral data of kanshone A (considered as a representative member of the nardosinane skeleton) has been discussed [52].



In the  $^1\text{H}$  NMR spectrum the protons of the methyl groups at C-4 and C-5 resonated at  $\delta$  1.02 and 1.08 respectively whereas the gem-dimethyls at C-11 appeared at  $\delta$  1.18 and 1.25. The methine protons at C-4 and C-6 appeared at  $\delta$  2.20 and 2.62 and the two olefinic protons at C-1 and C-8 were discernible at  $\delta$  7.0 and 6.16 respectively.

In the  $^{13}\text{C}$  NMR study the carbonyl carbon appeared at  $\delta$  187.9, the two quaternary carbons at C-5 and C-10 at  $\delta$  42.0 and 141.6 respectively. The olefinic carbons at C-1, C-7, and C-8 were discernible at  $\delta$  137.2, 151.1, and 128.9.

(vii) **Noreremophilane type:** To date, three sesquiterpenoids having the noreremophilane moiety (Figure 14) have been isolated from this genus. Of these, narchinol A and desoxo-narchinol A were isolated from *N. chinensis* whereas nardostachone was obtained from *N. jatamansi* roots. Narchinol A and desoxonarchinol A belong to the tris-nor-sesquiterpenoid class.




Figure 14

(viii) **Patchoulane type:** Three sesquiterpenes of the rearranged patchoulane type skeleton, *viz.*  $\alpha$ -patchoulene,  $\beta$ -patchoulene, and patchouli alcohol, were isolated from the roots of *N. jatamansi* and

*N. chinensis*. The structures of these three compounds are shown in Figure 15.





Figure 15

Along with all the sesquiterpenes mentioned, the presence of a spiranic sesquiterpene in the roots and oil of *N. jatamansi* was also reported. The structure of this spiro-sesquiterpene, spirojatamol, is shown in Figure 16.



Sesquiterpenes possessing a bridgehead structure (Figure 17) were also isolated from the roots of *N. jatamansi*. A literature survey revealed the presence of seychellene, seychellane, and nor-seychellane, all possessing [3, 3, 1] system.




Figure 17

Two sesquiterpenes, nardin (sesquiterpene acid) and valeranal (sesquiterpene aldehyde), were isolated from the roots of *N. jatamansi* (Figure 18).







Figure 18



**Table 8:** Coumarins (Figure 20) isolated from the genus *Nardostachys*.


Structural type	Compound	Mol. Formula & Physical State	Ref.	Source	Substituents	
					R <sub>1</sub>	R <sub>2</sub>
A	Angelicin	C <sub>11</sub> H <sub>6</sub> O <sub>3</sub> , m.p. 137°	[32, 38]	<i>N. jatamansi</i> (roots)	H	—
A	Oroselol	C <sub>14</sub> H <sub>12</sub> O <sub>4</sub> , m.p. 148-151°	[27]	<i>N. jatamansi</i>		—
B	Jatamansin	C <sub>19</sub> H <sub>20</sub> O <sub>5</sub> , m.p. 97-98°	[27, 32]	<i>N. jatamansi</i>	—	
B	Jatamansinol (Lomatin)	C <sub>14</sub> H <sub>14</sub> O <sub>4</sub> , m.p. 182°	[14, 15, 32]	<i>N. jatamansi</i> <i>N. chinensis</i>	—	OH (β)
C	7-Methoxy-8-isopentenyl coumarin	C <sub>15</sub> H <sub>16</sub> O <sub>3</sub> , m.p. 81°C	Unpublished Result	<i>N. jatamansi</i> (roots)	OMe	—

Two new monoterpenes nardostachin and 6-hydroxy-7-(hydroxymethyl)-4-methylene hexahydro cyclopenta [c] pyran-1(3H)-one were isolated from the roots and rhizomes of *N. chinensis* (Figure 19).

Nardostachin [63]  
C<sub>20</sub>H<sub>32</sub>O<sub>6</sub>6-hydroxy-7-(hydroxymethyl)-4-methylene hexahydro cyclopenta [c] pyran-1-(3H)-one  
C<sub>10</sub>H<sub>14</sub>O<sub>4</sub> [23]10-isopropyl-2,2,6-trimethyl-2,3,4,5-tetrahydronaphtha [1,8-bc] oxocine-5,11-diol C<sub>20</sub>H<sub>26</sub>O<sub>3</sub> [23]**Figure 19**

A new diterpene was isolated from the roots and rhizomes of *N. chinensis* (Figure 19) and characterized as 10-isopropyl-2,2,6-trimethyl-2,3,4,5-tetrahydronaphtha [1,8-bc] oxocine-5,11-diol [23].



(ix) **Coumarins:** Besides terpenes, coumarin derivatives were also present in *N. jatamansi* [Table 8]. The coumarin derivatives isolated have been subdivided into furanocoumarins (Structure A) and pyranocoumarins (Structure B) (Figure 20). Later studies revealed the presence of osthol (Structure C). Jatamansinol is the only coumarin isolated from *N. chinensis* to date.

**Figure 20**

(x) **Lignans and Neolignans:** Besides sesquiterpenes and coumarin derivatives, the presence of lignans and neolignans has also been reported from this genus.

The two-tetrahydrofuran lignans, isolated from the roots of *N. jatamansi*, were pinosresinol and 1-hydroxy pinosresinol (Figure 21).

Another class of compounds i.e. neolignans were also isolated from the roots of *N. jatamansi* (Figure 22).

**Figure 21****Figure 22**

Further literature survey revealed that only one alkaloid, i.e., actinidine (Figure 23), was obtained from the rhizomes of *N. jatamansi*.




Figure 23

GC-MS studies of *N. chinensis* and *N. jatamansi* revealed the presence of several different types of compounds in trace amounts. These are listed in Table 9.

Table 9: Minor compounds from *N. jatamansi*.

No.	Compound	Ref.	No.	Compound	Ref.
1	$\alpha$ -Pinene	[15,17, 32]	22	Ledol	[15]
2	$\beta$ -Pinene	[15,17, 32]	23	Cadina-1(10),6,8-triene	[15]
3	$\Delta^3$ -Carene	[32]	24	$\alpha$ -Copaene	[15]
4	$\beta$ -Sitosterol	[10,11, 13, 32, 66]	25	Valencene	[15]
5	<i>n</i> -Hexacosanol	[11]	26	$\alpha$ -Selinene	[15]
6	<i>n</i> -Hexacosane	[11]	27	$\alpha$ -Cubebene	[15, 17]
7	<i>n</i> -Hexacosanyl arachidate	[11]	28	$\beta$ -Copaene	[15]
8	<i>n</i> -Hexacosanyl isovalerate	[11]	29	7-Hexadecene	[15]
9	Isovaleric acid	[11]	30	Germacrene-D	[15, 17]
10	Formic acid	[15]	31	<i>cis</i> - $\beta$ -Farnesene	[15]
11	Bornyl acetate	[15]	32	$\beta$ -Elemene	[15]
12	Propionic acid	[15]	33	<i>trans</i> -Nerolidol	[15]
13	$\alpha$ -Terpineol	[15, 17]	34	Cycloprop(7,8) azulino(3a,4-b) oxirene	[15]
14	<i>p</i> -Cymene	[15, 17]	35	3,4-Dihydro- $\alpha$ -ionone	[15]
15	<i>n</i> -Hexane	[15]	36	( $\pm$ )4-Terpeneol	[15]
16	<i>n</i> -Nonane	[15]	37	$\beta$ -Selinene	[15, 17]
17	2-Ethylhexyl phthalate	[15]	38	$\beta$ -Caryophyllene	[15]
18	$\beta$ -Myrcene	[15, 17]	39	Cubebol	[15]
19	1,8-Cineole	[15]	40	$\alpha$ -Gurjunene	[15, 17]
20	Calamenene	[15]	41	$\gamma$ -Gurjunene	[15]
21	Myrtenol	[15]	42	$\alpha$ -Humulene	[15, 17]

Compounds, such as ursolic acid [63],  $\beta$ -Sitosterol [13], oleanolic acid [13], ethyl- $\beta$ -D-glucopyranoside [13] and  $\beta$ -ionone [17, 19] were also isolated from *N. chinensis*.

### Medicinal uses of *N. jatamansi* and *N. chinensis*:

*N. jatamansi* has been prescribed in India since 800 B.C. for different ailments such as epilepsy, chest pain and palpitation [67]. Recently, it was observed that *N. jatamansi* had a protective effect on cerebral ischemia [68]. The exact mechanism of this effect is still uncertain. In cerebral ischemia there is an excessive accumulation of glutamate in extra-cellular fluid [69] and of sodium and calcium ions in the intracellular fluid [70]. *N. jatamansi* enhances the function of  $\gamma$ -amino butyric acid [71-73] by stimulating the  $\gamma$ -amino butyric acid neurons which antagonize the effect of glutamate and saves the neurons from cerebral ischemia. However, in ischemia, accumulation of free radicals as well as decreased function of free radical scavengers like Superoxide Dismutase (SOD) was also observed [74-76]. The antioxidant property of *N. jatamansi* protects the neurons from ischemic injury [77].

Jatamansone, a sesquiterpene ketone, isolated from the rhizomes of *N. jatamansi*, showed antiarrhythmic and anticonvulsant activities [44]. This substance has also been used in febrile delirium and delirium tremors. Jatamansone semicarbazone was found to possess antiestrogenic activity, which was reflected in its estrogen-antagonizing action on the uterus [78].

A novel compound, code name BR-606, isolated from *N. jatamansi* roots, has been found to be useful in the treatment of osteoporosis and hypercalcemia [37]. The presence of the epoxy linkage in this compound is probably the key factor (Figure 9) as this is known to inhibit bone sorption and help to correct osteoporosis and hypercalcemia.

The roots and rhizomes of *N. chinensis* (Japanese crude drug name "Kanshoko") are used as a sedative and an analgesic in oriental medicines [24].

*N. chinensis* is known to be rich in terpenoids. Some of these possess antimalarial activity. Although, a number of medicines such as chloroquine and quinine are frequently used for the treatment of malaria, the rapid development of drug resistance poses a serious problem. Therefore new medicinal agents are required to overcome the drug resistance as well as to control epidemics caused by malarial parasites. During the search for new types of antimalarial compounds from plants, nardosinone [25] (Figure 13), an important constituent of *N. chinensis* roots, was found to exhibit antimalarial activity against *Plasmodium falcifarum* (EC<sub>50</sub> 4.5 x 10<sup>-6</sup> M). Further

research on this plant revealed the presence of three more guaiane type sesquiterpenoids, viz. nardoperoxide [35], isonardoperoxide [35] and nardoxide [36] (Table 6, Figure 7), all of which showed strong antimalarial activity. Further attempts to isolate antimalarial drugs led to the isolation of four guaiane-type endoperoxides, nardoguaianone A-D [36] from the roots. It was subsequently established that all these compounds have stronger antimalarial activity than the well-known antimalarial chloroquine. It is worthwhile to mention that nardoguaianone E to K [24, 34] (Table 5) does not exhibit any antimalarial activity. A close examination of their structures and the lack of the antimalarial activity of nardoguaianones E to K revealed that only those compounds having peroxy or oxo linkages exhibit antimalarial activity.

Among these antimalarial sesquiterpenoids, nardoperoxide, isonardoperoxide, nardoxide and

nardosinone showed cytotoxicities against FM3A and KB cells [35]. Malaria infected cells exhibit specific cytotoxic effects on treatment with these compounds and thereby these can be considered to be effective in the treatment of the disease.

From the above discussion it can be inferred that the genus '*Nardostachys*' is rich in several classes of compounds viz. sesquiterpenes, coumarins, lignans and neo-lignans. The never-ending requirement for the exploration of the rich medicinal armamentarium, with which nature has endowed mankind, stimulated the impetus to undertake in depth study of this genus. This resulted in the development of Ayush 56 [7] the combination-drug with *Marsilea minuta*, which finds use as an anti-stress, anti-convulsant and rehabilitation drug. The discovery of this combination-drug is a landmark in developing "alternate lines of treatment" leaving no side effects.

## References

- [1] Chittendon F. (1956) *RHS Dictionary of Plants plus Supplement*. Oxford University Press, 1951.
- [2] Chopra RN, Nayar SL, Chopra IC. (1956) *Glossary of Indian Medicinal Plants*. CSIR, New Delhi.
- [3] Chatterjee A, Pakrashi S. (1997) *The Treatise of Indian Medicinal Plants*. NISCAIR, New Delhi. Vol.5, 99.
- [4] Dharmananda S. (2000) *Valerian and Nardostachys*. [www.itmonline.org/arts/valerian.htm](http://www.itmonline.org/arts/valerian.htm)
- [5] Duke JA, Ayensu ES. (1985) *Medicinal Plants of China*. Reference Publications.
- [6] Stuart Rev GA. (1976) *Chinese Materia Medica*. Taipei.
- [7] Chatterjee A, Mukherjee, GD, Majhi SN, Bhattacharya P. (1984) CCRAS, New Delhi, Indian Patent No. 14170 dt.14.7.1976.
- [8] Sun HD, Ding JK, Lin ZW, Che FR. (1980) Study on the chemical constituents of the essential oils of *Nardostachys grandiflora* DC. and *N. chinensis* Batalin and their uses on the perfume. *Yunnan Zhiwu Yanjiu*, 2, 213-223.
- [9] Pesnelle P, Ourisson G. (1963) Aristolene isolation in nard essence. *Bulletin de la Societe Chimique de France*, 912-913.
- [10] Maheshwari ML, Saxena DB. (1980) 9-Aristolene-1 $\alpha$ -ol and 1(10)-aristolene-2-one from *Nardostachys jatamansi* DC. *Indian Perfumer*, 24, 171-174.
- [11] Sastry SD, Maheshwari ML, Chakravarti KK, Bhattacharyya SC. (1967) Terpenoids. CXV. Chemical constituents of *Nardostachys jatamansi*. *Perfumery and Essential Oil Record*, 58, 154-158.
- [12] Ruecker G, Kretzschmar U. (1971) 9-Aristolene-1- $\alpha$ -ol and 1,2,9,10-tetrahydroaristolane, new aristolane-type sesquiterpenes. *Justus Liebigs Annalen der Chemie*, 748, 214-217.
- [13] Shide L, Olbrich A, Mayer R, Ruecker G. (1987) Gansongone, a new aristolane ketone from *Nardostachys chinensis* and structure revision of an aristolene. *Planta Medica*, 53, 556-558.
- [14] Han Y, Xiao D, Xiang Y, Ye L, Cheng C. (2000) Study on the volatile oil of *Nardostachys chinensis*. *Zhong yao cai*, 23, 34-35.
- [15] Mahalwal VS, Ali M. (2002) Volatile constituents of the rhizomes of *Nardostachys jatamansi* DC. *Journal of Essential Oil-Bearing Plants*, 5, 83-89.
- [16] Buechi G, Greuter F, Tokoroyama T. (1962) Terpenes. XVII. Structure of calarene and stereochemistry of aristolone. *Tetrahedron Letters*, 827-833.
- [17] Costa R, De Fina MR, Valentino MR, Dugo P, Mondello L. (2007) Reliable identification of terpenoids and related compounds by using linear retention indices interactively with mass spectrometry search. *Natural Product Communications*, 2, 413-418.
- [18] Ruecker G. (1968) 1(10)-Aristolene-2-one and 1,8,9,10-tetrahydroaristolane-2-one, new sesquiterpene ketones of the aristolane type. *Justus Liebigs Annalen der Chemie*, 717, 221-224.
- [19] Ruecker G. (1969) Components of commercial Chinese spikenard oil (*Nardostachys jatamansi*). *Planta Medica*, 17, 32-34.
- [20] Ruecker G, Shide L, Olbrich A. (1990) Peroxides as plant constituents. VII. Formation of the sesquiterpene peroxide nardosinone from an aristolane precursor. *Archiv der Pharmazie*, 323, 171-175.

- [21] Itokawa H, Masuyama K, Morita H, Takeya K. (1993) Cytotoxic sesquiterpenes from *Nardostachys chinensis*. *Chemical and Pharmaceutical Bulletin*, **41**, 1183-1184.
- [22] Ruecker G. (1971) 9-hydroxy-delta-1(10)-aristolene-2-one (dehilon) from *Nardostachys chinensis*. *Planta Medica*, **19**, 16-18.
- [23] Zhang Y, Lu Y, Zhang L, Zheng Q-T, Xu L-Z, Yang S-L. (2005) Terpenoids from the Roots and Rhizomes of *Nardostachys chinensis*. *Journal of Natural Products*, **68**, 1131-1133.
- [24] Tanitsu M, Takaya Y, Akasaka M, Niwa M, Oshima Y. (2002) Guaiane-and aristolane-type sesquiterpenoids of *Nardostachys chinensis* roots. *Phytochemistry*, **59**, 845-849.
- [25] Schulte KE, Ruecker G, Glauch G. (1967) Some components of the subterranean parts of *Nardostachys chinensis*. *Planta Medica*, **15**, 274-281.
- [26] Hoerster H, Ruecker G, Tautges J. (1977) Valeranone content in the roots of *Nardostachys jatamansi* and *Valeriana officinalis*. *Phytochemistry*, **16**, 1070-1071.
- [27] Shanbhag SN, Mesta CK, Maheshwari ML, Paknikar SK, Bhattacharyya SC. (1964) Terpenoids. LII. Jatamansin, a new terpenic coumarin from *Nardostachys jatamansi*. *Tetrahedron*, **20**, 2605-2615.
- [28] Naves YR. (1963) Essential oils. CLXXXVII. Presence of maaliol and valeranone in oil of *Nardostachys jatamansi*. *Helvetica Chimica Acta*, **46**, 2139-2141.
- [29] Sastry SD, Maheshwari ML, Chakravarti KK, Bhattacharyya SC. (1967) Terpenoids. CVI. The structure of calarenol. *Tetrahedron*, **23**, 1997-2000.
- [30] Bagchi A, Oshima Y, Hikino H. (1988) The validity of oriental medicines. Part 127. Sesquiterpenoids. Part 62. Kanshone C, a sesquiterpenoid of *Nardostachys chinensis* roots. *Phytochemistry*, **27**, 2877-2879.
- [31] Bagchi A, Oshima Y, Hikino H. (1991) Jatamols A and B: sesquiterpenoids of *Nardostachys jatamansi* roots. *Planta Medica*, **57**, 282-283.
- [32] Shanbhag SN, Mesta CK, Maheshwari ML, Bhattacharyya SC. (1965) Terpenoids. LXXV. Constituents of *Nardostachys jatamansi* and synthesis of ( $\pm$ )-dihydrosamidin and visnadin from jatamansin. *Tetrahedron*, **21**, 3591-3597.
- [33] Masuyama K, Morita H, Takeya K, Itokawa H. (1993) Eudesm-11-en-2,4- $\alpha$ -diol from *Nardostachys chinensis*. *Phytochemistry*, **34**, 567-568.
- [34] Takaya Y, Akasaka M, Takeuji Y, Tanitsu M-a, Niwa M, Oshima Y. (2000) Novel Guaianoids, Nardoguaianone E-I, from *Nardostachys chinensis* Roots. *Tetrahedron*, **56**, 7679-7683.
- [35] Takaya Y, Kurumada KI, Takeuji Y, Kim HS, Shibata Y, Ikemoto N, Wataya Y, Oshima Y. (1998) Novel antimalarial guaiane-type sesquiterpenoids from *Nardostachys chinensis* roots. *Tetrahedron Letters*, **39**, 1361-1364.
- [36] Takaya Y, Takeuji Y, Akasaka M, Nakagawasai O, Tadano T, Kisara K, Kim HS, Wataya Y, Niwa M, Oshima Y. (2000) Novel Guaiane Endoperoxides, Nardoguaianone A-D, from *Nardostachys chinensis* Roots and their Antinociceptive and Antimalarial Activities. *Tetrahedron*, **56**, 7673-7678.
- [37] Kawashima A, Kishimoto M, Morimoto S, Akyama T, Maejima A, Kawada I. (1996) Extraction of novel compound BR-606 from *Nardostachys jatamansi* roots as bone sorption inhibitor for treatment of osteoporosis and hypercalcemia. *Japan Kokai Tokkyo Koho*, 5 pp.
- [38] Luo S, Zhang Q. (1997) Nardosaldehyde: new peroxide-sesquiterpene. *Tianran Chanwu Yanjiu Yu Kaifa*, **9**, 7-9.
- [39] Ruecker G, Tautges J. (1974) Structure of jatamansic acid. *Archiv der Pharmazie*, **307**, 791-795.
- [40] Ruecker G, Tautges J, Sieck A, Wenzl H, Graf E. (1978) Studies on isolation and pharmacodynamic activity of the sesquiterpene valeranone from *Nardostachys jatamansi* DC. *Arzneimittel-Forschung*, **28**, 7-13.
- [41] Govindachari TR, Pai BR, Purushothaman KK, Rajadurai S. (1959) Jatamansone. *Tetrahedron Letters*, **15**, 5-9.
- [42] Ruecker G. (1975) Structure of nardosinone. VII. Configuration of nardosinone. *Justus Liebigs Annalen der Chemie*, 311-315.
- [43] Govindachari TR, Rajadurai S, Pai BR. (1958) Structure of jatamanson. I. *Chemische Berichte*, **91**, 908-910.
- [44] Arora RB, Sharma PL, Kapila K. (1958) Antiarrhythmic and anticonvulsant activity of jatamansone. *Indian Journal of Medical Research*, **46**, 782-791.
- [45] Chaudhry GR, Sharma VN, Siddiqui S. (1951) Chemical constituents of *Nardostachys jatamansi*. I. Isolation of a crystalline acid and an essential oil. *Journal of Scientific and Industrial Research*, **10B**, 48.
- [46] Chaudhry GR, Dhar MM, Anand N, Dhar ML (1958) Structure of jatamansic acid. *Journal of Scientific and Industrial Research (India)*, **17B**, 159-162.
- [47] Sastry SD, Maheshwari ML, Bhattacharyya SC. (1966) Terpenoids. LXXXVII. Structure of nardol. *Tetrahedron Letters*, **10**, 1035-1042.
- [48] Chatterjee A, Basak B, Saha M, Dutta U, Mukhopadhyay C, Banerji J, Konda Y, Harigaya Y. (2000) Structure and Stereochemistry of Nardostachysin, a New Terpenoid Ester Constituent of the Rhizomes of *Nardostachys jatamansi*. *Journal of Natural Products*, **63**, 1531-1533.
- [49] Bagchi A, Oshima Y, Hikino H. (1988) The validity of the oriental medicines. Part 130. Sesquiterpenoids. Part 63. Kanshones D and E, sesquiterpenoids of *Nardostachys chinensis* roots. *Phytochemistry*, **27**, 3667-3669.

- [50] Ruecker G. (1970) Structure of nardosinone. IV. Constitution of isonardosinone. *Justus Liebigs Annalen der Chemie*, **733**, 152-158.
- [51] Shide L, Mayer R, Ruecker G. (1987) Nardonoxide, a new nardosinane-type sesquiterpene ether from *Nardostachys chinensis*. *Planta Medica*, **53**, 332-334.
- [52] Bagchi A, Oshima Y, Hikino H. (1988) Sesquiterpenoids. Part 61. Kanshones A and B, sesquiterpenoids of *Nardostachys chinensis*. *Phytochemistry*, **27**, 1199-1220.
- [53] Schulte KE, Glauch G, Rucker G. (1965) Nardosinone, a new constituent of *Nardostachys chinensis* Batalin. *Tetrahedron Letters*, **35**, 3083-3084.
- [54] Tori M, Furuta H, Asakawa Y. (1991) Total synthesis of kanshone A, a sesquiterpene isolated from *Nardostachys chinensis* (Valerianaceae). *Journal of the Chemical Society, Perkin Transaction I: Organic and Bio-organic Chemistry*, **8**, 1919-1922.
- [55] Hikino H, Hikino Y, Koakutsu S, Takemoto T. (1972) Sesquiterpenoids. XLV. Structure and absolute configuration of narchinol A. *Phytochemistry*, **11**, 2097-2099.
- [56] Saunders WD, Pinder AR. (1977) Further studies relating to the structure of nardostachone. *Tetrahedron Letters*, **20**, 1687-1690.
- [57] Ruecker G, Tautges J, Maheswari ML, Saxena DB. (1976) Norseychelanone,  $\alpha$  - and  $\beta$  -patchoulenes and patchouli alcohol from *Nardostachys jatamansi*. *Phytochemistry*, **15**, 224.
- [58] Rucker G, Kretzschmar U. (1972) Patchouli alcohol and beta patchoulene from *Nardostachys chinensis*. *Planta Medica*, **21**, 1-4.
- [59] Huang He, Forsyth CJ. (1995) Anti Selective Spirocarbomercuration: Synthesis and Stereochemistry of the Spirobicyclic Sesquiterpenes Spirojatamol and Erythrodiene. *Journal of Organic Chemistry*, **60**, 2773-2779.
- [60] Bagchi A, Oshima Y, Hikino H. (1990) Sesquiterpenoids. Part 64. Spirojatamol, a new skeletal sesquiterpenoid of *Nardostachys jatamansi* roots. *Tetrahedron*, **46**, 1523-1530.
- [61] Maheswari ML, Saxena DB. (1974) Seychellene and seychelane from *Nardostachys jatamansi*. *Indian Journal of Chemistry*, **12**, 1221-1222.
- [62] Chatterjee A, Basak B, Dutta U, Banerji J, Neuman A, Prangé T. (2005) Studies on the chemical constituents of *Nardostachys jatamansi* DC (Valerianaceae). *Indian Journal of Chemistry*, **44B**, 430-433.
- [63] Bagchi A, Oshima Y, Hikino H. (1988) Validity of the Oriental medicines. Part 123. Nardostachin, an iridoid of *Nardostachys chinensis*. *Planta Medica*, **54**, 87-88.
- [64] Bagchi A, Oshima Y, Hikino H. (1991) Neolignans and lignans of *Nardostachys jatamansi* roots. *Planta Medica*, **57**, 96-97.
- [65] Hirose Y, Yonemitsu K, Sonoda T. (1978) Chemical studies on the components of *Nardostachys jatamansi* de Candolle I. *Shoyakugaku Zasshi*, **32**, 121-122.
- [66] Singh V, Ali M. (2003) New phytoconstituents from *Nardostachys jatamansi* rhizomes. *Journal of Saudi Chemical Society*, **7**, 119-128.
- [67] Mody SB, Mody PD, Vasavada SA. (1997) An improved process for extraction of the Ayurvedic plant jatamansi (*Nardostachys jatamansi*), (J.B. Chemicals & Pharmaceuticals Ltd., India), 13pp.
- [68] Benveniste H. (1991) The excitotoxin hypothesis in relation to cerebral ischemia. *Cerebrovascular and Brain Metabolism Reviews*, **3**, 213-245.
- [69] Choi DW. (1988) Glutamate neurotoxicity and diseases of the nervous system. *Neuron*, **1**, 623-634.
- [70] Coyle JT, Puttfarcken P. (1993) Oxidative stress, glutamate, and neurodegenerative disorders. *Science*, **262**, 689-695.
- [71] Hamied KA, Bakshi VM, Aghara LP. (1962) Pharmacological investigation of *Nardostachys jatamansi* root. *Journal of Scientific and Industrial Research*, Section B: Physical Sciences, **21c**, 100-103.
- [72] Prabhu V, Karanth KS, Rao A. (1994) Effects of *Nardostachys jatamansi* on biogenic amines and inhibitory amino acids in the rat brain. *Planta Medica*, **60**, 114-117.
- [73] Houghton PJ. (1999) The scientific basis for the reputed activity of Valerian. *Journal of Pharmacy and Pharmacology*, **51**, 505-512.
- [74] Fridovich I. (1978) The biology of oxygen radicals. *Science*, **201**, 875-880.
- [75] Adlard BP, deSouza SW, Moon S. (1973) The effect of age, growth retardation and asphyxia on ascorbic acid concentrations in developing brain. *Journal of Neurochemistry*, **21**, 877-881.
- [76] Folbergrova J, Rehnrona S, Siesjoe BK. (1979) Oxidized and reduced glutathione in the rat brain under normoxic and hypoxic conditions. *Journal of Neurochemistry*, **32**, 1621-1627.
- [77] Salim S, Ahmad M, Zafar KS, Ahmad AS, Islam F. (2003) Protective effect of *Nardostachys jatamansi* in rat cerebral ischemia. *Pharmacology, Biochemistry and Behavior*, **74**, 481-486.
- [78] Agrawal SS, Sharma RC, Arora RB. (1973) Antioestrogenic activity of jatamansone semicarbazone. *Indian Journal of Experimental Biology*, **11**, 583.