

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Mathematical and Statistical Sciences Faculty
Publications and Presentations

College of Sciences

2020

OPTIMAL QUANTIZATION FOR DISCRETE DISTRIBUTIONS

Russel Cabasag

The University of Texas Rio Grande Valley

Samir Huq

The University of Texas Rio Grande Valley

Eric Mendoza

The University of Texas Rio Grande Valley

Mrinal Kanti Roychowdhury

The University of Texas Rio Grande Valley

Follow this and additional works at: https://scholarworks.utrgv.edu/mss_fac



Part of the [Mathematics Commons](#)

Recommended Citation

Cabasag, Russel, Samir Huq, Eric Mendoza, and Mrinal Kanti Roychowdhury. 2020. "Optimal Quantization for Discrete Distributions." ArXiv:2008.03255 [Math], August. <http://arxiv.org/abs/2008.03255>.

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

OPTIMAL QUANTIZATION FOR DISCRETE DISTRIBUTIONS

¹RUSSEL CABASAG, ²SAMIR HUQ, ³ERIC MENDOZA, AND ⁴MRINAL KANTI ROYCHOWDHURY

ABSTRACT. In this paper, we first determine the optimal sets of n -means and the n th quantization errors for all $1 \leq n \leq 6$ for two nonuniform discrete distributions with support the set $\{1, 2, 3, 4, 5, 6\}$. Then, for a probability distribution P with support $\{\frac{1}{n} : n \in \mathbb{N}\}$ associated with a mass function f , given by $f(x) = \frac{1}{2^k}$ if $x = \frac{1}{k}$ for $k \in \mathbb{N}$, and zero otherwise, we determine the optimal sets of n -means and the n th quantization errors for all positive integers up to $n = 300$. Further, for a probability distribution P with support the set \mathbb{N} of natural number associated with a mass function f , given by $f(x) = \frac{1}{2^k}$ if $x = k$ for $k \in \mathbb{N}$, and zero otherwise, we determine the optimal sets of n -means and the n th quantization errors for all positive integers n . At last we discuss for a discrete distribution, if the optimal sets are given, how to obtain the probability distributions.

1. INTRODUCTION

Quantization is the process of converting a continuous analog signal into a digital signal of k discrete levels, or converting a digital signal of n levels into another digital signal of k levels, where $k < n$. It is essential when analog quantities are represented, processed, stored, or transmitted by a digital system, or when data compression is required. It is a classic and still very active research topic in source coding and information theory. It has broad applications in engineering and technology (see [GG, GN, Z]). For mathematical treatment of quantization one is referred to Graf-Luschgy's book (see [GL]). Let \mathbb{R}^d denote the d -dimensional Euclidean space, $\|\cdot\|$ denote the Euclidean norm on \mathbb{R}^d for any $d \geq 1$, and $n \in \mathbb{N}$. Let P denote a Borel probability measure on \mathbb{R}^d . For a finite set $\alpha \subset \mathbb{R}^d$, the error $\int \min_{a \in \alpha} \|x - a\|^2 dP(x)$ is often referred to as the *cost* or *distortion error* for α , and is denoted by $V(P; \alpha)$. For any positive integer n , write $V_n := V_n(P) = \inf\{V(P; \alpha) : \alpha \subset \mathbb{R}^d, 1 \leq \text{card}(\alpha) \leq n\}$. Then, V_n is called the n th quantization error for P . Recently, optimal quantization for different uniform distributions have been investigated by several authors, for example, see [DR, R, RR, RS].

In this paper, we investigate the optimal quantization for finite, and infinite discrete distributions. In Section 3, we calculate the optimal sets of n -means and the n th quantization errors for all $1 \leq n \leq 6$ for two nonuniform discrete distributions with support $\{1, 2, 3, 4, 5, 6\}$ associated with two different probability vectors. In Section 4, first, for a probability distribution P with support $\{\frac{1}{n} : n \in \mathbb{N}\}$ associated with a mass function f , given by $f(x) = \frac{1}{2^k}$ if $x = \frac{1}{k}$ for $k \in \mathbb{N}$, and zero otherwise, we determine the optimal sets of n -means and the n th quantization errors for all positive integers up to $n = 300$. Then, for a probability distribution P with support the set \mathbb{N} of natural number associated with a mass function f , given by $f(x) = \frac{1}{2^k}$ if $x = k$ for $k \in \mathbb{N}$, and zero otherwise, we determine the optimal sets of n -means and the n th quantization errors for all positive integers n . In Section 5, we discuss for a discrete distribution, if the optimal sets are given, how to obtain the probability distributions.

2. BASIC PRELIMINARIES

Given a finite set $\alpha \subset \mathbb{R}^d$, the Voronoi region generated by $a \in \alpha$ is defined by

$$M(a|\alpha) = \{x \in \mathbb{R}^d : \|x - a\| = \min_{b \in \alpha} \|x - b\|\},$$

2010 *Mathematics Subject Classification.* 60Exx, 94A34.

Key words and phrases. Discrete distribution, optimal set of n -means, quantization error.

i.e., the Voronoi region generated by $a \in \alpha$ is the set of all elements in \mathbb{R}^d which are nearest to a , and the set $\{M(a|\alpha) : a \in \alpha\}$ is called the *Voronoi diagram* or *Voronoi tessellation* of \mathbb{R}^d with respect to α .

The following proposition is well-known (see [GG, GL]).

Proposition 2.1. Let α be an optimal set of n -means for P , and $a \in \alpha$. Then,

- (i) $P(M(a|\alpha)) > 0$, (ii) $P(\partial M(a|\alpha)) = 0$, (iii) $a = E(X : X \in M(a|\alpha))$,
where X is a random variable with distribution P .

Due to the above proposition, we see that if α is an optimal set and $a \in \alpha$, then a is the *conditional expectation* of the random variable X given that X takes values in the Voronoi region of a . In the sequel, we will denote the support of a probability distribution P by $\text{supp}(P)$. Let P be the uniform distribution defined on the set $\{1, 2, 3, 4, 5, 6\}$. Then, the random variable X associated with the probability distribution is a discrete random variable with probability mass function f given by

$$f(x) = P(X : X = x) = \frac{1}{6}, \text{ for all } x \in \{1, 2, 3, 4, 5, 6\}.$$

It is not difficult to show that if α_n is an optimal set of n -means for P , then

$$\begin{aligned} \alpha_1 &= \{3.5\}, \alpha_2 = \{2, 5\}, \alpha_3 = \{1.5, 3.5, 5.5\}, \alpha_4 = \{1.5, 3.5, 5, 6\}, \\ \alpha_5 &= \{1.5, 3, 4, 5, 6\}, \text{ and } \alpha_6 = \text{supp}(P). \end{aligned}$$

Remark 2.2. Optimal sets are not unique. For example, in the above, the set α_5 can be any one of the following sets:

$$\{1.5, 3, 4, 5, 6\}, \{1, 2.5, 4, 5, 6\}, \{1, 2, 3.5, 5, 6\}, \{1, 2, 3, 4.5, 6\}, \{1, 2, 3, 4, 5.5\}.$$

In the following sections we give our main results.

3. OPTIMAL QUANTIZATION FOR NONUNIFORM DISCRETE DISTRIBUTIONS

In this section, we determine the optimal sets of n -means for all $1 \leq n \leq 6$ for two nonuniform discrete distributions on the set $\{1, 2, 3, 4, 5, 6\}$ associated with two different probability vectors. Let X be the random variable associated with such a distribution. For $i, j \in \{1, 2, \dots, 6\}$ with $i \leq j$, write $a[i, j] := E(X : X \in \{i, i+1, \dots, j\})$. We give our results in two subsections.

3.1. Nonuniform distribution associated with the probability vector $(\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \frac{1}{2^5}, \frac{1}{2^5})$.

Let P be a nonuniform distribution defined on the set $\{1, 2, 3, 4, 5, 6\}$ with probability mass function f given by

$$f(j) = P(X : X = j) = \begin{cases} \frac{1}{2^j} & \text{if } j \in \{1, 2, 3, 4, 5\}, \\ \frac{1}{2^5} & \text{if } j = 6, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that $\text{supp}(P) = \{1, 2, 3, 4, 5, 6\}$. In this subsection, our goal is to calculate the optimal sets α_n of n -means and the n th quantization errors V_n for all $n = 1, 2, 3, 4, 5, 6$. Since

$$E(X) = \sum_{j=1}^6 jf(j) = \frac{63}{32},$$

the optimal set of one-mean is the set $\{\frac{63}{32}\}$ with quantization error the variance V of the random variable X , where

$$V = V_1 = E\|X - E(X)\|^2 = \sum_{j=1}^6 f(j)(j - \frac{63}{32})^2 = \frac{1695}{1024}.$$

Moreover, the optimal set α_6 of six-means is just the support of P , i.e., $\alpha_6 = \{1, 2, \dots, 6\}$. In the following propositions, we determine the optimal sets of n -means for $2 \leq n \leq 5$.

Proposition 3.1.1. *The optimal set of two-means is given by $\{a[1, 2], a[3, 6]\}$ with quantization error $V_2 = \frac{341}{768}$.*

Proof. Notice that $a[1, 2] = \frac{4}{3}$, and $a[3, 6] = \frac{31}{8}$. Let us consider the set $\beta := \{\frac{4}{3}, \frac{31}{8}\}$. Since $2 < \frac{1}{2}(\frac{4}{3} + \frac{31}{8}) = 2.60417 < 3$, the distortion error due to the set β is given by

$$\sum_{j=1}^6 f(j) \min_{a \in \beta} (j - a)^2 = \sum_{j=1}^2 f(j) \left(j - \frac{4}{3}\right)^2 + \sum_{j=3}^6 f(j) \left(j - \frac{31}{8}\right)^2 = \frac{341}{768}.$$

Since V_2 is the quantization error for two-means, we have $V_2 \leq \frac{341}{768} = 0.44401$. Let $\alpha := \{a_1, a_2\}$ be an optimal set of two-means. Without any loss of generality, we can assume that $1 \leq a_1 < a_2 \leq 6$. Notice that the Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 contains 3 as well. Then, as $a[1, 3] = \frac{11}{7}$, we have

$$V_2 \geq \sum_{j=1}^3 f(j) \left(j - \frac{11}{7}\right)^2 = \frac{13}{28} = 0.464286 > V_2,$$

which gives a contradiction. Hence, we can assume that the Voronoi region of a_1 does not contain 3. Next, suppose that the Voronoi region of a_1 contains only the point 1. Then, the Voronoi region of a_2 contains all the remaining points, and so

$$a_2 = a[2, 6] = \frac{47}{16},$$

implying

$$V_2 = \sum_{j=2}^6 f(j) \left(j - \frac{47}{16}\right)^2 = \frac{367}{512} = 0.716797 > V_2,$$

which yields a contradiction. Hence, we can assume that the Voronoi region of a_1 contains only the points 1 and 2, and the remaining points are contained in the Voronoi region of a_2 , implying

$$a_1 = a[1, 2] = \frac{4}{3}, \text{ and } a_2 = a[3, 6] = \frac{31}{8}$$

with quantization error $V_2 = \frac{341}{768}$. Thus, the proof of the proposition is complete. \square

Proposition 3.1.2. *The optimal set of three-means is given by $\{1, a[2, 3], a[4, 6]\}$ with quantization error $V_3 = \frac{65}{384}$.*

Proof. Notice that $a[2, 3] = \frac{7}{3}$, and $a[4, 6] = \frac{19}{4}$. The distortion error due to the set $\beta := \{1, \frac{7}{3}, \frac{19}{4}\}$ is given by

$$\sum_{j=1}^6 f(j) \min_{a \in \beta} (j - a)^2 = \sum_{j=2}^3 f(j) \left(j - \frac{7}{3}\right)^2 + \sum_{j=4}^6 f(j) \left(j - \frac{19}{4}\right)^2 = \frac{65}{384}.$$

Since V_3 is the quantization error for three-means, we have $V_3 \leq \frac{65}{384} = 0.169271$. Let $\alpha := \{a_1, a_2, a_3\}$ be an optimal set of three-means such that $1 \leq a_1 < a_2 < a_3 \leq 6$. Notice that the Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 also contains 3. Then,

$$V_3 \geq \sum_{j=1}^3 f(j) (j - a[1, 3])^2 = \frac{13}{28} > V_3,$$

which yields a contradiction. Thus, we can assume that the Voronoi region of a_1 does not contain 3. Suppose that the Voronoi region of a_1 contains only the two points 1 and 2. Then, the Voronoi region of a_2 must contain 3. The following two case can arise:

Case 1. The Voronoi region of a_2 does not contain 4.

Then, we must have $a_2 = 3$, and $a_4 = a[4, 6]$, yielding

$$V_3 \geq \sum_{j=1}^2 f(j)(j - a[1, 2])^2 + \sum_{j=4}^6 f(j)(j - a[4, 6])^2 = \frac{97}{384} = 0.252604 > V_3,$$

which is a contradiction.

Case 2. The Voronoi region of a_2 contains 4.

Then,

$$V_3 \geq \sum_{j=1}^2 f(j)(j - a[1, 2])^2 + \sum_{j=3}^4 f(j)(j - a[3, 4])^2 = \frac{5}{24} = 0.208333 > V_3,$$

which leads to a contradiction.

Hence, by Case 1 and Case 2, we can assume that the Voronoi region of a_1 contains only the point 1, i.e., $a_1 = 1$. Then, the Voronoi region of a_2 must contain 2. Suppose that the Voronoi region of a_2 also contains 4. Then,

$$V_3 \geq \sum_{j=2}^4 f(j)(j - a[2, 4])^2 = \frac{13}{56} = 0.232143 > V_3,$$

which yields a contradiction. Thus, we can assume that the Voronoi region of a_2 does not contain 4. Suppose that the Voronoi region of a_2 contains only the point 2. Then, the Voronoi region of a_3 must contain the remaining points, which yields

$$V_3 \geq \sum_{j=3}^6 f(j)(j - a[3, 6])^2 = \frac{71}{256} = 0.277344 > V_3,$$

which is a contradiction. Hence, we can assume that the Voronoi region of a_2 contains only the two points 2 and 3, implying the fact that the Voronoi region of a_3 contains the points 4, 5, and 6. Thus, we have

$$a_1 = 1, a_2 = a[2, 3] = \frac{7}{3}, \text{ and } a_3 = a[4, 6] = \frac{19}{4},$$

with quantization error $V_3 = \frac{65}{384}$, which yields the proposition. \square

Proposition 3.1.3. *The optimal set of four-means is $\{1, 2, a[3, 4], a[5, 6]\}$ with quantization error $V_4 = \frac{11}{192}$.*

Proof. The distortion error due to the set $\beta := \{1, 2, a[3, 4], a[5, 6]\}$ is given by

$$\sum_{j=1}^6 f(j) \min_{a \in \beta} (j - a)^2 = \sum_{j=3}^4 f(j)(j - a[3, 4])^2 + \sum_{j=5}^6 f(j)(j - a[5, 6])^2 = \frac{11}{192}.$$

Since V_4 is the quantization error for four-means, we have $V_4 \leq \frac{11}{192} = 0.0572917$. Let $\alpha := \{a_1, a_2, a_3, a_4\}$ be an optimal set of four-means. Without any loss of generality, we can assume that $1 \leq a_1 < a_2 < a_3 < a_4 \leq 6$. The Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 contains 2 as well. Then,

$$V_4 \geq \sum_{j=1}^2 f(j)(j - a[1, 2])^2 = \frac{1}{6} > V_4,$$

which gives a contradiction. Hence, we can assume that the Voronoi region of a_1 contain the point 1 only, i.e., $a_1 = 1$. Then, the Voronoi region of a_2 must contain 2. Suppose that the Voronoi region of a_2 also contains 3. Then,

$$V_4 \geq \sum_{j=2}^3 f(j)(j - a[2, 3])^2 = \frac{1}{12} = 0.0833333 > V_4,$$

which leads to a contradiction. Hence, the Voronoi region of a_2 does not contain 3, i.e., $a_2 = 2$. Then, the Voronoi region of a_3 must contain 3. Suppose that the Voronoi region of a_3 contains 5 as well. Then, we have

$$V_4 \geq \sum_{j=3}^5 f(j)(j - a[3, 5])^2 = \frac{13}{112} = 0.116071 > V_4,$$

which yields a contradiction. Thus, we can assume that the Voronoi region of a_3 does not contain 5. Suppose that the Voronoi region of a_3 contains 3 only. Then, the Voronoi region of a_5 contains 4, 5, 6, which implies

$$V_4 = \sum_{j=4}^6 f(j)(j - a[4, 6])^2 = \frac{11}{128} = 0.0859375 > V_4,$$

which gives a contradiction. Hence, the Voronoi region of a_3 contains 3 and 4, yielding $a_3 = a[3, 4]$, and $a_4 = a[5, 6]$. Thus, the optimal set of four-means is $\{1, 2, a[3, 4], a[5, 6]\}$ with quantization error $V_4 = \frac{11}{192}$. which is the proposition. \square

Using the similar technique as the previous proposition, the following proposition can be proved.

Proposition 3.1.4. *The optimal set of five-means is $\{1, 2, 3, 4, a[5, 6]\}$ with quantization error $V_5 = \frac{1}{64}$.*

3.2. Nonuniform distribution associated with a probability vector of the form $(x, (1-x)x, (1-x)^2x, (1-x)^3x, (1-x)^4x, (1-x)^5)$. Let P be a nonuniform distribution defined on the set $\{1, 2, 3, 4, 5, 6\}$ with probability mass function f given by

$$f(j) = P(X : X = j) = \begin{cases} x & \text{if } j = 1, \\ (1-x)^{j-1}x & \text{if } j \in \{2, 3, 4, 5\}, \\ (1-x)^5 & \text{if } j = 6, \\ 0 & \text{otherwise,} \end{cases}$$

where $0 < x < 1$. Notice that $\text{supp}(P) = \{1, 2, 3, 4, 5, 6\}$. Fix $x = \frac{7}{10}$. In this subsection, our goal is to calculate the optimal sets α_n of n -means and the n th quantization errors for all $n = 1, 2, 3, 4, 5, 6$ for the given mass function f with $x = \frac{7}{10}$. Since

$$E(X) = \sum_{j=1}^6 jf(j) = \frac{142753}{100000},$$

the optimal set of one-mean is the set $\{\frac{142753}{100000}\}$ with quantization error the variance V of the random variable X , where

$$V = V_1 = E\|X - E(X)\|^2 = \sum_{j=1}^6 f(j) \left(j - \frac{142753}{100000} \right)^2 = \frac{6007880991}{10000000000}.$$

Moreover, the optimal set α_6 of six-means is just the support of P , i.e., $\alpha_6 = \{1, 2, \dots, 6\}$. In the following propositions, we determine the optimal sets of n -means for $2 \leq n \leq 5$.

Proposition 3.2.1. *The optimal set of two-means is given by $\{1, a[2, 6]\}$ with quantization error $V_2 = \frac{174296997}{1000000000}$.*

Proof. The distortion error due to the set $\beta := \{1, a[2, 6]\}$ is given by

$$\sum_{j=2}^6 f(j) \min_{a \in \beta} (j - a)^2 = \sum_{j=2}^6 f(j) (j - a[2, 6])^2 = \frac{174296997}{1000000000}.$$

Since V_2 is the quantization error for two-means, we have $V_2 \leq \frac{174296997}{1000000000} = 0.174296997$. Let $\alpha := \{a_1, a_2\}$ be an optimal set of two-means. Without any loss of generality, we can assume that $1 \leq a_1 < a_2 \leq 6$. Notice that the Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 contains 3 as well. Then,

$$V_2 \geq \sum_{j=1}^3 f(j)(j - a[1, 3])^2 = \frac{4809}{13900} = 0.345971 > V_2,$$

which gives a contradiction. Hence, we can assume that the Voronoi region of a_1 does not contain 3. Next, suppose that the Voronoi region of a_1 contains 2. Then, the Voronoi region of a_2 contains all the remaining points, and so

$$V_2 = \sum_{j=1}^2 f(j)(j - a[1, 2])^2 + \sum_{j=3}^6 f(j)(j - a[3, 6])^2 = \frac{272139987}{1300000000} = 0.209338 > V_2,$$

which yields a contradiction. Hence, we can assume that the Voronoi region of a_1 contains only the point 1, and the remaining points are contained in the Voronoi region of a_2 , implying

$$a_1 = 1, \text{ and } a_2 = a[2, 6]$$

with quantization error $V_2 = \frac{174296997}{1000000000}$. Thus, the proof of the proposition is complete. \square

Proposition 3.2.2. *The optimal set of three-means is given by $\{1, 2, a[3, 6]\}$ with quantization error $V_3 = \frac{4779999}{1000000000}$.*

Proof. The distortion error due to the set $\beta := \{1, 2, a[3, 6]\}$ is given by

$$\sum_{j=3}^6 f(j) \min_{a \in \beta} (j - a)^2 = \sum_{j=3}^6 f(j)(j - a[3, 6])^2 = \frac{4779999}{1000000000} = 0.04779999.$$

Since V_3 is the quantization error for three-means, we have $V_3 \leq 0.04779999$. Let $\alpha := \{a_1, a_2, a_3\}$ be an optimal set of three-means such that $1 \leq a_1 < a_2 < a_3 \leq 6$. Notice that the Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 also contains 2. Then,

$$V_3 \geq \sum_{j=1}^2 f(j)(j - a[1, 2])^2 = \frac{21}{130} = 0.161538 > V_3,$$

which yields a contradiction. Thus, we can assume that the Voronoi region of a_1 contains only the point 1, i.e., $a_1 = 1$. The Voronoi region of a_2 contains 2. Suppose that the Voronoi region of a_2 also contains 3. Then,

$$V_3 \geq \sum_{j=2}^3 f(j)(j - a[2, 3])^2 = \frac{63}{1300} = 0.0484615 > V_3,$$

which is a contradiction. Hence, the Voronoi region of a_2 contains only the point 2, which yields $a_2 = 2$, and $a_3 = a[3, 6]$, with quantization error $V_3 = \frac{4779999}{1000000000}$. Thus, the proof of the proposition is complete. \square

Following the similar techniques as given in Proposition 3.2.2, we can prove the following two propositions.

Proposition 3.2.3. *The optimal set of four-means is given by $\{1, 2, 3, a[4, 6]\}$ with quantization error $V_4 = \frac{112833}{10000000}$.*

Proposition 3.2.4. *The optimal set of five-means is given by $\{1, 2, 3, 4, a[5, 6]\}$ with quantization error $V_5 = \frac{1701}{10000000}$.*

4. OPTIMAL QUANTIZATION FOR INFINITE DISCRETE DISTRIBUTIONS

In this section, for $n \in \mathbb{N}$, we investigate the optimal sets of n -means for two different infinite discrete distributions. We give them in the following two subsections.

4.1. Optimal quantization for an infinite discrete distribution with support $\{\frac{1}{n} : n \in \mathbb{N}\}$. Let $\mathbb{N} := \{1, 2, 3, \dots\}$ be the set of natural numbers, and let P be a Borel probability measure on the set $\{\frac{1}{n} : n \in \mathbb{N}\}$ with probability mass function f given by

$$f(x) = \begin{cases} \frac{1}{2^k} & \text{if } x = \frac{1}{k} \text{ for } k \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Then, P is a Borel probability measure on \mathbb{R} , and the support of P is given by $\text{supp}(P) = \{\frac{1}{n} : n \in \mathbb{N}\}$. In this section, our goal is to determine the optimal sets of n -means and the n th quantization errors for all positive integers n for the probability measure P . For $k, \ell \in \mathbb{N}$, where $k \leq \ell$, write

$$[k, \ell] := \{\frac{1}{n} : n \in \mathbb{N} \text{ and } k \leq n \leq \ell\}, \text{ and } [k, \infty) := \{\frac{1}{n} : n \in \mathbb{N} \text{ and } n \geq k\}.$$

Further, write

$$Av[k, \ell] := E\left(X : X \in [k, \ell]\right) = \frac{\sum_{n=k}^{\ell} \frac{1}{2^n} \frac{1}{n}}{\sum_{n=k}^{\ell} \frac{1}{2^n}}, \quad Av[k, \infty) := E\left(X : X \in [k, \infty)\right) = \frac{\sum_{n=k}^{\infty} \frac{1}{2^n} \frac{1}{n}}{\sum_{n=k}^{\infty} \frac{1}{2^n}},$$

$$Er[k, \ell] := \sum_{n=k}^{\ell} \frac{1}{2^n} \left(\frac{1}{n} - Av[k, \ell]\right)^2, \text{ and } Er[k, \infty) := \sum_{n=k}^{\infty} \frac{1}{2^n} \left(\frac{1}{n} - Av[k, \infty)\right)^2.$$

Notice that $E(X) := E(X : X \in \text{supp}(P)) = \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{1}{n} = Av[1, \infty) = \log(2)$, and so the optimal set of one-mean is the set $\{\log(2)\}$ with quantization error

$$V(P) = \sum_{n=1}^{\infty} \frac{1}{2^n} \left(\frac{1}{n} - \log(2)\right)^2 = Er[1, \infty) = \frac{1}{12} (\pi^2 - 18 \log^2(2)) = 0.101788.$$

Proposition 4.1.1. *The set $\{Av[2, \infty), 1\}$ forms the optimal set of two-means for the probability measure P with quantization error $V_2(P) = Er[2, \infty) = \frac{1}{12} (\pi^2 - 12 - 30 \log^2(2) + 24 \log(2)) = 0.0076288597$.*

Proof. Consider the set $\beta := \{Av[2, \infty), 1\}$. Since $\frac{1}{3} < \frac{1}{2}(Av[2, \infty) + 1) < 1$, the Voronoi region of 1 contains only the point 1, and the Voronoi region of $Av[2, \infty)$ contains the set $\{\frac{1}{n} : n \geq 2\}$. Hence, the distortion error due to the set β is given by

$$V(P; \beta) = Er[2, \infty) = \frac{1}{12} (\pi^2 - 12 - 30 \log^2(2) + 24 \log(2)) = 0.0076288597.$$

Since $V_2(P)$ is the quantization error for two-means, we have $V_2(P) \leq 0.0076288597$. Let $\alpha := \{a_2, a_1\}$ be an optimal set of two-means. Due to Proposition 2.1, we can assume that $0 \leq a_2 < a_1 \leq 1$. The Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 also contains $\frac{1}{2}$. Then,

$$V_2(P) \geq Er[1, 2] = \frac{1}{24} = 0.0416667 > V_2(P),$$

which leads to a contradiction. Hence, we can assume that the Voronoi region of a_1 does not contain $\frac{1}{2}$. Again, by Proposition 2.1, the Voronoi region of a_2 cannot contain the point 1. Thus, we have $a_2 = Av[2, \infty)$, and $a_1 = 1$, and the corresponding quantization error is $V_2(P) = Er[2, \infty) = 0.0076288597$. Thus, the proof of the proposition is complete. \square

Proposition 4.1.2. *The set $\{Av[3, \infty), \frac{1}{2}, 1\}$ forms the optimal set of three-means for the probability measure P with quantization error $V_3(P) = Er[3, \infty) = 0.00116437359$.*

Proof. Consider the set $\beta := \{Av[3, \infty), \frac{1}{2}, 1\}$. Since, $\frac{1}{3} < \frac{1}{2}(Av[3, \infty) + \frac{1}{2}) < \frac{1}{2}$, and $\frac{1}{2} < \frac{1}{2}(\frac{1}{2} + 1) < 1$, the distortion error due to the set $\beta := \{Av[3, \infty), \frac{1}{2}, 1\}$ is given by

$$V(P; \beta) = Er[3, \infty) = \frac{1}{24} (2\pi^2 - 51 - 108 \log^2(2) + 120 \log(2)) = 0.00116437359.$$

Since $V_3(P)$ is the quantization error for three-means, we have $V_3(P) \leq 0.00116437359$. Let $\alpha := \{a_3, a_2, a_1\}$ be an optimal set of three-means such that $0 \leq a_3 < a_2 < a_1 \leq 1$. Proceeding as Proposition 4.1.1, we can show that $a_1 = 1$. Suppose that the Voronoi region of a_2 contains $\frac{1}{2}$ and $\frac{1}{3}$. Then,

$$V_3(P) \geq Er[2, 3] = 0.002314814815 > V_3(P),$$

which is a contradiction. Hence, the Voronoi region of a_2 cannot contain $\frac{1}{3}$. Thus, we have

$$a_3 = Av[3, \infty), a_2 = \frac{1}{2}, \text{ and } a_1 = 1,$$

with quantization error $V_3(P) = Er[3, \infty) = 0.00116437359$. Thus, the proof of the proposition is complete. \square

Proposition 4.1.3. *The set $\{Av[4, \infty), \frac{1}{3}, \frac{1}{2}, 1\}$ forms the optimal set of four-means for the probability measure P with quantization error $V_4(P) = Er[4, \infty) = 0.0002418966477$.*

Proof. The proof of this proposition is similar to the proof of Proposition 4.1.2. \square

Proposition 4.1.4. *The set $\{Av[5, \infty), \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1\}$ forms the optimal set of five-means for the probability measure P with quantization error $V_5(P) = Er[5, \infty) = 0.00005991266593$.*

Proof. The distortion error due to the set $\beta := \{Av[5, \infty), \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1\}$ is given by

$$V(P; \beta) := Er[5, \infty) = Er[5, \infty) = 0.00005991266593.$$

Since $V_5(P)$ is the quantization error for five-means, we have $V_5(P) \leq 0.00005991266593$. Let $\alpha := \{a_5, a_4, a_3, a_2, a_1\}$ be an optimal set of five-means such that $0 \leq a_5 < a_4 < a_3 < a_2 < a_1 \leq 1$. Proceeding as Proposition 4.1.2, we can show that $a_1 = 1$, $a_2 = \frac{1}{2}$, and $a_3 = \frac{1}{3}$. We now show that $a_4 = \frac{1}{4}$. Suppose that the Voronoi region of a_4 contains $\frac{1}{4}$, $\frac{1}{5}$, and $\frac{1}{6}$. Then,

$$V_5(P) \geq Er[4, 6] = 0.0001116071429 > V_5(P),$$

which is a contradiction. Assume that the Voronoi region of a_4 contains only the points $\frac{1}{4}$, and $\frac{1}{5}$. Then, the Voronoi region of a_5 contains the set $[6, \infty)$, and so we have

$$V_5(P) = Er[6, \infty) + Er[4, 5] = 0.00006872664638 > V_5(P),$$

which leads to a contradiction. Hence, we can assume that the Voronoi region of a_4 contains only the point $\frac{1}{4}$. Thus, we have $a_5 = Av[5, \infty)$, $a_4 = \frac{1}{4}$, $a_3 = \frac{1}{3}$, $a_2 = \frac{1}{2}$, and $a_1 = 1$ with quantization error $V_5(P) = Er[5, \infty) = 0.00005991266593$. Thus, the proof of the Proposition is complete. \square

Proposition 4.1.5. *The set $\{Av[7, \infty), Av[5, 6], \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1\}$ forms the optimal set of six-means for the probability measure P with quantization error*

$$V_6(P) = Er[7, \infty) + Er[5, 6] = 0.00001658886625.$$

Proof. Notice that $\frac{1}{7} = 0.142857 < \frac{1}{2}(Av[7, \infty) + Av[5, 6]) = 0.158488 < 0.166667 = \frac{1}{6}$, and $\frac{1}{5} < \frac{1}{2}(Av[5, 6] + \frac{1}{4}) < \frac{1}{4}$. Hence, the distortion error due to the set $\beta := \{Av[7, \infty), Av[5, 6], \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1\}$ is given by

$$V(P; \beta) = Er[7, \infty) + Er[5, 6] = 0.00001658886625.$$

Since $V_6(P)$ is the distortion error for six-means, we have $V_6(P) \leq 0.00001658886625$. Let $\alpha := \{a_6, a_5, a_4, a_3, a_2, a_1\}$ be an optimal set of six-means such that $0 \leq a_6 < a_5 < \dots < a_1 \leq 1$. Proceeding in the similar way as in the proof of Proposition 4.1.2, we can show that $a_3 = \frac{1}{3}$, $a_2 = \frac{1}{2}$, and $a_1 = 1$. Proceeding in the similar way as in the proof of Proposition 4.1.4, we can show

that $a_4 = \frac{1}{4}$. We now show that $a_5 = Av[5, 6]$. Notice that the Voronoi region of a_5 must contain $\frac{1}{5}$. Suppose that the Voronoi region of a_5 contains $\frac{1}{7}$ and $\frac{1}{6}$ as well. Then,

$$V_6(P) \geq Er[5, 7] = 0.00002576328150 > V_6(P),$$

which leads to a contradiction. Suppose that the Voronoi region of a_5 contains only the point $\frac{1}{5}$, i.e., $a_5 = \frac{1}{5}$. Then,

$$V_6(P) = Er[6, \infty) = 0.00001664331305 > V_6(P),$$

which yields a contradiction. Hence, we can assume that the Voronoi region of a_5 contains only the two points $\frac{1}{6}$ and $\frac{1}{5}$. Thus, we have

$$a_6 = Av[7, \infty), a_5 = Av[5, 6], a_4 = \frac{1}{4}, a_3 = \frac{1}{3}, a_2 = \frac{1}{2}, \text{ and } a_1 = 1,$$

and the quantization error is $V_6(P) = Er[7, \infty) + Er[5, 6] = 0.00001658886625$. Thus, the proof of the proposition is complete. \square

In the following proposition, we calculate the optimal set of n -means and the n th quantization error for $n = 200$.

Proposition 4.1.6. *The set $\{Av[301, \infty), Av[299, 300], \frac{1}{298}, \frac{1}{297}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ forms the optimal set of 300-means for the probability measure P with quantization error $V_{300}(P) = Er[301, \infty) + Er[299, 300] = 1.564317642582409606174128 \times 10^{-100}$.*

Proof. Notice that $\frac{1}{301} = .003322259136 < \frac{1}{2}(Av[301, \infty) + Av[299, 300]) = 0.003326047849 < 0.003333333333 = \frac{1}{300}$, and $\frac{1}{299} = 0.003344481605 < \frac{1}{2}(Av[299, 300] + \frac{1}{298}) = 0.003348235106 < 0.003355704698 = \frac{1}{298}$. Hence, the distortion error due to the set

$$\beta := \{Av[301, \infty), Av[299, 300], \frac{1}{198}, \frac{1}{197}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$$

is given by

$$V(P; \beta) = Er[301, \infty) + Er[299, 300] = 1.564317642582409606174128 \times 10^{-100}.$$

Since $V_{300}(P)$ is the distortion error for 300-means, we have

$$V_{300}(P) \leq 1.564317642582409606174128 \times 10^{-100}.$$

Let $\alpha := \{a_{300}, a_{299}, \dots, a_3, a_2, a_1\}$ be an optimal set of 300-means such that $0 \leq a_{300} < a_{299} < \dots < a_1 \leq 1$. Proceeding in the similar way as in the proof of Proposition 4.1.2, we can show that $a_{297} = \frac{1}{297}$, $a_{296} = \frac{1}{296}$, \dots , $a_3 = \frac{1}{3}$, $a_2 = \frac{1}{2}$, and $a_1 = 1$. Proceeding in the similar way as in the proof of Proposition 4.1.4, we can show that $a_{298} = \frac{1}{298}$. We now show that $a_{299} = Av[299, 300]$. The Voronoi region of a_{299} must contain $\frac{1}{299}$. Suppose that the Voronoi region of a_{299} contains $\frac{1}{i}$ for $i = 299, 300, 301, 302$. Then,

$$V_{300}(P) \geq Er[299, 302] = 1.953916208081117722202350 \times 10^{-100} > V_{300}(P),$$

which leads to a contradiction. Assume that the Voronoi region of a_{299} contains only the points $\frac{1}{i}$ for $i = 299, 300, 301$. Then,

$$V_{300}(P) = Er[302, \infty) + Er[299, 301] = 1.698521119259119376459397 \times 10^{-100} > V_{300}(P),$$

which yields a contradiction. Assume that the Voronoi region of a_{299} contains only the point $\frac{1}{299}$. Then,

$$V_{300}(P) = Er[300, \infty) = 2.345910694878821203973953 \times 10^{-100} > V_{300}(P),$$

which gives a contradiction. Hence, we can assume that the Voronoi region of a_{299} contains only the two points $\frac{1}{299}$ and $\frac{1}{300}$. Thus, we have

$$a_{300} = Av[301, \infty), a_{299} = Av[299, 300], a_{298} = \frac{1}{298}, \dots, a_4 = \frac{1}{4}, a_3 = \frac{1}{3}, a_2 = \frac{1}{2}, \text{ and } a_1 = 1,$$

and the quantization error is given by

$$V_{300}(P) = Er[301, \infty) + Er[299, 300] = 1.564317642582409606174128 \times 10^{-100}.$$

Thus, the proof of the proposition is complete. \square

We now give the following theorem.

Theorem 4.1.7. *For any positive integer n , the sets $\{Av[n, \infty), \frac{1}{n-1}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$, where $1 \leq n \leq 5$, form the optimal sets of n -means for the probability measure P with quantization errors $V_n(P) := Er[n, \infty)$. For the positive integers n , where $6 \leq n \leq 300$, the sets $\{Av[n+1, \infty), Av[n-1, n], \frac{1}{n-2}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ form the optimal sets of n -means for the probability measure P with quantization errors*

$$V_n(P) = Er[n+1, \infty) + Er[n-1, n].$$

Proof. Due to Proposition 4.1.1 through Proposition 4.1.4, it follows that for $1 \leq n \leq 5$ the sets $\{Av[n, \infty), \frac{1}{n-1}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ form the optimal sets of n -means for the probability measure P with quantization errors $V_n(P) = Er[n, \infty)$. Proceeding in the similar way as Proposition 4.1.5 and Proposition 4.1.6, we can show that for any positive integer n , where $6 \leq n \leq 300$, the sets $\{Av[n+1, \infty), Av[n-1, n], \frac{1}{n-2}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ form the optimal sets of n -means for the probability measure P with quantization errors

$$V_n(P) = Er[n+1, \infty) + Er[n-1, n].$$

Thus, we complete the proof of the theorem. \square

We now give the following remark.

Remark 4.1.8. Proceeding in the similar way, as given in the proof of Theorem 4.1.7, it can be shown that the set $\{Av[n, \infty), \frac{1}{n-1}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ also gives an optimal set of n -means for $n = 301$. It is still not known whether the sets $\{Av[n+1, \infty), Av[n-1, n], \frac{1}{n-2}, \dots, \frac{1}{3}, \frac{1}{2}, 1\}$ give the optimal sets of n -means for all positive integers $n \geq 6$. If not, then the least upper bound of $n \in \mathbb{N}$ for which such sets give the optimal sets of n -means for the probability measure P is not known yet.

4.2. Optimal quantization for an infinite discrete distribution with support $\{n : n \in \mathbb{N}\}$. Let $\mathbb{N} := \{1, 2, 3, \dots\}$ be the set of natural numbers, and let P be a Borel probability measure on the set $\{n : n \in \mathbb{N}\}$ with probability density function f given by

$$f(x) = \begin{cases} \frac{1}{2^n} & \text{if } x = n \text{ for } n \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Then, P is a Borel probability measure on \mathbb{R} , and the support of P is the set \mathbb{N} of natural numbers. In this section, our goal is to determine the optimal sets of n -means and the n th quantization errors for all positive integers n for the probability measure P . For $k, \ell \in \mathbb{N}$, where $k \leq \ell$, write

$$[k, \ell] := \{n : n \in \mathbb{N} \text{ and } k \leq n \leq \ell\}, \text{ and } [k, \infty) := \{n : n \in \mathbb{N} \text{ and } n \geq k\}.$$

Further, write

$$Av[k, \ell] := E\left(X : X \in [k, \ell]\right) = \frac{\sum_{n=k}^{\ell} \frac{n}{2^n}}{\sum_{n=k}^{\ell} \frac{1}{2^n}}, \quad Av[k, \infty) := E\left(X : X \in [k, \infty)\right) = \frac{\sum_{n=k}^{\infty} \frac{n}{2^n}}{\sum_{n=k}^{\infty} \frac{1}{2^n}},$$

$$Er[k, \ell] := \sum_{n=k}^{\ell} \frac{1}{2^n} \left(n - Av[k, \ell]\right)^2, \text{ and } Er[k, \infty) := \sum_{n=k}^{\infty} \frac{1}{2^n} \left(n - Av[k, \infty)\right)^2.$$

Notice that $E(P) := E(X : X \in \text{supp}(P)) = \sum_{n=1}^{\infty} \frac{n}{2^n} = Av[1, \infty) = 2$, and so the optimal set of one-mean is the set $\{2\}$ with quantization error

$$V(P) = \sum_{n=1}^{\infty} \frac{1}{2^n} (n-2)^2 = Er[1, \infty) = 2.$$

Proposition 4.2.1. *The optimal set of two-means is given by $\{Av[1, 2], Av[3, \infty)\}$ with quantization error $V_2 = \frac{2}{3}$.*

Proof. We see that $Av[1, 2] = \frac{4}{3}$, and $Av[3, \infty) = 4$. Since $\frac{4}{3} < \frac{1}{2}(\frac{4}{3} + 4) < 4$, the distortion error due to the set $\beta := \{\frac{4}{3}, 4\}$ is given by

$$V(P; \beta) = Er[1, 2] + Er[3, \infty) = \frac{2}{3}.$$

Since V_2 is the quantization error for two-means, we have $V_2 \leq \frac{2}{3}$. Notice that the Voronoi region of a_1 must contain 1. Suppose that the Voronoi region of a_1 contains the set $\{1, 2, 3, 4\}$. Then,

$$V_2 \geq \sum_{j=1}^4 \frac{1}{2^j} (j - Av[1, 4])^2 = Er[1, 4] = \frac{97}{120} = 0.808333 > V_2,$$

which yields a contradiction. Hence, we can assume that the Voronoi region of a_1 contains only the set $\{1, 2, 3\}$, and so the Voronoi region of a_2 contains the set $\{n : n \geq 4\}$. Then, we have

$$V_2 = Er[1, 3] + Er[4, \infty) = \frac{5}{7} = 0.714286 > V_2,$$

which is a contradiction. Next, suppose that the Voronoi region of a_1 contains only the element 1, and so the Voronoi region of a_2 contains the set $\{n : n \geq 2\}$. Then, we have

$$V_2 = Er[2, \infty) = 1 > V_2,$$

which leads to a contradiction. Hence, we can assume that the Voronoi region of a_1 contains the set $\{1, 2\}$, and so the Voronoi region of a_2 contains $\{3, 4, 5, \dots\}$, yielding $a_1 = Av[1, 2]$, and $a_2 = Av[3, \infty)$, and the corresponding quantization error is $V_2 = \frac{2}{3}$. Thus, the proof of the proposition is complete. \square

Proposition 4.2.2. *The sets $\{1, Av[2, 3], Av[4, \infty)\}$, and $\{Av[1, 2], Av[3, 4], Av[5, \infty)\}$ form two optimal sets of three-means with quantization error $V_3 = \frac{1}{3}$.*

Proof. The distortion error due to set $\beta := \{1, Av[2, 3], Av[4, \infty)\}$ is given by

$$V(P; \beta) = Er[2, 3] + Er[4, \infty) = \frac{1}{3}.$$

Notice that the distortion error due to the set $\{Av[1, 2], Av[3, 4], Av[5, \infty)\}$ is also $\frac{1}{3}$. Since V_3 is the quantization error for three-means, we have $V_3 \leq \frac{1}{3}$. Let $\alpha := \{a_1, a_2, a_3\}$ be an optimal set of three-means, where $1 \leq a_1 < a_2 < a_3 < \infty$. Suppose that the Voronoi region of a_1 contains the set $\{1, 2, 3\}$. Then,

$$V_3 \geq \sum_{j=1}^3 \frac{1}{2^j} (j - Av[1, 3])^2 = \frac{13}{28} > \frac{1}{3} > V_3,$$

which leads to a contradiction. Hence, we can assume that the Voronoi region of a_1 contains either the set $\{1\}$, or the set $\{1, 2\}$. Consider the following two cases:

Case 1. *The Voronoi region of a_1 contains only the set $\{1\}$.*

In this case, the Voronoi region of a_2 must contain the element 2. Suppose that the Voronoi region of a_2 contains the set $\{2, 3, 4, 5\}$. Then,

$$V_3 \geq \sum_{j=2}^5 \frac{1}{2^j} (j - Av[2, 5])^2 = \frac{97}{240} = 0.404167 > V_3,$$

which yields a contradiction. Assume that the Voronoi region of a_2 contains only the set $\{2, 3, 4\}$, and so the Voronoi region of a_3 contains the set $\{n : n \geq 5\}$. Then, the distortion error is

$$V_3 = Er[2, 4] + Er[5, \infty) = \frac{5}{14} = 0.357143 > V_3,$$

which gives a contradiction. Next, assume that the Voronoi region of a_2 contains only the element 2, and so the Voronoi region of a_3 contains the set $\{n : n \geq 3\}$. Then, the distortion error is

$$V_3 = Er[3, \infty) = \frac{1}{2} > V_3,$$

which is a contradiction. Hence, in this case, we can conclude that the Voronoi region of a_2 contains only the set $\{2, 3\}$, yielding $a_1 = 1$, $a_2 = Av[2, 3]$, and $a_3 = Av[4, \infty)$ with quantization error $V_3 = \frac{1}{3}$.

Case 2. The Voronoi region of a_1 contains only the set $\{1, 2\}$.

In this case, the Voronoi region of a_2 must contain the element 3. Suppose that the Voronoi region of a_2 contains the set $\{3, 4, 5, 6\}$. Then,

$$V_3 \geq \sum_{j=1}^2 \frac{1}{2^j} (j - Av[1, 2])^2 + \sum_{j=3}^6 \frac{1}{2^j} (j - Av[3, 6])^2 = \frac{59}{160} = 0.36875 > V_3,$$

which yields a contradiction. Assume that the Voronoi region of a_2 contains only the set $\{3, 4, 5\}$, and so the Voronoi region of a_3 contains the set $\{n : n \geq 6\}$. Then, the distortion error is

$$V_3 = Er[1, 2] + Er[3, 5] + Er[6, \infty) = \frac{29}{84} = 0.345238 > V_3,$$

which gives a contradiction. Next, assume that the Voronoi region of a_2 contains only the element 3, and so the Voronoi region of a_3 contains the set $\{n : n \geq 4\}$. Then, the distortion error is

$$V_3 = Er[1, 2] + Er[4, \infty) = \frac{5}{12} = 0.416667 > V_3,$$

which yields a contradiction. Hence, in this case, we can conclude that the Voronoi region of a_2 contains only the set $\{3, 4\}$, yielding $a_1 = Av[1, 2]$, $a_2 = Av[3, 4]$, and $a_3 = Av[5, \infty)$ with quantization error $V_3 = \frac{1}{3}$.

By Case 1 and Case 2, the proof of the proposition is complete. \square

We need the following lemma.

Lemma 4.2.3. *Let $n \geq 4$, and let α_n be an optimal set of n -means. Then, α_n must contain the set $\{1, 2, \dots, (n-3)\}$.*

Proof. The distortion error due to the set $\beta := \{1, 2, \dots, (n-3), (n-2), Av[n-1, n], Av[n+1, \infty)\}$ is given by

$$V(P; \beta) = Er[n-1, n] + Er[n+1, \infty) = \frac{2^{3-n}}{3}.$$

Since V_n is the quantization error for n -means, we have $V_n \leq \frac{2^{3-n}}{3}$. Let $\alpha_n := \{a_1, a_2, \dots, a_n\}$ be an optimal set of n -means such that $1 \leq a_1 < a_2 < \dots < a_n < \infty$. We show that $a_1 = 1, a_2 = 2, \dots, a_{n-3} = n-3$. Notice that the Voronoi region of a_1 must contain the element 1. Suppose that the Voronoi region of a_1 also contains the element 2. Then,

$$V_n > \sum_{j=1}^2 \frac{1}{2^j} (j - Av[1, 2])^2 = \frac{1}{6} \geq \frac{2^{3-n}}{3} \geq V_n,$$

which is a contradiction. Hence, we can conclude that the Voronoi region of a_1 contains only the element 1, yielding $a_1 = 1$. Thus, we can deduce that there exists a positive integer k , where $1 \leq k < n-3$, such that $a_1 = 1, a_2 = 2, \dots, a_k = k$. We now show that $a_{k+1} = k+1$. Notice

that the Voronoi region of a_{k+1} must contain $k + 1$. Suppose that the Voronoi region of a_{k+1} also contains the element $k + 2$. Then, as $k < n - 3$, we have

$$V_n > \sum_{j=k+1}^{k+2} \frac{1}{2^j} (j - Av[k + 1, k + 2])^2 = \frac{2^{-k-1}}{3} \geq \frac{2^{3-n}}{3} \geq V_n,$$

which is a contradiction. Hence, we can conclude that the Voronoi region of a_{k+1} contains only the element $k + 1$, yielding $a_{k+1} = k + 1$. Thus, by the Principle of Mathematical Induction, we deduce that $a_1 = 1, a_2 = 2, \dots, a_{n-3} = n - 3$. Thus, the proof of the lemma is complete. \square

Theorem 4.2.4. *Let $n \geq 4$, and let α_n be an optimal set of n -means. Then, either $\alpha_n = \{1, 2, 3, \dots, n - 3, n - 2, Av[n - 1, n], Av[n + 1, \infty)\}$, or $\alpha_n = \{1, 2, 3, \dots, n - 3, Av[n - 2, n - 1], Av[n, n + 1], Av[n + 2, \infty)\}$ with quantization error $V_n = \frac{2^{3-n}}{3}$.*

Proof. As shown in the proof of Lemma 4.2.3, we have $V_n \leq \frac{2^{3-n}}{3}$. Let $\alpha_n := \{a_1, a_2, \dots, a_n\}$ be an optimal set of n -means such that $1 \leq a_1 < a_2 < \dots < a_n < \infty$. By Lemma 4.2.3, we have $a_1 = 1, a_2 = 2, \dots, a_{n-3} = n - 3$. Recall that $n \geq 4$. Suppose that the Voronoi region of a_{n-2} contains the set $\{n - 2, n - 1, n\}$. Then,

$$V_n \geq \sum_{j=n-2}^n \frac{1}{2^j} (j - Av[n - 2, n])^2 = \frac{13}{7} 2^{1-n} > \frac{2^{3-n}}{3} \geq V_n,$$

which leads to a contradiction. Hence, we can assume that the Voronoi region of a_{n-2} contains either the set $\{n - 2\}$, or the set $\{n - 2, n - 1\}$. Consider the following two cases:

Case 1. *The Voronoi region of a_{n-2} contains only the set $\{n - 2\}$.*

Proceeding along the similar lines as Case 1 in the proof of Proposition 4.2.2, we can show that the Voronoi region of a_{n-1} contains only the set $\{n - 1, n\}$, yielding $a_{n-2} = n - 2, a_{n-1} = Av[n - 1, n]$, and $a_n = Av[n + 1, \infty)$ with quantization error $V_n = \frac{2^{3-n}}{3}$.

Case 2. *The Voronoi region of a_{n-2} contains only the set $\{n - 2, n - 1\}$.*

Proceeding along the similar lines as Case 2 in the proof of Proposition 4.2.2, we can show that the Voronoi region of a_{n-1} contains only the set $\{n, n + 1\}$, yielding $a_{n-2} = Av[n - 2, n - 1]$, $a_{n-1} = Av[n, n + 1]$, and $a_n = Av[n + 2, \infty)$ with quantization error $V_n = \frac{2^{3-n}}{3}$.

By Case 1 and Case 2, the proof of the theorem is complete. \square

5. PROBABILITY DISTRIBUTIONS WHEN THE OPTIMAL SETS ARE GIVEN

Let P be a discrete probability measure on \mathbb{R} with support a finite or an infinite set $\{1, 2, 3, \dots\}$. Let (p_1, p_2, p_3, \dots) be a probability vector associated with $\{1, 2, 3, \dots\}$ such that the probability mass function f of P is given by $f(k) = p_k$ if $k \in \{1, 2, 3, \dots\}$, and zero otherwise. For $k, \ell \in \{1, 2, 3, \dots\}$ with $k \leq \ell$, write

$$[k, \ell] := \{n : k \leq n \leq \ell\}, \text{ and } [k, \infty) := \{k, k + 1, \dots\}.$$

For a random variable X with distribution P , let $Av[k, \ell]$ represent the conditional expectation of X given that X takes values on the set $\{k, k + 1, k + 2, \dots, \ell\}$, i.e.,

$$Av[k, \ell] = E(X : X \in [k, \ell]),$$

where $k, \ell \in \{1, 2, 3, \dots\}$ with $k \leq \ell$. On the other hand, by $Av[k, \infty)$ it is meant $Av[k, \infty) = E(X : X \in [k, \infty))$, where $k \in \{1, 2, 3, \dots\}$. Let α_n be an optimal set of n -means for P , where $n \in \mathbb{N}$. In this section, our goal is to find a set of probability vectors (p_1, p_2, p_3, \dots) such that for all $n \in \mathbb{N}$, the optimal sets of n -means are given by $\alpha_n = \{1, 2, 3, \dots, n - 1, Av[n, \infty)\}$.

Consider the following two cases:

Case 1. *$\{1, 2, 3, \dots\}$ is a finite set.*

In this case, there exists a positive integer m , such that the support of P is given by $\{1, 2, 3, \dots, m\}$. Notice that for any $k \in \{1, 2, \dots, m\}$, in this case by $[k, \infty)$ it meant the set $[k, m]$. If $m = 1$, then $\alpha_1 = \{1\}$; and if $m = 2$, then $\alpha_1 = \{Av[1, \infty)\}$, and $\alpha_2 = \{1, Av[2, \infty)\}$,

i.e., there is nothing to prove. So, we can assume that $m \geq 3$. Define the probability vector (p_1, p_2, \dots, p_m) as follows:

$$(1) \quad p_j = \begin{cases} x & \text{if } j = 1, \\ (1-x)^{j-1}x & \text{if } 2 \leq j \leq m-1, \\ (1-x)^{j-1} & \text{if } j = m. \end{cases}$$

For the sets α_n to form the optimal sets of n -means for all $1 \leq n \leq m$, we must have

$$(2) \quad (n-1) \leq \frac{1}{2}(n-1 + Av[n, \infty)) \leq n$$

for $2 \leq n \leq m$. The set of values of x obtained by solving the above inequalities does not guarantee that the sets α_n for $1 \leq n \leq m$ will form the optimal set of n -means. Thus, we need further investigation. Due to symmetry in the construction of the probability vectors, we can say that α_n for $1 \leq n \leq m$ will form the optimal sets of n -means if the following condition is also true:

$$(3) \quad V(P; \{1, Av[2, \infty)\}) \leq V(P; \{Av[1, 2], Av[3, \infty)\}).$$

Thus, we conjecture that the values of x , for which the inequalities given by (2) and (3) are true, form the set of probability vectors $(p_1, p_2, p_3, \dots, p_m)$, given by (1), for which the sets α_n for $1 \leq n \leq m$ form the optimal sets of n -means. By several examples, we verified that the conjecture is true, also see Example 5.1 and Example 5.2.

Case 2. $\{1, 2, 3, \dots\}$ is an infinite set.

Define the probability vector (p_1, p_2, p_3, \dots) as follows:

$$(4) \quad p_j = \begin{cases} x & \text{if } j = 1, \\ (1-x)^{j-1}x & \text{if } 2 \leq j. \end{cases}$$

For the sets α_n to form optimal sets of n -means for all $1 \leq n$, we must have

$$(5) \quad (n-1) \leq \frac{1}{2}(n-1 + a(n)) \leq n$$

for $2 \leq n$. The set of values of x obtained by solving the above inequalities does not guarantee that α_n for $1 \leq n$ will form an optimal set of n -means. Thus, we need further investigation. Due to symmetry in the construction of the probability vectors, we can say that the sets α_n for $1 \leq n$ will form the optimal sets of n -means if the following inequality is also true:

$$(6) \quad V(P; \{1, Av[2, \infty)\}) \leq V(P; \{Av[1, 2], Av[3, \infty)\}).$$

After some calculation, we see that there exists a real number y , the ten-digit rational approximation of which is 0.6666666667, such that the inequalities given by (5) and (6) are satisfied if $y \leq x < 1$. Thus, we conjecture that the sets α_n for $1 \leq n$ will form the optimal sets of n -means if the probability vector (p_1, p_2, p_3, \dots) is given by (4) for $0.6666666667 \leq x < 1$. By several examples, we verified that the conjecture is true.

Example 5.1. Let $m = 6$ in Case 1. Then, for $0 < x < 1$ we have

$$p_1 = x, p_2 = (1-x)x, p_3 = (1-x)^2x, p_4 = (1-x)^3x, p_5 = (1-x)^4x, \text{ and } p_6 = (1-x)^5.$$

After solving the inequalities given by (2), we have $0.4812099363 < x < 1$. Again, solving the inequality (3), we have $0.6628057756 \leq x < 1$. Notice that 0.4812099363 and 0.6628057756 are the ten-digit rational approximations of two real numbers. Thus, the inequalities given by (2) and (3) are true if $0.6628057756 \leq x < 1$. Hence, a set of probability vectors (p_1, p_2, \dots, p_6) for which the given sets α_n form the optimal sets of n means for $1 \leq n \leq 6$ is given by

$$\{(x, (1-x)x, (1-x)^2x, (1-x)^3x, (1-x)^4x, (1-x)^5) : 0.6628057756 \leq x < 1\}.$$

Example 5.2. Let $m = 7$ in Case 1. Then, proceeding as Example 5.1, we see that (2) and (3) are true if $0.6654212000 \leq x < 1$. Hence, a set of probability vectors (p_1, p_2, \dots, p_7) for which the given sets α_n form the optimal sets of n means for $1 \leq n \leq 7$ is given by

$$\{x, (1-x)x, (1-x)^2x, (1-x)^3x, (1-x)^4x, (1-x)^5x, (1-x)^6\}$$

where $0.6654212000 \leq x < 1$.

REFERENCES

- [DR] C.P. Dettmann and M.K. Roychowdhury, *Quantization for uniform distributions on equilateral triangles*, Real Analysis Exchange, Vol. 42(1), 2017, pp. 149-166.
- [GG] A. Gersho and R.M. Gray, *Vector quantization and signal compression*, Kluwer Academy publishers: Boston, 1992.
- [GL] S. Graf and H. Luschgy, *Foundations of quantization for probability distributions*, Lecture Notes in Mathematics 1730, Springer, Berlin, 2000.
- [GN] R. Gray and D. Neuhoff, *Quantization*, IEEE Trans. Inform. Theory, 44 (1998), pp. 2325-2383.
- [R] M.K. Roychowdhury, *Optimal quantizers for some absolutely continuous probability measures*, Real Analysis Exchange, Vol. 43(1), 2017, pp. 105-136.
- [RR] J. Rosenblatt and M.K. Roychowdhury, *Optimal quantization for piecewise uniform distributions*, Uniform Distribution Theory 13 (2018), no. 2, 23-55.
- [RS] M.K. Roychowdhury, and Wasila Salinas, *Quantization for a mixture of uniform distributions associated with probability vectors*, Uniform Distribution Theory 15 (2020), no. 1, 105-142.
- [Z] R. Zam, *Lattice Coding for Signals and Networks: A Structured Coding Approach to Quantization, Modulation, and Multiuser Information Theory*, Cambridge University Press, 2014.

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF TEXAS RIO GRANDE VALLEY,
1201 WEST UNIVERSITY DRIVE, EDINBURG, TX 78539-2999, USA.

E-mail address: $\{^1\text{russellloyd.cabasag01}, ^2\text{samir.huq01}, ^3\text{eric.mendoza01}, ^4\text{mrinal.roychowdhury}\}$
@utrgv.edu