University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

School of Medicine Publications and
Presentations

School of Medicine

Spring 4-29-2020

Disparity in Physician Recognition of QT interval duration and associated Mortality

Supraja Thunuguntla

The University of Texas Rio Grande Valley, supraja.thunuguntla@utrgv.edu

Mohammed Rizwan

Bhaskar Medical College

Charles Mild

The University of Texas Rio Grande Valley, chuckez@aol.com

Follow this and additional works at: https://scholarworks.utrgv.edu/som_pub

Part of the [Cardiovascular Diseases Commons](#)

Recommended Citation

Supraja Thunuguntla, Rizwan Mohammed, Charles Mild. "Disparity in Physician Recognition of QT interval duration and associated Mortality" UTRGV SOM Publications and Presentations (April, 2020).

This Article is brought to you for free and open access by the School of Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in School of Medicine Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Disparity in Physician Recognition of QT interval duration and associated Mortality

Supraja Thunuguntla [1], Mohammed Rizwan [2], Charles Mild [1]

1] UTRGV- VBMC, Internal Medicine Residency Program

2] Bhaskar Medical College, India

Background

QT interval prolongation is associated with all-cause mortality (1,2). American Heart Association/American College of Cardiology (AHA/ACC) recommends considering QTc interval above 99th percentile as abnormally prolonged, to prevent torsade de pointes (TdP).

Methods

Retrospective chart review of 291 EKG's (162 unique), from patients visiting our facility from Jan 2016 to Mar 2020. All EKG's were read, the measured QT interval was corrected for heart rate using Bazett formula. Simple descriptive statistics were created with all variables considered: patient age, gender, interpreting physician, report of long QT, presence of U-wave falsely prolonging QT and death. Correlation analysis was done, with linear mixed models to assess prolongation to account for repeated measures on some patients.

Results

The average age of patients was 65.7 years (SD=17.2, range:18-98) and 70.3% was male gender. EKG's were reviewed by a cardiologist or ER doctor 59.99%, 40.06% respectively. 4 EKG's (1.3%) were not reviewed by any physician. The 90th and 99th percentile QTc (ms) was 517 and 588.12 respectively. U-wave was included in the measurement of QT interval 1.7% times, contributed to false reporting of QTc above 99th percentile once. Using the 90th percentile cutoff, cardiologist documented prolonged QT (57.1%, 12/21) more often than an ER doctor (45.45%, 5/11) [p-value= .529]. Of the reported deaths (12/162), arrhythmia- related deaths (80%) are more likely associated with QTc above 50th percentile i.e., 475 ms than non-arrhythmia related death (14.7%) [p-value= .022]. QT prolongation was not documented in any death summary.

The Contingency tables below depict the (expected total) and the [chi-square statistic].

Long QT Reporting for QTc above 90th percentile

	Reported long QT	No report of long QT
Cardiologist	12 (11.16) [0.06]	9 (9.84) [0.07]
ER doctor	5 (5.84) [0.12]	6 (5.16) [0.14]

The chi-square statistic is 0.396. The p -value is .529148. *Not* significant at $p < .05$.

Deaths associated with QTc above 50th percentile

	Arrhythmia-related death	Non-Arrhythmia related death
QTc above 475 ms	4 (2.08) [1.76]	1 (2.92) [1.26]
QTc below 475 ms	1 (2.92) [1.26]	6 (4.08) [0.9]

The chi-square statistic is 5.182. The p -value is .022822. Significant at $p < .05$.

Conclusion

The observed trend is for cardiologists less likely to make an error than ER doctors, though not statistically significant. Based on the results of the study, this problem is likely not acted upon and maybe associated with increased mortality (more likely arrhythmia-related death). Long QT duration is a common problem and these results suggest further studies needed.

References

- 1) Charlotte Gibbs, Jacob Thalamus, Doris Tove Kristoffersen, Martin Veel Svendsen, Øystein L Holla, Kristian Heldal, Kristina H Haugaa, Jan Hysing, QT prolongation predicts short-term mortality independent of comorbidity, *EP Europace*, Volume 21, Issue 8, August 2019, Pages 1254–1260.
- 2) de Bruyne MC, Hoes AW, Kors JA, Hofman A, van Bommel JH, Grobbee DE. Prolonged QT interval predicts cardiac and all-cause mortality in the elderly. The Rotterdam Study. *Eur Heart J*. 1999 Feb;20(4):278-84.