

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Mathematical and Statistical Sciences Faculty
Publications and Presentations

College of Sciences

8-17-2016

EXISTENCE AND BEHAVIOR OF POSITIVE SOLUTIONS TO ELLIPTIC SYSTEM WITH HARDY POTENTIAL

Lei Wei

Xiyou Cheng

Zhaosheng Feng

The University of Texas Rio Grande Valley

Follow this and additional works at: https://scholarworks.utrgv.edu/mss_fac



Part of the [Mathematics Commons](#)

Recommended Citation

Wei, L., Cheng, X., & Feng, Z. (2016). EXISTENCE AND BEHAVIOR OF POSITIVE SOLUTIONS TO ELLIPTIC SYSTEM WITH HARDY POTENTIAL. *Electronic Journal of Differential Equations*, 2016, 1–14.

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

EXISTENCE AND BEHAVIOR OF POSITIVE SOLUTIONS TO ELLIPTIC SYSTEM WITH HARDY POTENTIAL

LEI WEI, XIYOU CHENG, ZHAOSHENG FENG

ABSTRACT. In this article, we study a class of elliptic systems with Hardy potentials. We analyze the possible behavior of radial solutions to the system when $p, t > 1$, $q, s > 0$ and $\lambda, \mu > (N - 2)^2/4$, and obtain the existence of positive solutions to the system with the Dirichlet boundary condition under certain conditions. When $\lambda, \mu \leq 0$, $p, t > 1$ and $q, s > 0$, we show that any radial positive solution is decreasing in r .

1. INTRODUCTION

We consider the elliptic system with singular potentials

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - u^p v^q, & x \in B_1(0) \setminus \{0\}, \\ -\Delta v &= \mu \frac{v}{|x|^2} - u^s v^t, & x \in B_1(0) \setminus \{0\}, \end{aligned} \quad (1.1)$$

where $p > 1$, $t > 1$, and $B_1(0) \subset \mathbb{R}^N$ ($N \geq 3$) is a unit ball centered at origin. The right hand side in (1.1) contains singular terms, which are usually called the singular inverse square potentials or Hardy potential in the literature. In this study, we investigate system (1.1) with the Dirichlet boundary condition

$$u(x) = v(x) = 0, \quad x \in \partial B_1(0). \quad (1.2)$$

Elliptic problems with Hardy potential have been an interesting topic in the field of singular partial differential equation for a long time. Let us briefly review some results with respect to the single equation with Hardy potential. Guerch and Véron [8] studied the equation

$$-\Delta u = V(x)u - h(u), \quad (1.3)$$

where $V(x) = \lambda|x|^{-2}$ and $h(u)$ has the property like u^p . They considered the classification of positive solutions as $\lambda \leq \frac{(N-2)^2}{4}$, and showed the behavior of positive solutions as $\lambda > \frac{(N-2)^2}{4}$ under some conditions. In a recent work [1], Cirstea studied

$$-\Delta u = \lambda \frac{u}{|x|^2} - b(x)f(u), \quad \forall x \in \Omega \setminus \{0\}, \quad (1.4)$$

where Ω is a domain containing the origin, and the function $b(x)$ is a continuous positive function (may vanish at the origin) or a singular function with a singular

2010 *Mathematics Subject Classification.* 35B40, 35J25.

Key words and phrases. Elliptic system; Hardy potential; existence; positive solution; blow up.

©2016 Texas State University.

Submitted January 12, 2016. Published August 17, 2016.

origin. It is well-known that $(N-2)^2/4$ is the Hardy constant corresponding to the Hardy potential. The function $f(u)$ has similar behavior like u^p ($p > 1$). When $\lambda \leq (\frac{N-2}{2})^2$, the qualitative properties of positive solutions at the origin were presented and a complete classification was provided.

The supercritical case has recently been studied. In [14] the authors considered the singular logistic equation

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - b(x)u^p, \quad x \in \Omega \setminus \{0\}, \\ u &= 0, \quad x \in \partial\Omega, \end{aligned} \quad (1.5)$$

where $\lambda > \frac{(N-2)^2}{4}$, $p > 1$ and $b(x)$ is a nonnegative continuous function over $\bar{\Omega}$. It shows that (1.5) has a minimal positive solution and a maximal positive solution if $b(x)$ is a positive function. Suppose that $b(x)$ has a vanishing set denoted as $\Omega_0 = \{x \in \Omega : b(x) = 0\}$. If $\Omega_0 \Subset \Omega$, the boundary of Ω belongs to C^μ , Ω_0 is a connected set and $b(0) \neq 0$, then problem (1.5) has a minimal positive solution and a maximal positive solution. In [14], the authors considered the problem

$$-\Delta u = \lambda \frac{u}{|x|^2} - |x|^\theta u^p, \quad x \in \Omega \setminus \{0\}, \quad (1.6)$$

where $\theta > -2$ and $\lambda > \frac{(N-2)^2}{4}$. The asymptotic estimate of positive solutions to (1.6) was

$$\lim_{|x| \rightarrow 0} \frac{u(x)}{|x|^{-\frac{2+\theta}{p-1}}} = \left[\lambda + \frac{2+\theta}{p-1} \left(\frac{2+\theta}{p-1} + 2 - N \right) \right]^{1/(p-1)}.$$

The uniqueness of positive solutions to (1.6) with $u = 0$ for all $x \in \partial\Omega$ was shown. In fact, a direct calculation yields

$$\lambda + \frac{2+\theta}{p-1} \left(\frac{2+\theta}{p-1} + 2 - N \right) > 0 \quad \text{when } \lambda > (N-2)^2/4.$$

This conclusion shows that any positive solution of (1.6) blows up at the origin when $\theta > -2$ and $\lambda > \frac{(N-2)^2}{4}$. When $\theta = -2$, positive solutions are uniformly bounded near the origin. When $\theta < -2$, any positive solution vanishes at the origin. So positive solutions of (1.6) have no singularity when $\theta \leq -2$. In addition, for problem (1.5), we can prove the uniqueness of positive solutions and give the exact behavior of the positive solution in a similar manner.

García-Melián and Rossi [7] considered the competitive type system

$$\begin{aligned} \Delta u &= u^p v^q, \quad x \in \Omega, \\ \Delta v &= u^s v^t, \quad x \in \Omega, \\ u &= v = \infty, \quad x \in \partial\Omega, \end{aligned} \quad (1.7)$$

where Ω is a bounded smooth domain, $p, t > 1$ and $q, s > 0$. The existence, uniqueness, blow-up rate and nonexistence of positive solutions were established under certain conditions. Li and Wang [9] considered an elliptic system in a smooth bounded domain,

$$\begin{aligned} -\Delta u &= u(a_1 - b_1 u^m - c_1 v^n) \quad x \in \Omega, \\ -\Delta v &= v(a_2 - b_2 u^p - c_2 v^q), \quad x \in \Omega, \\ u &= v = +\infty, \quad x \in \partial\Omega, \end{aligned} \quad (1.8)$$

where $a_i \geq 0$, b_i, c_i ($i = 1, 2$) are positive constants, $m, q > 0$ and $n, p \geq 0$. Based on the construction of certain sub-solution and upper-solution, some conditions on

the parameters and the exponents to ensure the existence of positive solutions were explored. García-Melián [6] extended the results in [7] to study the existence and uniqueness of positive solutions of

$$\begin{aligned}\Delta u &= a(x)u^p v^q, & x \in \Omega, \\ \Delta v &= b(x)u^r v^s, & x \in \Omega, \\ u = v &= \infty, & x \in \partial\Omega,\end{aligned}\tag{1.9}$$

where $a(x)$ and $b(x)$ satisfy

$$\begin{aligned}C_2 d(x, \partial\Omega)^{\gamma_1} &\leq a(x) \leq C_1 d(x, \partial\Omega)^{\gamma_1} \text{ for } x \in \Omega, \\ C_2 d(x, \partial\Omega)^{\gamma_2} &\leq b(x) \leq C_1 d(x, \partial\Omega)^{\gamma_2} \text{ for } x \in \Omega,\end{aligned}$$

for $\gamma_1 > -2$, $\gamma_2 > -2$.

This article is mainly devoted to the behavior analysis of positive solutions to (1.1) with the boundary condition (1.2). In our discussions, we apply some arguments and techniques with respect to boundary blow-up problems, which can be found in [2, 3, 4, 5, 10, 11, 12]. We also use the methods of subsolution and supersolution [13].

Let us summarize our main results into the following three theorems. For any radial positive solution $(u(x), v(x))$, the first theorem reveals the behavior of $u(x) + v(x)$ at the origin.

Theorem 1.1. *Suppose that $\lambda, \mu > \frac{(N-2)^2}{4}$, $p, t \geq 1$, $q, s > 0$ and (u, v) is an arbitrary radial positive solution of (1.1). Then*

$$\lim_{|x| \rightarrow 0} \{u(x) + v(x)\} = \infty.$$

When the positive solution (u, v) satisfies $u \rightarrow \infty$ and $v \rightarrow \infty$ as $|x| \rightarrow 0$, (u, v) is said to be a blow-up solution. The following theorem is regarding the existence of blow-up solution and an estimate of blow-up solution near the origin.

Theorem 1.2. *Suppose that $\lambda, \mu > \frac{(N-2)^2}{4}$, $p-1 > s > 0$ and $t-1 > q > 0$. Then the following two statements are true.*

- (i) *Problem (1.1) with condition (1.2) has at least one positive blow-up solution (U, V) .*
- (ii) *Assume that $\lambda > \max\{4\lambda_1[B_1(0)], (N-2)^2/4\}$ and (\hat{u}, \hat{v}) is a blow-up solution of the problem (1.1) with condition (1.2). Then for any $\tau > 0$ there exists $\delta := \delta(\tau) > 0$ and $C_1, C_2 > 0$, such that*

$$C_1 |x|^{-\frac{2(t-1-q)}{(p-1)(t-1)-qs} + \tau} \leq \hat{u}(x) \leq C_2 |x|^{-\frac{2(t-1-q)}{(p-1)(t-1)-qs} - \tau}, \quad \forall x \in B_\delta(0) \setminus \{0\}, \tag{1.10}$$

$$C_1 |x|^{-\frac{2(p-1-s)}{(p-1)(t-1)-qs} + \tau} \leq \hat{v}(x) \leq C_2 |x|^{-\frac{2(p-1-s)}{(p-1)(t-1)-qs} - \tau}, \quad \forall x \in B_\delta(0) \setminus \{0\}. \tag{1.11}$$

For the parameters $\lambda, \mu \leq 0$, we may obtain the behavior of positive solutions.

Theorem 1.3. *Suppose that $\lambda, \mu \leq 0$, $p, t > 1$, $q, s > 0$ and (u, v) is an arbitrary positive radial solution of (1.1) with condition (1.2). Then we have*

- (i) *$u'(r), v'(r) < 0$ for $r \in (0, 1]$. Moreover, $u(r)$ and $v(r)$ are convex functions; and*
- (ii) *$u(r) \rightarrow \infty$ and $v(r) \rightarrow \infty$ when $r \rightarrow 0$.*

The rest of this article is organized as follows. In Section 2, we introduce some preliminary results which will be used in the proofs of our main results. In Section 3, some rough estimates of positive solution of the problem (1.1) are established. Section 4 is dedicated to the existence of positive solution to problem (1.1) with boundary condition (1.2) and the estimate of blow-up rate of positive solution at the origin, i.e., the proof of Theorem 1.2. Finally, in Section 5 we show the proof of Theorem 1.3.

2. PRELIMINARY RESULTS

To make our discussions in a straightforward manner, we need the following three technical lemmas. The first lemma is regarding the comparison principle which can be found in [3, 5].

Lemma 2.1. *Suppose that Ω is a bounded domain in \mathbb{R}^N , $\alpha(x)$ and $\beta(x)$ are continuous functions in Ω with $\|\alpha\|_\infty < \infty$, and $\beta(x)$ is nonnegative and not identically zero. Let $u_1, u_2 \in C^1(\Omega)$ be positive in Ω and in the weak sense satisfy*

$$\begin{aligned} \Delta u_1 + \alpha(x)u_1 - \beta(x)g(u_1) &\leq 0 \leq \Delta u_2 + \alpha(x)u_2 - \beta(x)g(u_2), \quad x \in \Omega, \\ \limsup_{x \rightarrow \partial\Omega} (u_2 - u_1) &\leq 0, \end{aligned}$$

where $g(u)$ is continuous and such that $\frac{g(u)}{u}$ is strictly increasing with respect to u in the range of $\min\{u_1, u_2\} < u < \max\{u_1, u_2\}$. Then $u_2 \leq u_1$ holds.

For the Hardy potential, it is well-known that the Hardy constant $H = \frac{(N-2)^2}{4}$ and the Hardy inequality

$$\frac{(N-2)^2}{4} \int_{\Omega} \frac{\phi^2}{|x|^2} dx \leq \int_{\Omega} |\nabla\phi|^2 dx, \quad \forall \phi \in W_0^{1,2}(\Omega).$$

In addition, $\frac{(N-2)^2}{4}$ can be expressed as

$$\frac{(N-2)^2}{4} = \inf_{\phi \in W_0^{1,2}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla\phi|^2 dx}{\int_{\Omega} \frac{\phi^2}{|x|^2} dx}, \quad (2.1)$$

but it is not attained. Let $\lambda_1[a(x), \Omega]$ denote the first eigenvalue of

$$\begin{aligned} -\Delta u &= \lambda a(x)u, \quad x \in \Omega, \\ u &= 0, \quad x \in \partial\Omega. \end{aligned}$$

A close relation between the Hardy constant and the first eigenvalue of the Dirichlet eigenvalue problem is given in [14].

Lemma 2.2. *Suppose that Ω is a bounded smooth domain and $0 \in \Omega$. Then we have*

$$\begin{aligned} \lim_{\delta \rightarrow 0} \lambda_1 \left[\frac{1}{|x|^2}, \Omega_\delta \right] &= \frac{(N-2)^2}{4}; \\ \lim_{\epsilon \rightarrow 0} \lambda_1 \left[\frac{1}{|x|^2 + \epsilon}, \Omega \right] &= \frac{(N-2)^2}{4}, \end{aligned}$$

where $\Omega_\delta = \{x \in \Omega : |x| > \delta\}$.

By the uniqueness and blow-up rate of positive solutions for a single equation with the Hardy potential [14], we can obtain the following conclusion.

Lemma 2.3. *Suppose that $p > 1$, $\sigma > -2$, $\lambda > (N-2)^2/4$, Ω is a bounded smooth domain and $0 \in \Omega$.*

(i) *Assume that u is any positive solution of*

$$-\Delta u = \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, \quad x \in \Omega \setminus \{0\}.$$

Then u satisfies

$$\lim_{|x| \rightarrow 0} |x|^{\frac{2+\sigma}{p-1}} u(x) = \left[\lambda + \frac{2+\sigma}{p-1} \left(\frac{2+\sigma}{p-1} + 2 - N \right) \right]^{1/(p-1)}.$$

(ii) *Assume that c is a nonnegative constant ($c = +\infty$ is also allowed). Then the problem*

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, & x \in \Omega, \\ u &= c, & x \in \partial\Omega \end{aligned} \tag{2.2}$$

has a unique positive solution $U_{p,\sigma}$.

Proof. Firstly, Part (i) is the direct result of [14]. Now, we simply sketch the proof of Part (ii). Suppose that $c = 0$. Since $\lambda > \frac{(N-2)^2}{4}$, for any sufficiently small $\delta > 0$, by Lemma 2.2 and the standard arguments of logistic equations, it follows that

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, & x \in \Omega_\delta, \\ u &= 0, & x \in \partial\Omega_\delta \end{aligned}$$

has a unique positive solution u_δ . By the comparison principle and regularity arguments of elliptic equations, $w = \lim_{\delta \rightarrow 0} u_\delta$ is a positive solution of the problem (2.2). Suppose that $c > 0$ ($= \infty$), then a similar argument shows that (2.2) has a positive solution w . Suppose that ω is any positive solution of (2.2). By Part (i), we have

$$\lim_{|x| \rightarrow 0} |x|^{\frac{2+\sigma}{p-1}} \omega(x) = \left[\lambda + \frac{2+\sigma}{p-1} \left(\frac{2+\sigma}{p-1} + 2 - N \right) \right]^{1/(p-1)}.$$

If $c = \infty$, by the standard arguments of boundary blow-up problems, we can find the exact behavior near boundary for positive solutions. By the standard method, one can see the uniqueness of positive solutions. \square

3. BLOW-UP BEHAVIOR OF POSITIVE SOLUTIONS

In this section, we present some behavior analysis of positive solutions of (1.1) and prove Theorem 1.1.

Lemma 3.1. *Suppose that $p, t \geq 1$, $q, s > 0$, $\lambda, \mu > \frac{(N-2)^2}{4}$ and $(u(x), v(x))$ is an arbitrary solution of (1.1). Then*

$$\limsup_{|x| \rightarrow 0} \{u(x) + v(x)\} = \infty.$$

That is,

$$\limsup_{|x| \rightarrow 0} u(x) = \infty \quad \text{or} \quad \limsup_{|x| \rightarrow 0} v(x) = \infty.$$

Proof. By the way of contradiction, we suppose that $\limsup_{|x| \rightarrow 0^+} \{u(x) + v(x)\} = M$, where $M \in [0, \infty)$. Then there is a $\delta > 0$ such that $0 < v(x) \leq M + 1$ for all $x \in B_\delta(0) \setminus \{0\}$.

Case 1: $p > 1$. From the first equation of (1.1), we have

$$-\Delta u \geq \lambda \frac{u}{|x|^2} - (M+1)^q u^p \text{ in } B_\delta(0) \setminus \{0\}. \quad (3.1)$$

Let $w(x) = (M+1)^{\frac{q}{p-1}} u$, then we obtain

$$-\Delta w \geq \lambda \frac{w}{|x|^2} - w^p \text{ in } B_\delta(0) \setminus \{0\}.$$

Since $\lambda > \frac{(N-2)^2}{4}$, by Lemma 2.2, for any sufficiently small $\epsilon > 0$, the problem

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - u^p, & x \in B_\delta \setminus \overline{B}_\epsilon(0), \\ u &= 0, & |x| = \delta \text{ or } |x| = \epsilon \end{aligned}$$

has a unique positive solution u_ϵ . By the comparison principle, the function $u_*(x) := \lim_{\epsilon \rightarrow 0} u_\epsilon(x)$ is well defined in $B_\delta(0) \setminus \{0\}$, and hence by the regularity arguments, u_* satisfies

$$-\Delta u_* = \lambda \frac{u_*}{|x|^2} - u_*^p \text{ in } B_\delta(0) \setminus \{0\}.$$

By the comparison principle,

$$w(x) \geq u_\epsilon(x) \quad \text{for all } x \in B_\delta(0) \setminus \overline{B}_\epsilon(0).$$

Letting $\epsilon \rightarrow 0$, we find

$$w(x) \geq u_*(x) \quad \text{for all } x \in B_\delta(0) \setminus \{0\}.$$

By Lemma 2.3, $\lim_{|x| \rightarrow 0^+} u(x) = \infty$, which contradicts $\limsup_{|x| \rightarrow 0^+} \{u(x) + v(x)\} = M$.

Case 2: $p = 1$. Since $\lambda > \frac{(N-2)^2}{4}$ and using (3.1), there is $\tau \in (0, \delta)$ such that

$$-\Delta u \geq \frac{1}{2} \left(\lambda + \frac{(N-2)^2}{4} \right) \frac{u}{|x|^2} \quad \text{for } x \in B_\tau(0) \setminus \{0\}.$$

For each $\eta \in (0, \tau)$, u is a positive supersolution of

$$\begin{aligned} -\Delta \phi &= \frac{1}{2} \left(\lambda + \frac{(N-2)^2}{4} \right) \frac{\phi}{|x|^2}, & x \in B_\tau(0) \setminus \overline{B}_\eta(0), \\ \phi(x) &= 0, & x \in \partial B_\tau(0) \cup \partial B_\eta(0). \end{aligned}$$

So, we derive that

$$\lambda_1 \left[\frac{1}{|x|^2}, B_\tau(0) \setminus \overline{B}_\eta(0) \right] > \frac{1}{2} \left(\lambda + \frac{(N-2)^2}{4} \right),$$

which contradicts Lemma 2.2, namely

$$\lim_{\eta \rightarrow 0} \lambda_1 \left[\frac{1}{|x|^2}, B_\tau(0) \setminus \overline{B}_\eta(0) \right] = \frac{(N-2)^2}{4}.$$

□

Remark 3.2. Suppose that $p, t \geq 1$, $q, s > 0$ and $\lambda, \mu > (N-2)^2/4$. From the proof of Lemma 3.1, it follows that $\lim_{|x| \rightarrow 0} v(x) = \infty$ if u is bounded near the origin, and $\lim_{|x| \rightarrow 0} u(x) = \infty$ if v is bounded near the origin.

Proof of Theorem 1.1. Suppose that $(u(x), v(x))$ is an arbitrary positive radial solution of (1.1). By Lemma 3.1, we have

$$\limsup_{|x| \rightarrow 0} \{u(x) + v(x)\} = \infty.$$

It suffices to show that

$$\liminf_{|x| \rightarrow 0} \{u(x) + v(x)\} = \infty.$$

By the way of contradiction, we suppose $\liminf_{|x| \rightarrow 0} \{u(x) + v(x)\} < \infty$. For convenience of our statement, we denote $u(r) = u(x)$ and $v(r) = v(x)$ when $|x| = r$. In view of

$$\liminf_{|x| \rightarrow 0} \{u(x) + v(x)\} < \limsup_{|x| \rightarrow 0} \{u(x) + v(x)\},$$

there is $\{r_n\}$ with $r_n \rightarrow 0$ such that

$$\begin{aligned} u''(r_n) + v''(r_n) &\geq 0, & u'(r_n) + v'(r_n) &= 0, \\ \lim_{n \rightarrow \infty} \{u(r_n) + v(r_n)\} &= \liminf_{|x| \rightarrow 0} \{u(x) + v(x)\}. \end{aligned}$$

Without loss of generality, we assume that $\lambda \leq \mu$. It follows from (1.1) that

$$\frac{\lambda}{r_n^2} \leq \frac{u(r_n)^p v(r_n)^q + u(r_n)^s v(r_n)^t}{u(r_n) + v(r_n)}, \quad (3.2)$$

which implies

$$\frac{\lambda}{r_n^2} < u(r_n)^{p-1} v(r_n)^q + u(r_n)^s v(r_n)^{t-1}.$$

So, we have

$$u(r_n)^{p-1} v(r_n)^q + u(r_n)^s v(r_n)^{t-1} \rightarrow \infty \quad (n \rightarrow \infty).$$

This indicates that $\{u(r_n) + v(r_n)\}$ is an unbounded set. This contradicts the assumption of $\lim_{n \rightarrow \infty} (u(r_n) + v(r_n)) < \infty$. \square

We now show some analysis on the behavior of positive solutions to (1.1) near the origin.

Proposition 3.3. *Suppose that $\lambda, \mu > (N - 2)^2/4$, $t - 1 > q > 0$, $p - 1 > s > 0$, and $(u(x), v(x))$ is an arbitrary positive solution of the system (1.1) such that both $\lim_{|x| \rightarrow 0} u(x)$ and $\lim_{|x| \rightarrow 0} v(x)$ exist. Then at least one of following statements holds: (i) both $u(x)$ and $v(x)$ blow up at the origin; (ii) $u(x)$ blows up and $v(x)$ vanishes at the origin; and (iii) $u(x)$ vanishes and $v(x)$ blows up at the origin.*

Proof. We claim that either $\lim_{|x| \rightarrow 0} u(x) = \infty$ or $\lim_{|x| \rightarrow 0} u(x) = 0$ holds. Otherwise, we suppose $\lim_{|x| \rightarrow 0} u(x) = m \in (0, \infty)$. So there exists a $\delta > 0$ such that

$$m/2 \leq u(x) \leq 2m \quad \text{for } x \in B_\delta(0) \setminus \{0\}.$$

Then, we see that

$$-\Delta v \leq \mu \frac{v}{|x|^2} - (m/2)^s v^t \quad \text{for } x \in B_\delta(0) \setminus \{0\}.$$

Let $w_1 = (m/2)^{\frac{s}{t-1}} v$. Then

$$-\Delta w_1 \leq \mu \frac{w_1}{|x|^2} - w_1^t \quad \text{in } B_\delta(0) \setminus \{0\}.$$

By Lemma 2.3 and an analogous argument described in [14], it follows that

$$v(x) \leq C|x|^{-\frac{2}{t-1}} \text{ for } x \in B_\delta(0) \setminus \{0\}.$$

From the first equation of (1.1), we have

$$-\Delta u \geq \lambda \frac{u}{|x|^2} - C^q|x|^{-\frac{2q}{t-1}}u^p \text{ in } B_\delta(0) \setminus \{0\}.$$

Let $w_2 = C^{\frac{q}{p-1}}u$. Then

$$-\Delta w_2 \geq \lambda \frac{w_2}{|x|^2} - |x|^{-\frac{2q}{t-1}}w_2^p \text{ in } B_\delta(0) \setminus \{0\}.$$

Hence, by $t - 1 > q$ and Lemma 2.3, we deduce that

$$u(x) \geq C_1|x|^{-\frac{2-\frac{2q}{t-1}}{p-1}} = C_1|x|^{-\frac{2(t-1-q)}{(p-1)(t-1)}} \text{ on } B_\delta(0) \setminus \{0\},$$

which contradicts the assumption $\lim_{|x| \rightarrow 0} u(x) = m \in (0, \infty)$.

Similarly, we can prove that either $\lim_{|x| \rightarrow 0} v(x) = \infty$ or $\lim_{|x| \rightarrow 0} v(x) = 0$ holds. □

Proposition 3.4. *Suppose that $p, t \geq 1$, $q, s > 0$ and $\lambda, \mu > (N - 2)^2/4$ and $(u(x), v(x))$ is an arbitrary positive solution of (1.1). Then both $u^{p-1}v^q$ and $u^s v^{t-1}$ are unbounded near the origin. In particular, when $p = 1$ and $t = 1$, both u and v are unbounded near the origin.*

Proof. Suppose that $u^{p-1}v^q$ is bounded near the origin. Using the first equation in (1.1), we have

$$-\Delta u + (u^{p-1}v^q)u = \lambda \frac{u}{|x|^2} \text{ in } B_1(0) \setminus \{0\}.$$

Since $u^{p-1}v^q$ is bounded near the origin, there is a sufficiently small $\delta_0 > 0$ such that for any $\delta \in (0, \delta_0]$,

$$-\Delta \geq \frac{1}{2} \left[\lambda + \frac{(N - 2)^2}{4} \right] \frac{u}{|x|^2} \text{ in } B_{\delta_0}(0).$$

This implies

$$\lambda_1 \left[\frac{1}{|x|^2}, B_{\delta_0}(0) \setminus B_\delta(0) \right] \geq \frac{1}{2} \left[\lambda + \frac{(N - 2)^2}{4} \right] \text{ for } \delta \in (0, \delta_0),$$

which is a contradiction to

$$\lim_{\delta \rightarrow 0} \lambda_1 \left[\frac{1}{|x|^2}, B_{\delta_0}(0) \setminus B_\delta(0) \right] = \frac{(N - 2)^2}{4}.$$

□

4. EXISTENCE AND ESTIMATES OF BLOW-UP SOLUTIONS

In this section, we deal with the existence of positive blow-up solutions of the problem (1.1) with the boundary condition (1.2) and establish estimates of positive blow-up solutions near the origin.

Suppose that $\lambda > 4\lambda_1[B_1(0)]$ and $\alpha \in (0, 1)$ satisfies $\lambda \frac{\alpha^2}{(1+\alpha)^2} > \lambda_1[B_1(0)]$. For the case $\sigma \geq 0$, let $u_{0,\alpha}$ denote the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{\alpha^2}{(1+\alpha)^2} u - \alpha^2(1+\alpha)^\sigma u^p, \quad x \in B_1(0) \\ u(x) &= 0, \quad x \in \partial B_1(0). \end{aligned}$$

For convenience, for the case of $-2 < \sigma < 0$, by $u_{0,\alpha}$ we denote the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{\alpha^2}{(1+\alpha)^2} u - \alpha^2 (1-\alpha)^\sigma u^p, & x \in B_1(0) \\ u(x) &= 0, & x \in \partial B_1(0). \end{aligned}$$

When $\sigma \geq 0$, let $u_{\infty,\alpha}$ denote the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{\alpha^2}{(1-\alpha)^2} u - \alpha^2 (1-\alpha)^\sigma u^p, & x \in B_1(0) \\ u(x) &= \infty, & x \in \partial B_1(0). \end{aligned}$$

For the case of $-2 < \sigma < 0$, let $u_{\infty,\alpha}$ denote the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{\alpha^2}{(1-\alpha)^2} u - \alpha^2 (1+\alpha)^\sigma u^p, & x \in B_1(0) \\ u(x) &= \infty, & x \in \partial B_1(0). \end{aligned}$$

In fact, there is a relation between $u_{0,\alpha}$, $u_{\infty,\alpha}$ and α . For our convenience, we denote $u_{0,\alpha}$ and $u_{\infty,\alpha}$ by u_0 and u_∞ , respectively. In some places, we will write $u_\infty(x; \lambda)$ and $u_0(x; \lambda)$ instead of u_∞ and u_0 , to clearly indicate the relation depending on the parameter λ .

Lemma 4.1. *Let $p > 1$, $\sigma > -2$, $\lambda > 4\lambda_1[B_1(0)]$ and U be a positive solution of*

$$-\Delta U = \lambda \frac{U}{|x|^2} - |x|^\sigma U^p \quad \text{on } B_R(0) \setminus \{0\}.$$

Then we have

$$u_0(0)|x|^{-\frac{2+\sigma}{p-1}} \leq U(x) \leq u_\infty(0)|x|^{-\frac{2+\sigma}{p-1}} \quad \text{for } x \in B_{R/2}(0) \setminus \{0\}. \quad (4.1)$$

Proof. Suppose that $x_0 \in B_{R/2}(0) \setminus \{0\}$ is an arbitrary point. Since $\lambda > 4\lambda_1[B_1(0)]$, we can choose $\alpha \in (0, 1)$ close to 1 such that $\lambda \frac{\alpha^2}{(1+\alpha)^2} > \lambda_1[B_1(0)]$. Let

$$\tilde{U}(x) = |x_0|^{\frac{2+\sigma}{p-1}} U(x_0 + \alpha|x_0|x), \quad x \in B_1(0).$$

So, for all $x \in B_1(0)$, we obtain

$$(1-\alpha)|x_0| \leq |(x_0 + \alpha|x_0|x)| \leq (1+\alpha)|x_0|.$$

When $\sigma \geq 0$, a straightforward calculation leads to

$$\begin{aligned} -\Delta \tilde{U} &\leq \lambda \frac{\alpha^2}{(1-\alpha)^2} \tilde{U} - \alpha^2 (1-\alpha)^\sigma \tilde{U}^p & \text{in } x \in B_1(0), \\ -\Delta \tilde{U} &\geq \lambda \frac{\alpha^2}{(1+\alpha)^2} \tilde{U} - \alpha^2 (1+\alpha)^\sigma \tilde{U}^p & \text{in } x \in B_1(0). \end{aligned}$$

We find that if $0 > \sigma > -2$, then

$$\begin{aligned} -\Delta \tilde{U} &\leq \lambda \frac{\alpha^2}{(1-\alpha)^2} \tilde{U} - \alpha^2 (1+\alpha)^\sigma \tilde{U}^p & x \in B_1(0), \\ -\Delta \tilde{U} &\geq \lambda \frac{\alpha^2}{(1+\alpha)^2} \tilde{U} - \alpha^2 (1-\alpha)^\sigma \tilde{U}^p & x \in B_1(0). \end{aligned}$$

By the comparison principle, we obtain

$$u_0(x) \leq \tilde{U}(x) \leq u_\infty(x) \quad \text{for all } x \in B_1(0).$$

Particularly, choosing $x = 0$ and by the arbitrariness of x_0 , we arrive at (4.1). \square

Lemma 4.2. *Suppose that $C > 0$, $R > 0$ and $\lambda > \max\{4\lambda_1[B_1(0)], (N - 2)^2/4\}$.*

(i) *If $\mathcal{U} > 0$ satisfies*

$$-\Delta \mathcal{U} \geq \lambda \frac{\mathcal{U}}{|x|^2} - C|x|^\sigma \mathcal{U}^p, \quad x \in B_R(0) \setminus \{0\},$$

then

$$\mathcal{U}(x) \geq C^{-\frac{1}{p-1}} u_0(0) |x|^{-\frac{2+\sigma}{p-1}} \quad \text{for } x \in B_{R/2}(0) \setminus \{0\}. \quad (4.2)$$

(ii) *If $\mathcal{U} > 0$ satisfies*

$$-\Delta \mathcal{U} \leq \lambda \frac{\mathcal{U}}{|x|^2} - C|x|^\sigma \mathcal{U}^p, \quad x \in B_R(0) \setminus \{0\}, \quad (4.3)$$

then

$$\mathcal{U}(x) \leq C^{-\frac{1}{p-1}} u_\infty(0) |x|^{-\frac{2+\sigma}{p-1}} \quad \text{for } x \in B_{R/2}(0) \setminus \{0\}. \quad (4.4)$$

Proof. From the condition in case (i), it follows that $C^{\frac{1}{p-1}} \mathcal{U}$ is a supersolution of

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, \quad x \in B_R(0) \setminus \{0\}, \\ u &= 0, \quad x \in \partial B_R(0). \end{aligned} \quad (4.5)$$

Let U be the unique positive solution of (4.5), then

$$U(x) = \lim_{\delta \rightarrow 0} U_\delta(x),$$

where U_δ is the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, \quad x \in B_R(0) \setminus \overline{B}_\delta(0), \\ u &= 0, \quad x \in \partial B_R(0) \cup \partial B_\delta(0). \end{aligned} \quad (4.6)$$

So, the comparison principle yields

$$U_\delta(x) \leq C^{\frac{1}{p-1}} \mathcal{U}(x) \quad \text{for all } x \in B_R(0) \setminus B_\delta(0).$$

Further, letting $\delta \rightarrow 0$, we have

$$U(x) \leq C^{\frac{1}{p-1}} \mathcal{U}(x) \quad \text{for all } x \in B_R(0).$$

In view of Lemma 4.1, we arrive at (4.2).

Now, we prove conclusion (ii). Suppose that $\mathcal{U} > 0$ satisfies (4.3). Let V_δ be the unique positive solution of

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - |x|^\sigma u^p, \quad x \in B_R(0) \setminus \overline{B}_\delta(0), \\ u &= \infty, \quad x \in \partial B_R(0) \cup \partial B_\delta(0). \end{aligned} \quad (4.7)$$

From [14], we know that

$$\hat{U}(x) := \lim_{\delta \rightarrow 0} V_\delta(x) \quad \text{for all } x \in B_R(0) \setminus \{0\} \text{ is well defined,}$$

and that \hat{U} satisfies

$$\begin{aligned} -\Delta \hat{U} &= \lambda \frac{\hat{U}}{|x|^2} - |x|^\sigma \hat{U}^p, \quad x \in B_R(0) \setminus \{0\}, \\ \hat{U} &= \infty, \quad x \in \partial B_R(0). \end{aligned}$$

So by the comparison principle, it is easy to see

$$C^{\frac{1}{p-1}}\mathcal{U}(x) \leq \hat{U}(x) \quad \text{for } x \in B_R(0) \setminus \{0\}.$$

Using Lemma 4.1 again, we arrive at (4.4). □

Next, we give the proof of Theorem 1.2, i.e., we explore the existence of blow-up solution of the problem (1.1) with the boundary condition (1.2), and give the estimate of positive solutions near the origin.

Proof of Theorem 1.2. Since $s < p - 1$ and $q < t - 1$, we may choose $\gamma, \sigma \in (-2, 0)$ such that

$$-\frac{\gamma}{q} = \frac{2 + \sigma}{t - 1} \quad \text{and} \quad -\frac{\sigma}{s} = \frac{2 + \gamma}{p - 1}. \tag{4.8}$$

Denote by $U_{p,\gamma}$ the unique positive solution of (4.5) with σ being replaced by γ and $R = 1$, and denote by $U_{t,\sigma}$ the unique positive solution of (4.5) with p being replaced by t and $R = 1$. Choose $\eta \in (s/(t - 1), (p - 1)/q)$. A simple calculation shows that for sufficiently small $\epsilon > 0$ and sufficiently large $M > 0$, we have

$$\begin{aligned} -\Delta(\epsilon U_{p,\gamma}) &\leq \lambda \frac{\epsilon U_{p,\gamma}}{|x|^2} - (\epsilon U_{p,\gamma})^p (\epsilon^{-\eta} U_{t,\sigma})^q \quad \text{in } B_1(0) \setminus \{0\}, \\ -\Delta(\epsilon^{-\eta} U_{t,\sigma}) &\geq \lambda \frac{\epsilon^{-\eta} U_{t,\sigma}}{|x|^2} - (\epsilon U_{p,\gamma})^s (\epsilon^{-\eta} U_{t,\sigma})^t \quad \text{in } B_1(0) \setminus \{0\}, \\ -\Delta(M U_{p,\gamma}) &\geq \lambda \frac{M U_{p,\gamma}}{|x|^2} - (M U_{p,\gamma})^p (M^{-\eta} U_{t,\sigma})^q \quad \text{in } B_1(0) \setminus \{0\}, \\ -\Delta(M^{-\eta} U_{t,\sigma}) &\leq \lambda \frac{M^{-\eta} U_{t,\sigma}}{|x|^2} - (M U_{p,\gamma})^s (M^{-\eta} U_{t,\sigma})^t \quad \text{in } B_1(0) \setminus \{0\}. \end{aligned}$$

So, $(\epsilon U_{p,\gamma}, M^{-\eta} U_{t,\sigma})$ and $(M U_{p,\gamma}, \epsilon^{-\eta} U_{t,\sigma})$ are a pair of subsolution and supersolution of

$$\begin{aligned} -\Delta u &= \lambda \frac{u}{|x|^2} - u^p v^q, \quad x \in B_1(0) \setminus \overline{B}_{1/n}(0), \\ -\Delta v &= \mu \frac{v}{|x|^2} - u^s v^t, \quad x \in B_1(0) \setminus \overline{B}_{1/n}(0), \\ u &= v = 0, \quad x \in \partial B_1(0), \\ u &= \epsilon U_{p,\gamma}, \quad v = M^{-\eta} U_{t,\sigma}, \quad x \in \partial B_{1/n}(0). \end{aligned} \tag{4.9}$$

By the supersolution and subsolution method, we know that (4.9) has at least one solution $(U_n(x), V_n(x))$ satisfying

$$\begin{aligned} \epsilon U_{p,\gamma} &\leq U_n(x) \leq M U_{p,\gamma}(x) \quad \text{for } x \in B_1(0) \setminus \overline{B}_{1/n}(0), \\ M^{-\eta} U_{t,\sigma}(x) &\leq V_n(x) \leq \epsilon^{-\eta} U_{t,\sigma}(x) \quad \text{for } x \in B_1(0) \setminus \overline{B}_{1/n}(0). \end{aligned}$$

In view of the regularity arguments of elliptic equations, (U_n, V_n) has a convergent subsequence in $C_{loc}^2(\overline{B}_1(0) \setminus \{0\}) \times C_{loc}^2(\overline{B}_1(0) \setminus \{0\})$. We by (u, v) denote the limit functions, and hence (u, v) is a solution of the problem (1.1). Clearly, since both $U_{p,\gamma}$ and $U_{t,\sigma}$ blow up at the origin, we can see that both u and v blow up at the origin. The proof of (i) is complete.

Now, we prove the conclusion (ii). By hypothesis, both $\hat{u}(x)$ and $\hat{v}(x)$ have positive bound from below in $B_{1/2}(0)$. Denote $m_0 = \inf\{\hat{v}(x) : x \in B_{1/2}(0)\}$, and hence \hat{u} satisfies

$$-\Delta \hat{u} \leq \lambda \frac{\hat{u}}{|x|^2} - m_0^q \hat{u}^p, \quad x \in B_{1/2}(0) \setminus \{0\}.$$

In view of Lemma 4.2, we have

$$\hat{u}(x) \leq m_0^{-\frac{q}{p-1}} u_\infty(0; \lambda) |x|^{-\frac{2}{p-1}} \quad \text{for } x \in B_{1/4}(0) \setminus \{0\}.$$

Let $a_1 = m_0^{-\frac{q}{p-1}} u_\infty(0; \lambda)$ and $\alpha_0 = \frac{2}{p-1}$. Then we find that

$$\hat{u}(x) \leq a_1 |x|^{-\alpha_0} \quad \text{for } x \in B_{1/4}(0) \setminus \{0\}.$$

By the second equation in (1.1), it follows that

$$-\Delta \hat{v} \geq \mu \frac{\hat{v}}{|x|^2} - a_1^s |x|^{-\frac{2s}{p-1}} \hat{v}^t, \quad x \in B_{1/4}(0) \setminus \{0\}.$$

Since $s < p - 1$, we have $-\frac{2s}{p-1} > -2$. Take $a_2 = a_1^{-\frac{s}{t-1}}$. In view of Lemma 4.2 again, we obtain

$$\hat{v}(x) \geq a_2 |x|^{-\frac{2-s\alpha_0}{t-1}} \quad \text{for } x \in B_{2^{-3}}(0) \setminus \{0\}.$$

Choose $\beta_1 = \frac{2-s\alpha_0}{t-1}$, and hence we see that

$$\hat{v}(x) \geq a_2 |x|^{-\beta_1} \quad \text{for } x \in B_{2^{-3}}(0) \setminus \{0\}.$$

Let $a_3 = a_2^{-\frac{q}{p-1}}$ and $\alpha_2 = \frac{2-q\beta_1}{p-1}$. Using the first equation in (1.1) and Lemma 4.2 again, we have

$$\hat{u} \leq a_3 |x|^{-\frac{2-q\beta_1}{p-1}} \quad \text{in } B_{2^{-4}}(0) \setminus \{0\}.$$

Proceeding in this way inductively, we obtain

$$\hat{u}(x) \leq a_{2n-1} |x|^{-\frac{2-q\beta_{n-1}}{p-1}}, \quad x \in B_{2^{-2n}}(0) \setminus \{0\},$$

$$\hat{v}(x) \geq a_{2n} |x|^{-\frac{2-q\alpha_{n-1}}{t-1}}, \quad x \in B_{2^{-(2n+1)}}(0) \setminus \{0\},$$

where $n \geq 1$, and

$$\begin{aligned} \alpha_0 &= \frac{2}{p-1}, \quad \beta_0 = 0, \quad \alpha_n = \frac{2-q\beta_n}{p-1}, \quad \beta_n = \frac{2-s\alpha_{n-1}}{t-1}, \\ a_1 &= m_0^{-\frac{q}{p-1}} u_\infty(0; \lambda), \quad a_{2n} = a_{2n-1}^{-\frac{s}{t-1}}, \quad a_{2n+1} = a_{2n}^{-\frac{q}{p-1}}. \end{aligned}$$

From the relation between α_n and β_n , we can deduce that $\{\alpha_n\}$ is a decreasing sequence and $\{\beta_n\}$ is an increasing sequence. By letting $n \rightarrow \infty$, it follows that

$$\lim_{n \rightarrow \infty} \alpha_n = \frac{2(t-1-q)}{(p-1)(t-1)-qs} \quad \text{and} \quad \lim_{n \rightarrow \infty} \beta_n = \frac{2(p-1-s)}{(p-1)(t-1)-qs}.$$

From the relation between a_n and a_{n+1} , we can deduce that

$$a_{2n+1} = a_1^{\frac{q^n s^n}{(t-1)^n (p-1)^n}}.$$

In view of $(p-1)(t-1) > qs$ and $\min\{a_1, 1\} \leq a_{2n+1} \leq \max\{a_1, 1\}$ for all n , we have

$$\min\{a_1, a_1^{-\frac{s}{t-1}}\} \leq a_n \leq \max\{a_1, a_1^{-\frac{s}{t-1}}\} \quad \text{for all } n.$$

Thus, for any $\tau > 0$ there are $\delta := \delta(\tau)$ and $C_1, C_2 > 0$ such that

$$C_1 |x|^{-\frac{2(p-1-s)}{(p-1)(t-1)-qs} + \tau} \leq \hat{v}(x) \quad \text{for } x \in B_\delta(0) \setminus \{0\},$$

$$\hat{u}(x) \leq C_2 |x|^{-\frac{2(t-1-q)}{(p-1)(t-1)-qs} - \tau} \quad \text{for } x \in B_\delta(0) \setminus \{0\}.$$

Consequently, the second inequality of (1.10) and the first inequality of (1.11) hold. By a similar argument, we can prove the remaining cases of (1.10) and (1.11). \square

Remark 4.3. From the proof of (i) in Theorem 1.2, it follows that (1.1) has a positive solution (u, v) satisfying

$$\begin{aligned} C_1|x|^{-\frac{2+\gamma}{p-1}} &\leq u(x) \leq C_2|x|^{-\frac{2+\gamma}{p-1}} \quad \text{for } x \in \Omega \setminus \{0\}, \\ C_1|x|^{-\frac{2+\sigma}{i-1}} &\leq v(x) \leq C_2|x|^{-\frac{2+\sigma}{i-1}} \quad \text{for } x \in \Omega \setminus \{0\}, \end{aligned}$$

where γ and σ satisfies (4.8). In fact, a simple calculation shows that

$$\frac{2+\gamma}{p-1} = \frac{2(t-1-q)}{(p-1)(t-1)-qs} \quad \text{and} \quad \frac{2+\sigma}{t-1} = \frac{2(p-1-s)}{(p-1)(t-1)-qs}.$$

5. PROOF OF THEOREM 1.3

Proof. Suppose that $(u(x), v(x))$ is a positive radial solution of (1.1) with condition (1.2). For convenience, we denote $u(r) = u(x)$ and $v(r) = v(x)$ as $|x| = r$. Define

$$t_0 = \inf\{r_0 \in (0, 1) : u'(r) < 0 \text{ and } v'(r) < 0 \text{ for } r \in (r_0, 1)\}.$$

By Hopf's lemma, when $r < 1$ is close to 1, we have $u'(r) < 0$ and $v'(r) < 0$. So, this leads to $t_0 < 1$. We claim $t_0 = 0$. Otherwise, we will see that

Case (1) $u'(t_0) = 0$, $u'(r) < 0$ and $v'(r) < 0$ for $r \in (t_0, 1)$, and

Case (2) $v'(t_0) = 0$, $v'(r) < 0$ and $u'(r) < 0$ for $r \in (t_0, 1)$.

Without loss of generality, we assume that Case (1) occurs. Denote

$$\tilde{u} = u(t_0) - u(x) \quad \text{for } t_0 < |x| < 1.$$

Then, $\tilde{u}(x) > 0$ when $t_0 < |x| < 1$ and $\tilde{u}'(t_0) = \tilde{u}(t_0) = 0$. A simple calculation gives

$$-\Delta\tilde{u} - \lambda\frac{\tilde{u}}{|x|^2} = -\lambda\frac{u(t_0)}{|x|^2} + u^p v^q > 0 \quad \text{when } t_0 < |x| < 1.$$

By Hopf's lemma, there holds $\tilde{u}'(t_0) > 0$, which contradicts $\tilde{u}'(t_0) = 0$. The conclusion $t_0 = 0$ implies that

$$u'(r) < 0 \text{ and } v'(r) < 0 \text{ for } r \in (0, 1).$$

Since $u'(r) < 0$ in $(0, 1)$ and $\lambda \leq 0$, we have $u'' \geq u^p v^q$ in $(0, 1)$, which implies that $u(r)$ is a convex function in $(0, 1)$. Analogously, we may show that v is a convex function. So, the conclusion (i) is proved.

Now we prove the conclusion (ii). From (1.1), we see that

$$-(r^{N-1}u')' = \lambda r^{N-3}u - r^{N-1}u^p v^q \text{ for } r \in (0, r).$$

Integrating on $[r, 1]$ gives

$$r^{N-1}u'(r) - u'(1) = \lambda \int_r^1 \tau^{N-3}u(\tau)d\tau - \int_r^1 \tau^{N-1}u(\tau)^p v(\tau)^q d\tau \quad \text{for } r \in (0, 1).$$

That is,

$$\begin{aligned} u'(r) &= r^{1-N}u'(1) + \lambda r^{1-N} \int_r^1 \tau^{N-3}u(\tau)d\tau \\ &\quad - r^{1-N} \int_r^1 \tau^{N-1}u(\tau)^p v(\tau)^q d\tau \quad \text{for } r \in (0, 1). \end{aligned}$$

Integrating on $[r, 1]$ yields

$$u(r) = - \int_r^1 s^{1-N}u'(1)ds - \lambda \int_r^1 s^{1-N} \int_s^1 \tau^{N-3}u(\tau)d\tau ds$$

$$\begin{aligned}
& + \int_r^1 s^{1-N} \int_s^1 \tau^{N-1} u(\tau)^p v(\tau)^q d\tau ds \\
& \geq - \int_r^1 s^{1-N} u'(1) ds.
\end{aligned}$$

This inequality implies that $u(r) \rightarrow \infty$ as $r \rightarrow 0$. In a similar manner, one can see that $v(r) \rightarrow \infty$ as $r \rightarrow 0$. \square

Acknowledgements. This work is supported by NSF of China, No. 11471148.

REFERENCES

- [1] F. Cîrstea; A complete classification of the isolated singularities for nonlinear elliptic equations with inverse square potentials, *Memoirs of AMS*, Vol. 227 Number 1068, 2014.
- [2] F. Cîrstea, V. D. Rădulescu; Existence and uniqueness of blow-up solutions for a class of logistic equations, *Commun. Contemp. Math.*, **4** (2002), 559-586.
- [3] Y. Du; Order structure and topological methods in nonlinear partial differential equations, *Vol I Maximum Principle and Applications*, World Scientific Publishing, 2006.
- [4] Y. Du, Q. Huang; Blow-up solutions for a class of semilinear elliptic and parabolic equations, *SIAM J. Math. Anal.* **31** (1999), 1-18.
- [5] Y. Du, L. Ma; Logistic type equations on \mathbb{R}^N by a squeezing method involving boundary blow-up solutions, *J. London Math. Soc.* **64** (2001), 107-124.
- [6] J. García-Melián; A remark on uniqueness of large solutions for elliptic systems of competitive type, *J. Math. Anal. Appl.*, **331** (2007) 608-616.
- [7] J. García-Melián, J. D. Rossi; Boundary blow-up solutions to elliptic systems of competitive type, *J. Differential Equations*, **206** (2004) 156-181.
- [8] B. Guerch, L. Véron; Local properties of stationary solutions of some nonlinear singular Schrödinger equations, *Revista Matemática Iberoamericana*, **7** (1991), 65-114.
- [9] H. Li, M. Wang; Existence and uniqueness of positive solutions to the boundary blow-up problem for an elliptic system, *J. Differential Equations*, **234** (2007) 246-266.
- [10] J. López-Gómez; The boundary blow-up rate of large solutions, *J. Differential Equations*, **195** (2003), 25-45.
- [11] J. López-Gómez; Uniqueness of large solutions for a class of radially symmetric elliptic equations, in: S. Cano-Casanova, J. López-Gómez, C. Mora-Corral (Eds.), *Spectral Theory and Nonlinear Analysis with Applications to Spatial Ecology*, World Scientific, (2005) 75-110.
- [12] M. Marcus, L. Véron; Existence and uniqueness results for large solutions of general nonlinear elliptic equations, *J. Evol. Equ.*, **3** (2003), 637-652.
- [13] C. Pao; *Nonlinear parabolic and elliptic equations*, Plenum, New York, 1992.
- [14] L. Wei, Z. Feng; Isolated singularity for semilinear elliptic equations, *Discrete Contin. Dyn. Syst.*, **35** (2015), 3239-3252.

LEI WEI

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGSU NORMAL UNIVERSITY, XUZHOU 221116, CHINA
E-mail address: wlxznu@163.com

XIYOU CHENG

SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU 730000, CHINA
E-mail address: chengxy@lzu.edu.cn

ZHAOSHENG FENG

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS-RIO GRANDE VALLEY, EDINBURG, TX 78539, USA
E-mail address: zhaosheng.feng@utrgv.edu