University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Biology Faculty Publications and Presentations

College of Sciences

10-21-2021

Review of the Invasive Forage Grass, Guinea Grass (*Megathyrsus maximus*): Ecology and Potential Impacts in Arid and Semi-Arid Regions

Pushpa Soti

The University of Texas Rio Grande Valley, pushpa.soti@utrgv.edu

Vanessa Thomas

The University of Texas Rio Grande Valley

Follow this and additional works at: https://scholarworks.utrgv.edu/bio_fac

Part of the [Biology Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Soti, P. & Thomas, V. (2021) Review of the invasive forage Grass, Guinea grass (*Megathyrsus maximus*): Ecology and potential impacts in arid and semi-arid regions. *Weed Research*, 00, 1– 7. <https://doi.org/10.1111/wre.12512>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

**Review of the Invasive Forage Grass, Guinea Grass (*Megathyrsus maximus*): Ecology and
Potential Impacts in Arid and Semi-Arid Regions**

Pushpa Soti¹, Vanessa Thomas¹

¹Biology Department, University of Texas Rio Grande Valley, Edinburg, Texas 78539

Correspondence

Pushpa Soti, Biology Department, University of Texas Rio Grande Valley, Edinburg, Texas
78539

Pushpa.soti@utrgv.edu

Abstract

Several introduced African grasses are known to present recurring patterns of invasiveness and cause a severe impact on the diversity and functioning of ecosystems worldwide. *Megathyrsus maximus* (Guinea grass), a forage grass species native to South Africa, is reported to be highly invasive and pose a serious threat to native biodiversity in the introduced range. Despite the severe ecological threats posed by *M. maximus* worldwide, there is a dearth of information on the ecological and agroecological impact of *M. maximus* when growing in unintended areas. In this review, we present general information on *M. maximus*, its distribution and ecological threats it poses, particularly in arid and semi-arid regions. We highlight the gaps in current knowledge on the impact on recipient communities, challenges in effective management, and potential impacts due to climate change, particularly changes in rainfall patterns. We also stress the need for public awareness about the threats posed by *M. maximus* to prevent its invasion in unintended areas.

Keywords: biological invasion, African forage grasses, semi-arid subtropics, habitat degradation, Guinea grass

Introduction

With the introduction of species outside their native range, humans have caused a significant impact on the composition of biological communities worldwide. While a significant portion of introduced species do not get established in proportions that can have ecological impacts, a few become highly successful in invading recipient habitats. These invasive non-native species pose a serious threat to native species and potentially alter the ecosystem functions. Invasive plants are known to threaten biodiversity, reduce carbon storage, and influence the fundamental ecosystem processes such as fire regimes and nutrient cycling. Invasive species not only pose considerable harm to the native ecosystem and biodiversity but also have a significant economic impact. For example, Pimentel *et al.* (2005) estimated the annual cost of invasive plants in the United States to be at least US\$27B. A recent study by Diagne *et al.* (2021) reported the total cost of biological invasion world-wide to be a minimum of US \$1.288 trillion (2017 dollars) over the past few decades. While the high cost of invasive species control is one of the challenges faced by land managers, researchers have also acknowledged the social dimensions of invasive species management (Pimentel *et al.* 2005). This challenge is further compounded when invasive plant species have commercial value. For example, managing invasive grasses with agronomic value for farmers results in conflicts between farmers who want to exploit them as grazing grasses and conservationists who are concerned about ecological impacts (CABI 2019).

Invasive non-native grasses, originally introduced as forage grasses, are known to cause a significant impact on the functioning and stability of ecosystems (D'antonio and Vitousek, 1992). They also pose a threat to agriculture as major agronomic weeds (Parker *et al.* 2013). Invasive grasses of African origin are particularly known to cause a severe impact on the

diversity and functioning of ecosystems worldwide. These grasses have evolved under the high pressure of herbivory (Cerling *et al.* 2015) and adapted to a wide range of environmental conditions (Baruch, 1994) which gives them a competitive advantage against the native plants in terms of colonizing ruderal habitats. The life history traits (e.g., high growth rates and tolerance to herbivory, soil nutrient status, pH, and salinity) that make them valuable as forage grasses are also the ones that promote invasiveness in these grasses (Overholt and Franck, 2017).

Here we present the ecology, economic and ecological threats, and challenges in the management of *Megathyrsus maximus*, [Jacq.] B.K. Simon & S.W.L. Jacobs (Poaceae), previously *Panicum maximum* and *Urochloa maxima* [Jacq.] (Guinea grass), introduced to the tropics and subtropics as a forage grass. In the introduced regions, *M. maximus* has escaped from the cultivated rangelands and invaded disturbed sites, roadsides, untended areas, and grazing pastures at alarming rates. Despite the severe ecological threats posed by *M. maximus*, there is limited information on the ecology of and potential threats posed by *M. maximus* in the invaded regions, particularly in the tropical and sub-tropical regions around the globe, where it poses a significant threat in both agricultural fields and natural areas. The aim of this review is to highlight the potential threats posed by *M. maximus* in the introduced range if the grass grows out of confinement in ranches and pastures and infests nearby areas.

Origins and Distribution

Megathyrsus maximus, a forage grass native to tropical and sub-tropical Africa, was introduced across Asia, Europe, North America, and South America for hay and silage production but has caused significant ecological impacts. *Megathyrsus maximus* has become an invasive species in

tropical areas and warm temperate areas including the United States, India, Australia, and Brazil (Daehler *et al.* 1998; Sarkar *et al.* 2018) (Fig. 1). By 1915, *M. maximus* was present in the United States, Mexico, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Trinidad, Bermuda, Bahama, Cuba, Jamaica, and Haiti (Hitchcock and Chase, 1917). In the United States, *M. maximus* was first introduced to Florida and across the Gulf Coast in the early 1800s as grazing fodder for cattle and sheep, which then further spread into Southern Mexico (Vasey, 1887). By 1907, *M. maximus* was reported to grow along the coast of Alabama, Mississippi, and Texas as a forage grass. *Megathyrsus maximus* was cited to have been growing along the bank of the Guadeloupe river in Texas as early as 1894 (Arthur, 1894). It is now a prominent invasive plant in south Texas (Soti *et al.* 2020) and all major islands of Hawaii (Ammond *et al.* 2013).

Morphology and Seed Biology

Megathyrsus maximus is a deep-rooted C4 perennial bunchgrass. It grows in erect clumps with a clump radius ranging from 0.21 m to 2.89 m. Stems are cylindrical and 2.5m -3.5m tall (Aganga and Tshwenyane, 2004; Everitt, 2011) with a slightly flattened base. However, the height is dependent on other habitat conditions such as soil moisture, nutrients, shade, etc. Stems are streaked with white wax at the nodes and internodes with leaf blades growing from the lower nodes (Moore, 2010). Leaf-blades are clustered, 20-35cm long and 7-20mm wide, with few appressed hairs. When the leaves are older, the ends curl and dry (Gould, 1975; Everitt, 2011). Roots are dense and fibrous with extensive root hairs near the surface but continue to grow deeper, up to 4.5m (Sumiyoshi, 2012). The dense rhizomes and roots, which can grow up to 1m

deep, allow *M. maximus* to survive drought conditions (Aganga and Tshwenyane, 2004) and tolerate fire.

Reproduction occurs through seed as well as vegetative propagation. It is a prolific seed producer, with each plant producing up to 9000 seeds, however, seed yields are low due to seed shattering and small seed size (Sidhu, 1992). While plant biomass is reported to be significantly higher under shade, seed production is reported to be low (Sidhu, 1992). The germination rate of *M. maximus* seeds is reported to be relatively low (Mishra *et al.* 2008). However, the seed viability may be well over 80% if they are dried gradually to 10% moisture (Muir and Jank 2004). The seeds have been reported to experience dormancy for more than 3 years. Optimal seeding depth for *M. maximus* varies by soil type, 1cm in heavy soils and 1.5 cm in sandy soils (Muir and Jank 2004).

Habitat preferences

In its native range, subtropical southern Africa, *M. maximus* is adapted to grow under trees. It is reported to grow well under 25-50% shading, but growth declines at 75% shading (Malaviya *et al.* 2020). Under shaded conditions, *M. maximus* is reported to have a higher nitrogen concentration in the tissue (Paciullo *et al.* 2017). In south Texas, it does well under both shade and open canopy (authors' personal observation). This could potentially explain high *M. maximus* growth under mesquite trees, a leguminous plant. *Megathyrsus maximus* tolerates a wide range of temperature, 12.2 – 27.8°C. The optimum temperature for seed germination is estimated at 19.1-22.9°C but plant growth and biomass accumulation are higher in higher temperatures, with temperature having a strong positive correlation with root biomass (Muir and Jank, 2004). *Megathyrsus maximus* grown under high temperatures is also reported to have a

strong association with mycorrhizal fungi, leading to higher phosphorus uptake (Řezáčová *et al.* 2018).

Megathyrsus maximus is generally reported as a drought-tolerance species. However, soil moisture is reported as the major limiting factor for *M. maximus* growth. It grows well in areas with a total annual rainfall of 87-100 cm and grows moderately in drier soils. Under low soil moisture conditions, leaf biomass production declines significantly (Viciedo *et al.* 2019).

Megathyrsus maximus is known to grow well in a wide range of soil conditions. It prefers well-drained light-textured soils, sandy loams, or loams (Holm *et al.* 1977). *Megathyrsus maximus* is reported to tolerate seasonal inundation and the seeds can survive some flooding, but prolonged water logging can reduce seed viability and germination rate (Muir, 2004). In Malaysia, *M. maximus* is reported to grow on peat (Gajaweera, *et al.* 2011), while in Sri Lanka, it is reported to do well in low humic gley soils with very high-water holding capacity. *Megathyrsus maximus* also has a wide pH tolerance range, with optimal growth at soil pH 5-7. In south Texas, it grows in soils with pH greater than 8, while in Sri Lanka it grows in pH 5.5-7.7, and in Malaysia in 3.0-3.5 (Chew *et al.* 1980). Though it has a wide pH tolerance range, biomass production in *M. maximus* is reported to decline in soil pH >8 and <4 (Bernardes *et al.* 2018). It has high nitrogen demand and is highly competitive in nitrogen-rich soils, producing higher biomass than the cooccurring natives.

Ecological impact

Megathyrsus maximus invades both agricultural fields and natural areas, causing a significant impact on the ecosystem functioning and processes by altering the fire regime and soil quality as well as attracting pests and diseases of crops (Mantoani *et al.* 2016). It has been reported to be a

major pest in both annual and perennial crops such as rice, corn, sugarcane, coffee, citrus, and other fruit orchards causing a major reduction in crop yield (Table 1). *Megathyrsus maximus* has been associated with agronomic pests such as *Bipolaris yamadae*, a leaf spot disease infecting sugarcane, serving as a refuge during the otherwise fallow season.

Not only is *M. maximus* fire resistant but it is also reported to alter fire regime in the dry tropical forests of Hawaii (Ellsworth *et al.* 2014) and other tropical and subtropical landscapes. Tall *M. maximus* plants growing under trees add a high fuel load and can act as fire ladders carrying fire from the surface to tree canopies during the dry season causing lasting damage in the invaded systems (Best, 2006). Because *Megathyrsus maximus* is fire-tolerant and can rapidly regenerate from rhizomes after fires, it creates a positive feedback loop favoring its own growth in post-fire, high nutrient ash beds (Aganga and Tshwenyane, 2004). In Queensland, Australia, the dense tussocks of *M. maximus* growing along rivers and floodplains are known to outcompete the native species and displace them (Calvert 1998). In south Texas, the native plant restoration project in the Tamaulipan thornscrub has been significantly impacted by the extensive invasion by *M. maximus*. *M. maximus*, which can grow up to 2 meters tall, can overgrow and shade out the transplanted seedlings of native plants and outcompete them (Dick 2015).

While there is not much information on the impact of *M. maximus* on wildlife, it has been reported to degrade the northern bobwhite (*Colinus virginianus*) habitat in Texas. It also displaces native seed producing plants eaten by Gambel's quail (*Callipepla gambelii*) and other bird species (Kuvlesky *et al.* 2012). In addition, the shift in fire regime causes a decline in native arthropod communities in the habitats nearby (Warren *et al.* 1987). In Puerto Rico, *M. maximus*

is known to cause a decline in ground-dwelling insects, while in Australia, it is reported to reduce the larval survival rates of *Mycalesis* spp butterflies feeding on its leaves (See Table 1).

Table 1. Summary of ecological and agronomic impacts of *M. maximus* in the introduced range.

Location/Region	Ecological and Agronomic Impacts	References
Australia	Reduction in the larval survival rates of <i>Mycalesis</i> spp butterflies.	Braby <i>et al.</i> 1995
Argentina	Major weed in sugar cane fields leading up to 60% crop loss.	Cabrera <i>et al.</i> 2020
Brazil	Aggressive invader of annual and perennial crops, including rice, sugarcane, coffee, citrus, and other fruit orchards.	Durigan, 1992
Ecuador	Reduction in biodiversity of the Northern Ecuadorian Amazon area.	González <i>et al.</i> 2021
India	Host of a major pest, fall armyworm (<i>Spodoptera frugiperda</i>)	Maruthadurai and Ramesh, 2020
Sri Lanka	Invades natural areas such as forests and scrublands and disturbed degraded lands negatively impacting forestry and agriculture.	Weerawardane and Dissanayake, 2005
Hawaii	Reduction of native grasses and woody plant communities. Adds fuel to brush fires.	Cabin <i>et al.</i> 2002; Ellsworth, 2014

	Pose a threat to crops such as <i>Jatropha curcas</i> i.e	
	Barbados nut directly by influencing the fire regime	
	and indirectly by changing soil nutrient status.	
Florida, Texas,	Major weed in cotton and sugarcane.	Overholt and
and Louisiana	Serves as an alternate host for sugarcane aphid	Franck, 2017;
	<i>Melanaphis sacchari</i> .	Schenck and
	Serves as host for <i>Bipolaris yamadae</i> , leaf spot disease,	Lehrer, 2000; de
	which infects sugar cane crops.	Souza <i>et al.</i> 2019
	Degradation of the northern bobwhite <i>Colinus</i>	Adhikari <i>et al.</i>
	<i>virginianus</i> habitat.	2020; Moore, 2010
Puerto Rico	Decrease in the abundance of ground-dwelling	Moreno <i>et al.</i> 2014
	arthropods.	

168

169 **Economic and Environmental Value**

170 *Megathyrsus maximus* was universally introduced as a fodder species for its high protein content
171 and high tolerance to grazing and environmental stresses (Habermann *et al.* 2019). Because it is
172 one of the most productive forage grasses and highly palatable to cattle, it is frequently planted
173 by ranchers. Since it is a perennial bunchgrass with dense root growth, it has the potential to
174 reduce soil erosion (Maass *et al.* 1988; Mishra *et al.* 2008) and add soil organic matter.
175 *Megathyrsus maximus* has also been reported to be a moderate metal accumulator and has the
176 potential to be used as a phytoremediation/phytoextraction candidate in soil and wastewater
177 treatment projects (Olatunji *et al.* 2014, de Sousa *et al.* 2019, Anigbogu *et al.* 2020). In low
178 rainfall areas in Africa, *M. maximus* mulch is used as a drought management strategy (Wade and

Sahchez, 1983; Manu *et al.* 2017). In addition, *M. maximus* incorporated into the corn-legume cropping cycle is reported to increase soybean yields, improve forage quality, minimize nutrient loss, and thus maintain soil fertility in tropical conditions (Costa *et al.* 2021). *Megathyrus maximus* can also potentially host predatory arthropods including earwigs and spiders and could be utilized as a trap plant in maize fields to reduce spotted stem borer, *Chilo partellus*, eggs and larva (Koji *et al.* 2017).

Climate Change and Range Expansion Potential

Given the agronomic value of *M. maximus*, human mediated dispersal and propagule pressure are two major factors for its range expansion. However, climate change, which is projected to influence the rainfall pattern and temperatures leading to increased temperatures and prolonged drought periods, particularly in the sub-tropics, can also influence the distribution of *M. maximus* in this region. While *M. maximus* is reported to be tolerant to drought and high temperatures, there is some evidence that the above ground biomass growth is limited by soil moisture levels (Viciedo *et al.* 2019). These results show a mixed outcome for *M. maximus* under climate change scenarios. It can potentially both increase and or decrease suitable habitats for *M. maximus*. Under increasing temperatures, combined with drought conditions, *M. maximus* might reduce its expansion in natural areas in the arid and semi-arid regions. There is also a possibility of decrease in habitat suitability in arid and subtropical regions as well as the northward expansion of the species. However, irrigated agricultural fields, which are rich in soil nitrogen, are at a higher risk of *M. maximus* invasion in the topical, subtropical, and warm temperature regions (Kariyawasam *et al.* 2021). There is also a possibility of decrease in the suitable habitats in the arid and semi-arid tropics and subtropics and northward expansion. Thus,

further species distribution models projecting the potential response of *M. maximus* to changes in rainfall and temperature could be important in developing long-term management plans.

Management

The characteristics of *M. maximus*, such as high growth rate and tolerance to heavy grazing, shade, drought, salinity, and soil pH, which make it preferred forage grass species, also make it an aggressive invader in non-target habitats. In addition, prolific seed production and ability to rapidly regrow from rhizomes after fire make this species difficult to manage in the arid and semi-arid regions where prescribed burning is typically used for invasive species control (Johnson and Di Tomaso, 2006). Mechanical removal/mowing is reported to be ineffective as the plant can grow back from rhizomes. At a local scale, when the growth is limited, manual removal can be effective, but in larger areas it is expensive and labor intensive. Furthermore, because of its high agronomic value, complete eradication of *M. maximus* from introduced regions is impossible and/or highly controversial. Clearly, there is no single strategy to effectively manage this invasive grass. Several efforts to introduce biological control agents for *M. maximus* management have had mixed results. While the fungal pathogens *Dreschlera gigantea*, *Exserohilum rostratum*, and *E. longirostratum* have shown promising results in managing *M. maximus* in sugarcane fields in Florida (Chandramohan *et al.* 2004), a recent effort to introduce stem boring moths, *Buakea kaeuae* Moyal *et al.*, which is specific to small *M. maximus* of south-central Kenya, was reported to be unsuccessful (Vacek *et al.* 2021). Along with biocontrol, treating with 1% glyphosate is reported to be effective *M. maximus* management (Smith *et al.* 2012). However, there are conflicting reports on successful management with glyphosate treatment. While treatment with glyphosate only is reported to be effective for spot

control when the plants are at a younger stage, glyphosate mixed with flazasulfuron is reported to provide up to 95% *M. maximus* control in citrus groves (Singh *et al.* 2012). In south Texas, management strategies have generally involved a combination of cattle grazing and prescribed burning. It has been reported to reduce *M. maximus* density and increase native plant species richness (Ramirez-Yanez, 2005; Ramirez-Yanez *et al.* 2007). Thus, effective management of *M. maximus* in the introduced range, particularly in the semi-arid tropics and subtropics, can be achieved through a combination of public awareness and integrated pest management including cattle grazing, post emergent herbicides, and prescribed burning followed by pre-emergent herbicides.

Conclusions

M. maximus is a forage grass species with high agronomic value, widely distributed in the tropics and sub-tropics where it is now considered as a highly invasive species. Given its high tolerance for biotic and abiotic stresses, it is likely to further expand its distribution. Because of its agronomic value and extent of spread, complete eradication of *M. maximus* from the introduced range is not desirable nor possible. Thus, the primary strategy for *M. maximus* management should be to reduce its impact on native communities and crops in agricultural fields. Site specific strategies based on the habitat environmental conditions need to be developed for the effective management of *M. maximus*. In areas where *M. maximus* has not extensively invaded cropping fields and native grasslands, it can be managed by well-planned grazing. In areas where *M. maximus* is already established, management can potentially be achieved through the integration of biocontrol (including planned grazing) as well as cultural, chemical, and mechanical methods. While the effectiveness of habitat manipulation has mixed results and is

site dependent (Huston 2004), its invasion in agricultural fields can be prevented and/or minimized through proper management of nitrogen fertilizer and precision irrigation. Further comprehensive studies on seed viability, germination, and site-specific *M. maximus* physiology and growth analysis are necessary for effective management. In addition, habitat modeling, incorporating habitat preferences to identify potential impacts of changes in climatic variables, could be important in preventing further spread of *M. maximus* while still allowing for economic uses where feasible.

Acknowledgements

We thank Dr. John Goolsby, Research Entomologist, USDA- ARS, for his input during the preparation of this manuscript.

References

- Adhikari, A., Wang, X., Lane, B., Harmon, P.F. and Goss, E., 2020. First report of *Bipolaris yamadae* leaf spot disease on Guinea grass (*Panicum maximum*) in Florida. *Plant Disease*, (ja),
- Aganga, A.A. and Tshwenyane, S., 2004. Potentials of guinea grass (*Panicum maximum*) as forage crop in livestock production. *Pakistan journal of nutrition*, 3(1), pp. 1-4.
- Ammond, S.A., Litton, C.M., Ellsworth, L.M. and Leary, J.K., 2013. Restoration of native plant communities in a Hawaiian dry lowland ecosystem dominated by the invasive grass *Megathyrsus maximus*. *Applied Vegetation Science*, 16(1), pp. 29-39.
- Anigbogu, N.M., Agida, C.A., Chukwurah, C.B., Afam-Ibezim, E.M. and Efuribe, B.E., 2020. Economic impact and growth rate of Red Sokoto goats fed treated municipal organic waste and grass/legume mixtures. *Nigerian Journal of Animal Science and Technology (NJAST)*, 3(2), pp. 253-265.
- Arthur., 1984 Botanical Explorations in Southern Texas during the Season of 1894. Lancaster, PA: New Era Printing House, Herbarium of Franklin and Marshal College. Vol. 01, pp. 11.
- Baruch, Z. 1994. Responses to drought and flooding in tropical forage grasses: I. Biomass allocation, leaf growth and mineral nutrients. *Plant and Soil* 164, pp. 87–96.
- Bernardes, T.F., Daniel, J., Adesogan, A.T., Mcallister, T.A., Drouin, P., Nussio, L.G., Huhtanen, P., Tremblay, G.F., Bélanger, G. and Cai, Y., 2018. Silage review: Unique challenges of silages made in hot and cold regions. *Journal of dairy science*, 101(5), pp. 4001-4019.

Best, C., 2006. Fighting weeds with weeds: battling invasive grasses in the Rio Grande delta of Texas. *Invasive plants on the move: controlling them in North America*, 307.

Braby, M.F., 1995. Larval and adult food plants for some tropical satyrine butterflies in northern Queensland. *Australian Entomologist, The*, 22(1), pp. 5-13.

Cabrera, D., Sobrero, M.T., Pece, M. and Chaila, S., 2020. Effect of environmental factors on the germination of *Megathyrus maximus*: an invasive weed in sugarcane in Argentina. *Planta Daninha*, 38.

Cabin, R.J., Weller, S.G., Lorence, D.H., Cordell, S., Hadway, L.J., Montgomery, R., Goo, D. and Urakami, A., 2002. Effects of light, alien grass, and native species additions on Hawaiian dry forest restoration. *Ecological Applications*, 12(6), pp. 1595-1610.

Calvert, G. 1998. Guinea grass. Paper presentation at the Society for Growing Australian Plants: Queensland Regional Conference. *Townsville*. 28 June-5 July, 1998.

Cerling, T., Andanje, S., Blumenthal, S., Brown, F., Chritz, K., Harris, J., Hart, J., Kierra, F., Kaleme, P., Leakey, L., and Leakey, M., 2015. Dietary changes of large herbivores in the Turkana Basin, Kenya from 4 to 1 Ma. *PNAS* 112, pp.11467–11472.

Chandramohan, S., Shine, J.M., Duchrow, M.J., Roskopf, E.N. and Charudattan, R., 2004. A bioherbicide system to manage Guinea grass (*Panicum maximum* Jacq.) in sugarcane in Florida. *Sugar Cane International*, pp. 28-29.

Chew, W.Y., Williams, C.N., Ismail, L. and Ramli, K., 1980. Effects of liming and soil pH on Guinea grass (*Panicum maximum*) in Malaysian tropical oligotrophic peat. *Experimental Agriculture*, 16(3), pp. 263-267.

Costa, N.R., Andreotti, M., Crusciol, C.A.C., Pariz, C.M., Bossolani, J.W., Castilhos, A.M.D., Nascimento, Carlos Antonio Costa Do, Lima, César Diagne, C., Leroy, B., Vaissière, A.

305 C., Gozlan, R. E., Roiz, D., Jarić, I., ... and Courchamp, F. (2021). High and rising
 306 economic costs of biological invasions worldwide. *Nature*, 592(7855), 571-576.

307 Daehler, C., and Carino, D., 1998. Recent replacement of native pili grass (*Heteropogon*
 308 *contortus*) by invasive African grasses in the Hawaiian Islands. *Pacific Science*, vol. 52,
 309 no. 3: pp. 220-227.

310 D'antonio, C. M., and Vitousek, P.M., 1992. Biological invasions by exotic grasses, the
 311 grass/fire cycle, and global change. *Annual review of ecology and systematics* 23.1, pp.
 312 63-87.

313 De Souza, M.F. and Davis, J.A., 2019. Determining potential hosts of *melanaphis sacchari*
 314 (Hemiptera: Aphididae) in the Louisiana agroecoscape. *Environmental*
 315 *Entomology*, 48(4), pp. 929-934.

316 Dick, K.N. 2015. Restoring Semi-arid Thornscrub Forests: Seedling Growth and Survival in
 317 Response to Shelter Tubes, Grass-Specific Herbicide, and Herbivore Exclosures. MS
 318 thesis. Brownsville: University of Texas at Brownsville

319 Johnson, D.W. and Di Tomaso, J.M. 2006. The use of fire as a tool for controlling invasive
 320 plants. *Cal-IPC publication*, 1, pp. 56.

321 Durigan, J.C., 1992. Efeito de adjuvantes na calda e do estágio de desenvolvimento das plantas,
 322 no controle do capim-colonião (*Panicum maximum*) com glyphosate. *Planta*
 323 *Daninha*, 10(1-2), pp. 39-44.

324 Ellsworth, L.M., Litton, C.M., Dale, A.P. and Miura, T., 2014. Invasive grasses change
 325 landscape structure and fire behaviour in Hawaii. *Applied Vegetation Science*, 17(4), pp.
 326 680-689.

327 Everitt, J.H., Drawe, D.L., Little, C.R. and Lonard, R.I., 2011. *Grasses of South Texas: a guide*
328 *to their identification and value*. Texas Tech University Press Lubbock.

329 Gajaweera, C. J., Weerasinghe, B., and Seresinhe, T. 2011. Preliminary study on the adaptation
330 of wild Guinea grass (*Panicum maximum*) to different agro-climatic regions in
331 Hambanthota District. *Proceedings of International Forestry and Environment*
332 *Symposium*. Vol. 16.

333 Gould, F.W., 1975. *The grasses of Texas*. Texas A&M University, Texas Agricultural
334 Experiment Station.

335 Habermann, E., San Martin, Juca Abramo Barrera, Contin, D.R., Bossan, V.P., Barboza, A.,
336 Braga, M.R., Groppo, M. and Martinez, C.A., 2019. Increasing atmospheric CO₂ and
337 canopy temperature induces anatomical and physiological changes in leaves of the C₄
338 forage species *Panicum maximum*. *PLoS One*, 14(2), pp. e0212506.

339 Hitchcock, A. S., and Chase, A. 1917. Grasses of the west Indies. *Systematic investigations of*
340 *tropical American plants*.

341 Holm, L.G., Plucknett, D.L., Pancho, J.V. and Herberger, J.P., 1977. *The world's worst weeds*.
342 *Distribution and biology*. University press of Hawaii.

343 Huston, M., 2004 "Management strategies for plant invasions: manipulating productivity,
344 disturbance, and competition." *Diversity and distributions* 10.3, pp. 167-178.

345 Kariyawasam, C. S., Kumar, L., and Ratnayake, S. S., 2021. Potential risks of invasive alien
346 plant species on agriculture under climate change scenarios in Sri Lanka. *Current*
347 *Research in Environmental Sustainability*, 3, 100051.

348 Koji, S., Khan, Z.R. and Midega, C.A., 2007. Field boundaries of *Panicum maximum* as a
 349 reservoir for predators and a sink for *Chilo partellus*. *Journal of Applied*
 350 *Entomology*, 131(3), pp. 186-196.

351 Kuvlesky Jr, W. P., Brennan, L. A., Fulbright, T. E., Hernandez, F., DeMaso, S. J., Sands, J. P.,
 352 ... and Hardin, J. B. 2012. Impacts of invasive, exotic grasses on quail of southwestern
 353 rangelands: a decade of progress?. In National Quail Symposium Proceedings (Vol. 7,
 354 No. 1, p. 55).

355 Maass, J.M., Jordan, C.F. and Sarukhan, J., 1988. Soil erosion and nutrient losses in seasonal
 356 tropical agroecosystems under various management techniques. *Journal of Applied*
 357 *Ecology*, pp. 595-607.

358 Malaviya, D.R., Roy, A.K. and Kaushal, P., 2020. Comparative leaf anatomical studies reveal
 359 photosynthetic pathways type in guinea grass (*Panicum maximum* Jacq.) accessions and
 360 their interspecific lineage. *Current science*, 119(5), pp. 808.

361 Mantoani, M.C. and Torezan, J.M.D., 2016. Regeneration response of Brazilian Atlantic Forest
 362 woody species to four years of *Megathyrsus maximus* removal. *Forest Ecology and*
 363 *Management*, 359, pp. 141-146.

364 Manu, V., Whitbread, A. and Blair, G., 2017. Mulch effect on successive crop yields and soil
 365 carbon in T onga. *Soil Use and Management*, 33(1), pp. 98-105.

366 Maruthadurai, R. and Ramesh, R., 2020. Occurrence, damage pattern and biology of fall
 367 armyworm, *Spodoptera frugiperda* (JE smith) (Lepidoptera: Noctuidae) on fodder crops
 368 and green amaranth in Goa, India. *Phytoparasitica*, 48(1), pp. 15-23.

369 Mishra, S., Sharma, S. and Vasudevan, P., 2008. Comparative effect of biofertilizers on fodder
370 production and quality in guinea grass (*Panicum maximum* Jacq.). *Journal of the science*
371 *of food and agriculture*, 88(9), pp. 1667-1673.

372 Moore, S.F., 2010. Effects of Guinea grass on northern bobwhite habitat use, *Masters Abstracts*
373 *International* 2010.

374 Muir, J.P. and Jank, L., 2004. Guinea grass. *Warm-season (C4) grasses*, 45, pp. 589-621.

375 Olatunji, O.S., Ximba, B.J., Fatoki, O.S. and Opeolu, B.O., 2014. Assessment of the
376 phytoremediation potential of *Panicum maximum* (guinea grass) for selected heavy metal
377 removal from contaminated soils. *African Journal of Biotechnology*, 13(19), pp. 1979-
378 1984.

379 Overholt, W.A. and Franck, A.R., 2017. The invasive legacy of forage grass introductions into
380 Florida. *Natural Areas Journal*, 37(2), pp. 254-264.

381 Paciullo, D., Gomide, C.D.M., Castro, C.D., Maurício, R.M., Fernandes, P.B. and Morenz, M.,
382 2017. Morphogenesis, biomass, and nutritive value of *Panicum maximum* under different
383 shade levels and fertilizer nitrogen rates. *Grass and Forage Science*, 72(3), pp. 590-600.

384 Parker., 2013. Do invasive species perform better in their new ranges? *Ecology* 94.5 pp. 985-994.

385 Pimentel, D., Zuniga, R., and Morrison, D. 2005. Update on the environmental and economic
386 costs associated with alien-invasive species in the United States. *Ecological*
387 *economics*, 52(3), 273-288.

388 Ramirez-Yanez, L.E., 2005. Prescribed fire and intensive grazing to control invasive guinea
389 grass on native pastures in South Texas. Texas A&M University-Kingsville.

390 Ramirez-Yanez, L.E., Ortega-S, J.A., Brennan, L.A., and Rasmussen, G.A., 2007. Use of
391 prescribed fire and cattle grazing to control Guinea grass, *Proceedings of the 23rd Tall*

392 *Timbers Fire Ecology Conference: Fire in Grassland and Shrubland Ecosystems*, Tall
393 Timbers Research Station Tallahassee, FL, USA, pp. 240-245.

394 Řezáčová, V., Slavíková, R., Zemková, L., Konvalinková, T., Procházková, V., Šťovíček, V.,
395 Hršelová, H., Beskid, O., Hujslová, M. and Gryndlerová, H., 2018. Mycorrhizal
396 symbiosis induces plant carbon reallocation differently in C3 and C4 *Panicum*
397 grasses. *Plant and Soil*, 425(1), pp. 441-456.

398 Sarkar, D., Meitei, C. B., Das, A., Ghosh, P. K., & Mandal, B. 2018. Changes in soil organic
399 carbon pools in a long-term trial with perennial fodder crops in acid soils of north-east
400 India. *Grass and Forage Science*, 73(2), 473-481.

401 Schenck, S. and Lehrer, A.T., 2000. Factors affecting the transmission and spread of sugarcane
402 yellow leaf virus. *Plant Disease*, 84(10), pp. 1085-1088.

403 Smith, H., Ferrell, J., and Sellers, B., 2012. Identification and Control of Johnsongrass,
404 Vaseygrass, and Guinea Grass in Pastures. EDIS, 2012(8).

405 Soti, P., Goolsby, J. A., and Racelis, A., 2020. Agricultural and environmental weeds of South
406 Texas and their management. *Subtropical Agriculture and Environments*, 71.

407 Sidhu, B.S., 1992. "Inter-relationships among total crude protein and digestible dry matter
408 production and their component traits in Guinea grass." *Euphytica* 64.1, pp. 59-63.

409 Sumiyoshi, Y. 2012. *Belowground impact of napier and Guinea grasses grown for biofuel*
410 *feedstock production* (Doctoral dissertation, [Honolulu]: [University of Hawaii at
411 Manoa], [December 2012]).

412 Vacek, A. T., Goolsby, J. A., Calatayud, P. A., Le Ru, B., Musyoka, B., and Kariyat, R. R. 2021.
413 Importation and Preliminary Evaluation of the Stem Boring Moth *Buakea kaeuae* as a

414 Potential Biological Control Agent of Invasive Guinea grass, *Megathyrsus*
 415 *maximus*. *Southwestern Entomologist*, 46(1), 257-260.
 416 Vasey, G. 1887. Grasses of the South: A Report on Certain Grasses and Forage Plants for
 417 Cultivation in the South and Southwest. *Washington Printing Office*, pp. 9-10.
 418 Vicedo, D.O., De Mello Prado, R., Martínez, C.A., Habermann, E. and De Cássia Piccolo, M.,
 419 2019. Short-term warming and water stress affect *Panicum maximum* Jacq.
 420 stoichiometric homeostasis and biomass production. *Science of the Total*
 421 *Environment*, 681, pp. 267-274.
 422 Wade, M.K. and Sanchez, P.A., 1983. Mulching and Green Manure Applications for Continuous
 423 Crop Production in the Amazon Basin 1. *Agronomy Journal*, 75(1), pp. 39-45.
 424 Warren, S.D., Scifres, C.J. and Teel, P.D., 1987. Response of grassland arthropods to burning: a
 425 review. *Agriculture, Ecosystems & Environment*, 19(2), pp. 105-130.
 426 Weerawardane, N. and Dissanayake, J., 2005. Status of forest invasive species in Sri Lanka. *The*
 427 *unwelcome guests*, pp. 114-120.