

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

School of Mathematical and Statistical  
Sciences Faculty Publications and  
Presentations

College of Sciences

---

1-22-2014

## On cubic multisections of Eisenstein series

Andrew Alaniz

*The University of Texas Rio Grande Valley*

Timothy Huber

*The University of Texas Rio Grande Valley*, [timothy.huber@utrgv.edu](mailto:timothy.huber@utrgv.edu)

Follow this and additional works at: [https://scholarworks.utrgv.edu/mss\\_fac](https://scholarworks.utrgv.edu/mss_fac)



Part of the [Mathematics Commons](#)

---

### Recommended Citation

Alaniz, A., Huber, T. On cubic multisections of Eisenstein series. *Ramanujan J* 35, 391–403 (2014).  
<https://doi.org/10.1007/s11139-013-9540-9>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in School of Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

# ON CUBIC MULTISECTIONS OF EISENSTEIN SERIES

ANDREW ALANIZ AND TIM HUBER

ABSTRACT. A systematic procedure for generating cubic multisections of Eisenstein series is given. The relevant series are determined from Fourier expansions for Eisenstein series by restricting the congruence class of the summation index modulo three. We prove that the resulting series are rational functions of  $\eta(\tau)$  and  $\eta(3\tau)$ , where  $\eta$  is the Dedekind eta function. A more general treatment of cubic dissection formulas is given by describing the dissection operators in terms of linear transformations. These operators exhibit properties that mirror those of similarly defined quintic operators.

## 1. INTRODUCTION

We begin with an identity observed by F. Garvan and communicated to the authors by B.C. Berndt. The purpose of this note is to place this identity in a larger context.

**Theorem 1.1.**

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} d \right) q^n = \frac{3(q^3; q^3)_{\infty}^6}{(q; q)_{\infty}^2}, \quad |q| < 1. \quad (1.1)$$

Here, we employ the usual notation for the  $q$ -expansion of the Dedekind eta function defined by  $q^{-1/24}\eta(\tau) = \prod_{n=1}^{\infty} (1 - q^n) = (q; q)_{\infty}$ ,  $q = e^{2\pi i\tau}$ ,  $\text{Im } \tau > 0$ . In this paper we prove Garvan's identity and place (1.1) in the context of several infinite classes of relations for dissections of Eisenstein series. Some examples of formulas that ensue are

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} d^3 \right) q^n = (q; q)_{\infty}^8 + 3^4 q \frac{(q^3; q^3)_{\infty}^{12}}{(q; q)_{\infty}^4},$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} d^7 \right) q^n = 3 \cdot 43 (q; q)_{\infty}^{16} + 110 \cdot 3^6 q (q; q)_{\infty}^4 (q^3; q^3)_{\infty}^{12} + 41 \cdot 3^{10} q^2 \frac{(q^3; q^3)_{\infty}^{24}}{(q; q)_{\infty}^8},$$

and, if  $\left(\frac{\cdot}{3}\right)$  denotes the Jacobi symbol modulo three,

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} \left(\frac{d}{3}\right) \right) q^n = \frac{(q^3; q^3)_{\infty}^3}{(q; q)_{\infty}}, \quad (1.2)$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} \left(\frac{d}{3}\right) d^4 \right) q^n = -15 (q; q)_{\infty}^7 (q^3; q^3)_{\infty}^3 - 3^6 q \frac{(q^3; q^3)_{\infty}^{15}}{(q; q)_{\infty}^5}.$$

---

2010 *Mathematics Subject Classification.* Primary 11F11; Secondary 11F33.

*Key words and phrases.* Eisenstein series; cubic theta functions; cubic multisections; t-cores.

The preceding relations are similar to dissections for Eisenstein series of level five appearing in [9] involving Dirichlet characters modulo five. We have the following dissection formulas for the quintic Dirichlet character defined by  $\langle \chi(n) \rangle_{n=0}^4 = \langle 0, 1, -i, i, -1 \rangle$ .

**Theorem 1.2.**

$$\begin{aligned} \sum_{n=0}^{\infty} \left( \sum_{d|5n+4} \chi(d) \right) q^n &= \frac{i(q; q)_{\infty} (q^5; q^5)_{\infty}}{(q^2; q^5)_{\infty}^3 (q^3; q^5)_{\infty}^3}, & \sum_{n=0}^{\infty} \left( \sum_{d|5n+3} \chi(d) \right) q^n &= \frac{(1+i)(q^5; q^5)_{\infty}^2}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}}, \\ \sum_{n=0}^{\infty} \left( \sum_{d|5n+2} \chi(d) \right) q^n &= \frac{(1-i)(q^5; q^5)_{\infty}^2}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}}, & \sum_{n=0}^{\infty} \left( \sum_{d|5n+1} \chi(d) \right) q^n &= \frac{(q; q)_{\infty} (q^5; q^5)_{\infty}}{(q; q^5)_{\infty}^3 (q^4; q^5)_{\infty}^3}. \end{aligned}$$

The analogous dissection formulas we derive for Eisenstein of level three are consequences of a triple of fortuitous relations between generators for the relevant spaces of modular forms. To derive the expansions, we employ well known properties of the Borwein's cubic theta functions  $a(q)$ ,  $b(q)$ , and  $c(q)$ , defined by

$$a(q) = \sum_{m,n=-\infty}^{\infty} q^{n^2+nm+m^2}, \quad b(q) = \sum_{m,n=-\infty}^{\infty} \omega^{n-m} q^{n^2+nm+m^2}, \quad (1.3)$$

$$c(q) = \sum_{m,n=-\infty}^{\infty} q^{(n+\frac{1}{3})^2+(n+\frac{1}{3})(m+\frac{1}{3})+(m+\frac{1}{3})^2}, \quad \omega = e^{2\pi i/3}. \quad (1.4)$$

**Theorem 1.3.**

$$a^3(q) = b^3(q) + c^3(q) \quad (1.5)$$

$$a(q) = a(q^3) + 2c(q^3), \quad b(q) = a(q^3) - c(q^3). \quad (1.6)$$

Equivalent formulations of all three identities appear in Ramanujan's notebooks [12]. The first proof of (1.5) was given by J. Borwein and P. Borwein [4]. Transformation formulas resulting from Theorem 1.3 appear in a different form in [3] and motivate cubic analogues of Jacobi's Principles of Duplication and Dimidiation [10]. Such formulas are subsumed in Ramanujan's "theories of elliptic functions to alternative bases" and termed Processes of Triplication and Trimidiation [3]. The utility of these transformations in studying the coefficients of modular forms is apparent in light of the fact that  $a(q)$  and  $c(q)$  serve as generators for the principal congruence subgroup of level three [2]. In fact, Sebbar [13] proved that the principal congruence subgroups of level three and five are two of precisely six congruence subgroups of  $SL(2, \mathbb{Z})$  whose graded ring of modular forms is a polynomial ring with two generators, each of degree one.

The remainder of the paper is organized as follows. In Section 2, we prove Garvan's identity (1.1) and generalize the dissection technique to Eisenstein series of arbitrary weight and cubic character. We identify classes of Eisenstein series whose cubic dissections are expressible as rational functions of the Dedekind eta function at argument  $q$  and  $q^3$ . These representations demonstrate general congruence properties for divisor sums and related arithmetic functions that are stated at the beginning of Section 2 and proven in the conclusion of Section 2. In Section 3 we provide explicit matrix representations for dissection operators on certain vector spaces. We use the operators to derive new congruence properties for the coefficients of corresponding eigenforms.

## 2. CUBIC DISSECTIONS FOR EISENSTEIN SERIES

In this section, we demonstrate that dissections formulas like (1.1) result from parameterizations for Eisenstein series in terms of the cubic theta functions  $a(q)$ ,  $b(q)$ , and  $c(q)$ . The combinatorial consequences of our work include several new results for twisted divisor sums and place a number of known results in context. In particular, the ensuing dissection formulas exhibit the following combinatorial interpretations.

**Theorem 2.1.** *Suppose  $\ell \in \mathbb{N}$  is odd and  $3^s \parallel \ell$ . Then for  $k \in \mathbb{N}$ ,*

$$\sum_{d|3k+2} \left(\frac{d}{3}\right) d^{2\ell} \equiv 0 \pmod{3^{s+1}}, \quad \sum_{d|3k+1} \left(\frac{d}{3}\right) d^{2\ell} \equiv c_3(k) \pmod{3^{s+1}}, \quad (2.1)$$

$$\sum_{d|3k+2} d^\ell \equiv 0 \pmod{3^{s+1}}, \quad \sum_{d|3k+1} d^\ell \equiv c_3(k) \pmod{3^{s+1}}, \quad (2.2)$$

where  $c_3(k)$  is the number of 3-core partitions of  $k$ ; i.e., the number of partitions of  $k$  satisfying the condition that no hook number in the Ferrers graph is divisible by 3.

If  $\ell = s = 0$  in the latter congruence of (2.1), we obtain an equality corresponding to (1.2), appearing in Granville and Ono's work [8]. The left congruences from (2.2) are proven without reference to Eisenstein series in [11]. The remaining congruences from Theorem 2.1 appear to be new. At the end of this section, we show that each of the generating functions for (2.1)–(2.2) reduce modulo  $3^{s+1}$  to dissected Eisenstein series of low weight. Moreover, our work results in explicit eta function expansions for the generating functions corresponding to the trisected divisor sums in Theorem 2.1. The product formulations follow from the product expansions [4] for  $b(q)$  and  $c(q)$

$$b(q) = \frac{(q; q)_\infty^3}{(q^3; q^3)_\infty}, \quad c(q) = 3q^{1/3} \frac{(q^3; q^3)_\infty^3}{(q; q)_\infty}. \quad (2.3)$$

To derive the aforementioned expansions for generating functions in terms of cubic theta functions, two classes of Eisenstein series will be appropriate to our discussion. The first are normalized Eisenstein series of weight  $k$  for the full modular group

$$E_{2k}(q) = 1 + \frac{2}{\zeta(1-2k)} \sum_{n=1}^{\infty} \frac{n^{2k-1} q^n}{1-q^n}, \quad (2.4)$$

where  $\zeta$  is the analytic continuation of the Riemann  $\zeta$ -function. We will also refer to the Hecke Eisenstein series associated with the Dirichlet character  $\chi$  modulo three

$$E_{k,\chi}(q) = 1 + \frac{2}{L(1-k, \chi)} \sum_{n=1}^{\infty} \chi(n) \frac{n^{k-1} q^n}{1-q^n}, \quad (2.5)$$

where  $L(1-k, \chi)$  denotes associated Dirichlet  $L$ -series. Since we build cubic Eisenstein expansions inductively, we require representations for low weight series from [7, 14]

$$E_4(q) = a^4(q) + 8ac^3(q), \quad E_6(q) = a^6(q) - 20a^3(q)c^3(q) - 8c^6(q), \quad E_{3,(\frac{\cdot}{3})}(q) = b^3(q).$$

In the proof of the next Theorem, we show that Garvan's formula (1.1) arises from a dissection of the Hecke Eisenstein series of weight two and trivial character  $\mathbf{1}$ . The technique applied here is representative of the methods used in the rest of the paper.

**Theorem 2.2.** Let  $\left(\frac{\cdot}{3}\right)$  denote the Jacobi symbol modulo three. Then

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} d \right) q^n = \frac{3(q^3; q^3)_{\infty}^6}{(q; q)_{\infty}^2}, \quad (2.6)$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} \left(\frac{d}{3}\right) \right) q^n = \frac{(q^3; q^3)_{\infty}^3}{(q; q)_{\infty}}, \quad \text{and} \quad \sum_{n=0}^{\infty} \left( \sum_{d|3n+2} \left(\frac{d}{3}\right) \right) q^n = 0. \quad (2.7)$$

*Proof.* To derive (2.6), we start with the cubic parameterization for the Eisenstein series of weight two [7, Theorem 11.10]

$$\frac{3}{2}E_2(q^3) - \frac{1}{2}E_2(q) = a^2(q). \quad (2.8)$$

Apply Theorem 1.3 to (2.8) to derive

$$\frac{3}{2}E_2(q) - \frac{1}{2}E_2(q^{1/3}) = a^2(q) + 4a(q)c(q) + 4c^2(q). \quad (2.9)$$

Because of the unique determination of the Maclaurin expansions of each side of (2.9), we may equate terms on each side with indices congruent to 2 modulo 3 to derive

$$-12 \sum_{n=0}^{\infty} \left( \sum_{d|3n+2} d \right) q^{(3n+2)/3} = 4c^2(q) = 36q^{2/3} \frac{(q^3; q^3)_{\infty}^6}{(q; q)_{\infty}^2}. \quad (2.10)$$

The product representations appearing in last equality follow from (2.3). A proof of (2.7) is likewise obtained by applying Theorem 1.3 to the identity [7, Equation (2.33)]

$$E_{1, \left(\frac{\cdot}{3}\right)}(q) = 1 + 6 \sum_{n=1}^{\infty} \binom{n}{3} \frac{q^n}{1 - q^n} = a(q). \quad \square$$

Since  $a(q) \equiv 1 \pmod{3}$ , we may equate terms on both sides of (2.9) of index congruent to one modulo three to obtain the  $\ell = 1$  case of the latter congruence in (2.2). By applying the triple product identity to the theta expansion for  $a(q)$  [5, Lemma 2.1]

$$a(q) = \theta_3(q)\theta_3(q^3) + \theta_2(q)\theta_2(q^3), \quad \theta_2(q) = \sum_{n=-\infty}^{\infty} q^{(n+\frac{1}{2})^2}, \quad \theta_3 = \sum_{n=-\infty}^{\infty} q^{n^2}, \quad (2.11)$$

we obtain a companion expansion to (2.6). Most product formulations for dissections involving the function  $a(q)$  are similarly unwieldy and will not be further studied here.

**Theorem 2.3.**

$$\begin{aligned} \sum_{n=0}^{\infty} \left( \sum_{d|3n+1} d \right) q^n &= \frac{(-q; q^2)_{\infty}^2 (-q^3; q^6)_{\infty}^2 (q^3; q^3)_{\infty}^3 (q^2; q^2)_{\infty} (q^6; q^6)_{\infty}}{(q; q)_{\infty}} \\ &\quad + \frac{4q(q^4; q^4)_{\infty} (q^{12}; q^{12})_{\infty} (q^3; q^3)_{\infty}^3}{(q^2; q^4)_{\infty} (q^6; q^{12})_{\infty} (q; q)_{\infty}}. \end{aligned}$$

In order to achieve the claimed rational expansions in  $\eta(q)$  and  $\eta(q^3)$  for dissections of higher weight Eisenstein series, we characterize Eisenstein series of level three whose trisections lie in the complex span of homogeneous polynomials in  $c(q)$  and  $b(q)$ .

**Lemma 2.4.** *If  $\bar{n}$  denotes the least positive residue class of  $n$  modulo 3, then the following series are, respectively, homogeneous polynomials in  $c(q)$  and  $b(q)$  over  $\mathbb{Q}$ :*

$$\sum_{k=0}^{\infty} \left( \sum_{d|3k+2\bar{n}} d^{2n-1} \right) q^{(3k+2\bar{n})/3}, \quad n \not\equiv 0 \pmod{3},$$

$$\sum_{k=0}^{\infty} \left( \sum_{d|3k+2\bar{n}+1} \left( \frac{d}{3} \right) d^{2n} \right) q^{(3k+2\bar{n}+1)/3}, \quad n \not\equiv 1 \pmod{3}.$$

*Proof.* An induction argument along with recursion formulas from Lemma 2.5 imply

$$E_{2n+1,(\frac{\cdot}{3})}(q) = f_{2n+1}(a(q), b(q)) \quad \text{and} \quad E_{2n,1}(q) = g_{2n}(a(q), b(q)) \quad (2.12)$$

for homogeneous polynomials  $f_{2n+1}$  and  $g_{2n}$  in  $a(q)$  and  $b(q)$  of degrees at most  $2n+1$  and  $2n$ , respectively. By replacing  $q$  by  $q^{1/3}$  and applying Theorem 1.3, we derive expansions for  $f_{2n+1}(a(q^{1/3}), b(q^{1/3}))$  and  $g_{2n}(a(q^{1/3}), b(q^{1/3}))$  as homogeneous polynomials in  $a(q)$  and  $c(q)$  with rational coefficients. From (1.5), and since we have assumed  $2n+1 \not\equiv 0 \pmod{3}$ , the  $\overline{2n+1}$ -dissection of  $f_{2n+1}(a(q^{1/3}), b(q^{1/3}))$  equals, for  $\lambda_k \in \mathbb{Q}$ ,

$$\sum_{k=0}^{\infty} \left( \sum_{d|3k+2\bar{n}+1} \left( \frac{d}{3} \right) d^{2n} \right) q^{\frac{3k+2\bar{n}+1}{3}} = \sum_{k=0}^{2n+1} \lambda_k (b^3(q) + c^3(q))^{\frac{2n+1-2\bar{n}+1-3k}{3}} c^{\overline{2n+1}+3k}(q).$$

Since  $m - \bar{m} \equiv 0 \pmod{3}$ , each exponent of  $(b^3(q) + c^3(q))$  in the last summand is a natural number. Therefore, the  $\overline{2n+1}$ -dissection of  $f_{2n+1}(a(q^{1/3}), b(q^{1/3}))$  is a homogeneous polynomial in  $b(q)$  and  $c(q)$  of degree at most  $2n+1$ . A similar calculation shows the  $\overline{2n}$ -dissection of the polynomial  $g_{2n}(a(q^{1/3}), b(q^{1/3}))$  is of the required form.  $\square$

The Hecke Eisenstein series of appropriate weight may be expressed as polynomials in the cubic theta functions by employing recursion formulas for cubic Eisenstein series derived by S. Cooper and others [7]. To avoid a conflict of notation, we adopt the conventions

$$G_{2k,1}(q) = \frac{L(1-2k, \mathbf{1})}{2} E_{2k,1}(q), \quad G_{2k,(\frac{\cdot}{3})}(q) = \frac{L(-2k, (\frac{\cdot}{3}))}{2} E_{2k+1,(\frac{\cdot}{3})}(q), \quad (2.13)$$

$$G_{2k}(q) = \frac{\zeta(1-2k)}{2} E_{2k}(q). \quad (2.14)$$

**Lemma 2.5.** *Let  $\chi$  and  $\mathbf{1}$  denote, respectively, the Jacobi symbol and the principal character modulo three. Then, for each integer  $n \geq 1$ ,*

$$G_{2n+2,\chi}(q) = -9(2n+1)(2n+2)G_{0,\chi}^2(q)G_{2n,\chi}(q)$$

$$- 2(2n+1)(2n+2) \sum_{j=1}^{n-1} \binom{2n}{2j} G_{2j,\chi}(q)G_{2n+2-2j}(q),$$

$$G_{2n+2,1}(q) = 18G_{0,\chi}(q)G_{2n,\chi}(q) + 6 \sum_{j=1}^{n-1} \binom{2n}{2j} G_{2j,\chi}(q)G_{2n-2j,\chi}(q).$$

By applying Lemma 2.5, the cubic theta function parameterizations for Eisenstein series of low weight [7, 14], as well as the familiar recursion formula for Eisenstein series on the full modular group [6, p. 30], we may generate each expansion displayed in the introduction and derive corresponding dissections for Eisenstein series of higher weight

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} \left( \frac{d}{3} \right) d^6 \right) q^n = \frac{(q; q)_{\infty}^{17}}{(q^3; q^3)_{\infty}^3} + 3^4 \cdot 50q(q; q)_{\infty}^5 (q^3; q^3)_{\infty}^9 + \frac{3^9 \cdot 7q^2 (q^3; q^3)_{\infty}^{21}}{(q; q)_{\infty}^7},$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} d^9 \right) q^n = \frac{(q; q)_{\infty}^{26}}{(q^3; q^3)_{\infty}^6} + 3^5 \cdot 23 \cdot 47q(q; q)_{\infty}^{14} (q^3; q^3)_{\infty}^6 \quad (2.15)$$

$$+ 3^9 \cdot 2237q^2(q; q)_{\infty}^2 (q^3; q^3)_{\infty}^{18} + \frac{3^{13} \cdot 11 \cdot 61q^3 (q^3; q^3)_{\infty}^{30}}{(q; q)_{\infty}^{10}},$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} \left( \frac{d}{3} \right) d^{10} \right) q^n = -\frac{3 \cdot 341(q; q)_{\infty}^{25}}{(q^3; q^3)_{\infty}^3} - 3^7 \cdot 4477q(q; q)_{\infty}^{13} (q^3; q^3)_{\infty}^9 \quad (2.16)$$

$$- 3^{10} \cdot 20317q^2(q; q)_{\infty} (q^3; q^3)_{\infty}^{21} - \frac{3^{15} \cdot 1847q^3 (q^3; q^3)_{\infty}^{33}}{(q; q)_{\infty}^{11}}$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} \left( \frac{d}{3} \right) d^{12} \right) q^n = \frac{(q; q)_{\infty}^{35}}{(q^3; q^3)_{\infty}^9} + 3^4 \cdot 207076q(q; q)_{\infty}^{23} (q^3; q^3)_{\infty}^3 \quad (2.17)$$

$$+ 3^9 \cdot 722810q^2(q; q)_{\infty}^{11} (q^3; q^3)_{\infty}^{15} + \frac{3^{13} \cdot 722812q^3 (q^3; q^3)_{\infty}^{27}}{(q; q)_{\infty}} + \frac{3^{18} \cdot 55601q^4 (q^3; q^3)_{\infty}^{39}}{(q; q)_{\infty}^{13}},$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+2} d^{13} \right) q^n = \frac{3 \cdot 2731(q; q)_{\infty}^{34}}{(q^3; q^3)_{\infty}^6} + 3^6 \cdot 1674872q(q; q)_{\infty}^{22} (q^3; q^3)_{\infty}^6 \quad (2.18)$$

$$+ 3^{10} \cdot 9766130q^2(q; q)_{\infty}^{10} (q^3; q^3)_{\infty}^{18} + \frac{3^{15} \cdot 2790064q^3 (q^3; q^3)_{\infty}^{30}}{(q; q)_{\infty}^2} + \frac{3^{19} \cdot 597871q^4 (q^3; q^3)_{\infty}^{42}}{(q; q)_{\infty}^{14}},$$

$$\sum_{n=0}^{\infty} \left( \sum_{d|3n+1} d^{15} \right) q^n = \frac{(q; q)_{\infty}^{44}}{(q^3; q^3)_{\infty}^{12}} + 3^4 \cdot 13 \cdot 1019729q(q; q)_{\infty}^{32} \quad (2.19)$$

$$+ 3^{10} \cdot 80982274q^2(q; q)_{\infty}^{20} (q^3; q^3)_{\infty}^{12} + 3^{14} \cdot 228972994q^3(q; q)_{\infty}^8 (q^3; q^3)_{\infty}^{24}$$

$$+ \frac{3^{18} \cdot 152647045q^4 (q^3; q^3)_{\infty}^{36}}{(q; q)_{\infty}^4} + \frac{3^{22} \cdot 28621321q^5 (q^3; q^3)_{\infty}^{48}}{(q; q)_{\infty}^{16}}.$$

To prove the congruences on line (2.1), apply Euler's theorem  $a^{\varphi(n)} \equiv 1 \pmod{n}$ ,  $\gcd(a, n) = 1$ , with  $n = 3^{s+1}$  to reduce the divisor sums to one of the identities of Theorem 2.2. The congruences from (2.2), follow from the fact that the sums from (2.2) reduce to those of (2.1) modulo  $3^{s+1}$ , since for each odd prime  $p$ , [11, Lemma 1]

$$n^{\frac{1}{2}\varphi(p^\lambda)} = \left( \frac{n}{p} \right) \pmod{p^\lambda}, \quad p \nmid n, \quad \lambda \in \mathbb{N}. \quad (2.20)$$

## 3. GENERALIZED CUBIC DISSECTIONS

We now consider cubic multisections of modular forms beyond the Eisenstein series studied in the previous section. Our purpose here is to study the dissection operators

$$\Omega_{3,k} \left( \sum_{n=0}^{\infty} \nu_n q^n \right) = \sum_{n=0}^{\infty} \nu_{3n+k} q^n, \quad \pi \left( \sum_{n=0}^{\infty} \nu_n q^n \right) = \sum_{n=0}^{\infty} \nu_n q^{n/3} \quad (3.1)$$

$k = 0, 1, 2$ , on subspaces of homogeneous polynomials in the cubic theta functions. We construct explicit matrix representations for these operators and discuss their spectral properties. Some of the integer eigenvalue and eigenvector pairs induce interesting congruences for the coefficients of corresponding eigenforms modulo powers of three.

**Theorem 3.1.** *Let*

$$\frac{3^5}{504}(E_6(q^3) - E_6(q)) = \sum_{n=0}^{\infty} u_n q^n, \quad 27q(q^3; q^3)_{\infty}^6(q; q)_{\infty}^6 = \sum_{n=0}^{\infty} v_n q^n, \quad (3.2)$$

then

$$u_{3^{\ell}n} \equiv 0 \pmod{3^{5\ell}}, \quad v_{3^{\ell}n} \equiv 0 \pmod{3^{2\ell}}. \quad (3.3)$$

To prove the last theorem, we first derive expansions for more general operators.

**Theorem 3.2.** *let  $\mathfrak{D}_n(x, y)$  denote the complex span of homogeneous polynomials in  $x$  and  $y$  of degree  $d$  and let  $B_d$  be the matrix whose  $(r, k)$ th entry equals*

$$\sum_{j=0}^k \sum_{\ell=0}^{3d-3j} \binom{k}{j} \binom{3d-3j}{\ell} \binom{3j}{r-\ell} 2^{\ell} (-1)^{3j+\ell-r}. \quad (3.4)$$

Suppose that  $f(q) \in \mathfrak{D}_d(a^3, c^3)$  with

$$f(q) = \sum_{n=0}^{\infty} v_n q^n = \sum_{k=0}^d \alpha_k a^{3(d-k)}(q) c^{3k}(q), \quad \alpha_n \in \mathbb{C}, \quad (3.5)$$

$$B_d(\alpha_0, \alpha_1, \dots, \alpha_d)^T = (\beta_0, \beta_1, \dots, \beta_d)^T. \quad (3.6)$$

If  $\mathbf{1}$  denotes the principal Dirichlet character modulo three, then, for  $m = 0, 1, 2$ ,

$$\Omega_{3,m}(f) = \sum_{n=0}^{\infty} k_{3n+m} q^n = q^{-m/3} \sum_{k=0}^{d-1(m)} \beta_{3k+m} a^{3d-3k-m}(q) c^{3k+m}(q). \quad (3.7)$$

The first few matrices  $B_d$  for  $d = 1, 2, 3$  from Theorem 3.2 are given explicitly by

$$\begin{pmatrix} 1 & 6 & 12 & 8 \\ 0 & 9 & 9 & 9 \end{pmatrix}^T, \quad \begin{pmatrix} 1 & 12 & 60 & 160 & 240 & 192 & 64 \\ 0 & 9 & 63 & 171 & 234 & 180 & 72 \\ 0 & 0 & 81 & 162 & 243 & 162 & 81 \end{pmatrix}^T, \quad (3.8)$$

$$\begin{pmatrix} 1 & 18 & 144 & 672 & 2016 & 4032 & 5376 & 4608 & 2304 & 512 \\ 0 & 9 & 117 & 657 & 2088 & 4140 & 5328 & 4464 & 2304 & 576 \\ 0 & 0 & 81 & 648 & 2187 & 4212 & 5265 & 4374 & 2268 & 648 \\ 0 & 0 & 0 & 729 & 2187 & 4374 & 5103 & 4374 & 2187 & 729 \end{pmatrix}^T. \quad (3.9)$$



*Proof of 3.2.* Suppose that  $f(q) \in \mathfrak{D}_n(a^3, c^3)$  has the expansions given by (3.5). Then, from the Borwein's identity (1.5) and the binomial theorem we deduce

$$f(q) = \sum_{k=0}^d \alpha_k a^{3(d-k)} (a^3 - b^3)^k = \sum_{k=0}^d \alpha_k \sum_{j=0}^k \binom{k}{j} (-1)^{3j} a^{3d-3j} b^{3j} \quad (3.10)$$

Applying Theorem (1.3) and the binomial theorem, we see that  $\pi(f) = f(q^{1/3})$  equals

$$\begin{aligned} & \sum_{k=0}^d \alpha_k \sum_{j=0}^k \binom{k}{j} (-1)^{3j} (a+2c)^{3d-3j} (a-c)^{3j} \\ &= \sum_{k=0}^d \sum_{j=0}^k \sum_{\ell=0}^{3d-3j} \sum_{i=0}^{3j} \binom{k}{j} \binom{3d-3j}{\ell} \binom{3j}{i} (-1)^{3j-i} 2^\ell a^{3d-\ell-i} c^{\ell+i} \alpha_k. \end{aligned} \quad (3.11)$$

Therefore, for  $0 \leq k \leq d$  and  $0 \leq r \leq 3d$ , the coefficient of  $\alpha_k a^{3d-r} c^r$  equals the expression on line (3.4). Since the contribution to the  $m$ -dissection  $\sum_{n=0}^{\infty} v_n q^{(3n+m)/3}$  of  $\pi(f) = f(q^{1/3})$  arises entirely from terms in (3.11) with  $r \equiv m \pmod{3}$  we have shown that the matrices defined by (3.4) generate the requisite expansions for  $\Omega_{3,m}(f)$ .  $\square$

From Theorem 3.2, we deduce that the operator  $\Omega_{3,0}$  over  $\mathfrak{D}_d(a^3, c^3)$  corresponds to a  $(d+1) \times (d+1)$  matrix whose entries come from rows  $1, 4, \dots, 3d+1$  of  $B_d$ . In particular, the respective matrix representations for  $\Omega_{3,0}$  on  $\mathfrak{D}_d(a^3, c^3)$ ,  $d = 1, 2, 3$  are

$$\begin{pmatrix} 1 & 0 \\ 8 & 9 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 \\ 160 & 171 & 162 \\ 64 & 72 & 81 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 & 0 \\ 672 & 657 & 648 & 729 \\ 5376 & 5328 & 5265 & 5103 \\ 512 & 576 & 648 & 729 \end{pmatrix}. \quad (3.12)$$

The eigenvectors  $x_i$  and eigenvalues  $\lambda_i$  for the matrix corresponding to  $\mathfrak{D}_2(a^3, c^3)$  are

$$x_i = (0 \ 9 \ 4)^T, \quad (0 \ 1 \ -1)^T, \quad (121 \ -152 \ 40)^T, \quad \lambda = 243, 9, 1, \quad i = 1, 2, 3.$$

We may apply the cubic theta function parameterizations for Eisenstein series from [7] to deduce that the eigenform for  $x_1$  equals the Fourier expansion at  $\tau = 0$  for the Hecke Eisenstein series corresponding to the principal character modulo three

$$9a^3(q)c^3(q) + 4c^6(q) = \sum_{n=1}^{\infty} \frac{n^5(q^n + q^{2n})}{1 - q^{3n}} = \frac{3^5}{504} (E_6(q^3) - E_6(q)), \quad (3.13)$$

while  $x_2$  and  $x_3$  correspond, respectively, to the eigenforms

$$121a^6(q) - 152a^3(q)c^3(q) + 40c^6(q) = 121E_{5,\chi_{3,1}}(q), \quad a^3(q)c^3(q) - c^6(q) = c^3(q)b^3(q).$$

This proves Theorem 3.1. From divisor sum expansions for (2.5), it follows that the Eisenstein series for primitive Dirichlet character  $\chi$  at the cusp  $\tau = \infty$ ,  $E_{3j,\chi}(q)$ , are eigenforms for  $\Omega_{3,0}$  on  $\mathfrak{D}_{3j}(a^3, c^3)$  with eigenvector 1, while the companion eigenforms

$$\sum_{n=1}^{\infty} \frac{n^{3j-1}(\chi(n)q^n + \chi(2)q^{2n})}{1 - q^{3n}} = \sum_{n=1}^{\infty} \left( \sum_{d|n} \chi(n/d) d^{3j-1} \right) q^n, \quad j \geq 1, \quad \chi(3j) = (-1)^j,$$

corresponding to the Eisenstein series at the cusp  $\tau = 0$ , have eigenvalue  $3^{3j-1}$ .

The matrices corresponding to  $\Omega_{3,0}$  on  $\mathfrak{D}_{3j}(a^3, c^3)$  have other interesting properties paralleling those of corresponding quintic operators from [9]. We conjecture that, up to sign, the determinants are powers of three. A proof of the conjecture may follow from an analysis of the spectral structure of  $\Omega_{3,0}$  by way of classical Hecke operators.

**Conjecture 3.3.** *Let  $C_d$  denote the matrix representation for  $\Omega_{3,0}$  on the vector space  $\mathfrak{D}_{3d}(a^3, c^3)$  of homogeneous polynomials of degree  $3d$  in  $a^3(q)$  and  $c^3(q)$  over  $\mathbb{C}$ . Then  $\det C_n = \pm 3^{w(n)}$  for some  $w(n) \in \mathbb{N}$ . For even indices, we have  $w(2n) = 4n(6n - 2)$ .*

Similar formulas may be obtained for the determinants of a class of matrix representations for  $\pi(f)$  corresponding to the embedding  $\pi(\mathfrak{D}_n(a, b))$  in  $\mathfrak{D}_n(a, c)$ . We give a precise construction for these trimidiation arrays in Theorem 3.4. Our proof of the theorem is similar to that of Theorem 3.2, and depends primarily on the Borwein's identity (1.5), the transformation formulas Theorem 1.3, and the binomial theorem.

**Theorem 3.4.** *Let  $f(q) = \sum_{n=0}^{\infty} \nu_n q^n$  be a homogeneous polynomial in  $a(q)$  and  $b(q)$ ,*

$$f(q) = \sum_{n=0}^d \alpha_n a^n(q) b^{(d-n)}(q), \quad (3.14)$$

where  $\alpha_n \in \mathbb{C}$ . Then there exists a  $(d+1) \times (d+1)$  matrix  $\mathcal{B}_d$  over  $\mathbb{Z}$  such that if

$$\mathcal{B}_d(\alpha_0, \alpha_1, \dots, \alpha_d)^T = (\beta_0, \beta_1, \dots, \beta_d)^T \quad (3.15)$$

then

$$\pi(f) = f(q^{1/3}) = \sum_{n=0}^d \beta_n a^n(q) c^{(d-n)}(q), \quad (3.16)$$

Moreover, the  $(r, n)$ th entry of  $\mathcal{B}_d$  equals

$$2^{d-n} (-1)^{n-r} \binom{n}{r} {}_2F_1 \left( n-d, -r; n-r+1; -\frac{1}{2} \right). \quad (3.17)$$

The hypergeometric function  ${}_2F_1$ , defined in [1, p. 61], arises from the the expansion

$$\sum_{k=0}^{d-n} 2^{(d-n)-k} (-1)^{n-(r-k)} \binom{d-n}{k} \binom{n}{r-k} \quad (3.18)$$

for the entries of  $\mathcal{B}_d$  as sums of binomial coefficients. The first few matrices  $\mathcal{B}_d$  defined in Theorem 3.4 are

$$\mathcal{B}_1 = \begin{pmatrix} 2 & -1 \\ 1 & 1 \end{pmatrix}, \quad \mathcal{B}_2 = \begin{pmatrix} 4 & -2 & 1 \\ 4 & 1 & -2 \\ 1 & 1 & 1 \end{pmatrix}, \quad \mathcal{B}_3 = \begin{pmatrix} 8 & -4 & 2 & -1 \\ 12 & 0 & -3 & 3 \\ 6 & 3 & 0 & -3 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

As with the matrices in Theorem 3.2, we observe that the determinants of the matrices from Theorem 3.4 are certain powers of three. This leads to a more general conjecture.

**Conjecture 3.5.** *Let  $\mathcal{B}_d$  be the trimidiation matrix defined in Theorem 3.4. Then*

$$\det \mathcal{B}_n = 3^{n(n+1)/2}.$$

## REFERENCES

- [1] G. E. Andrews, R. Askey, and R. Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.
- [2] Eiichi Bannai, Masao Koike, Akihiro Munemasa, and Jiro Sekiguchi. Some results on modular forms—subgroups of the modular group whose ring of modular forms is a polynomial ring. In *Groups and combinatorics—in memory of Michio Suzuki*, volume 32 of *Adv. Stud. Pure Math.*, pages 245–254. Math. Soc. Japan, Tokyo, 2001.
- [3] B. C. Berndt, S. Bhargava, and F. Garvan. Ramanujan’s theories of elliptic functions to alternative bases. *Trans. Amer. Math. Soc.*, 347(11):4163–4244, 1995.
- [4] J. M. Borwein and P. B. Borwein. A cubic counterpart of Jacobi’s identity and the AGM. *Trans. Amer. Math. Soc.*, 323(2):691–701, 1991.
- [5] J. M. Borwein, P. B. Borwein, and F. G. Garvan. Some cubic modular identities of Ramanujan. *Trans. Amer. Math. Soc.*, 343(1):35–47, 1994.
- [6] K. Chandrasekharan. *Elliptic functions*, volume 281 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1985.
- [7] S. Cooper. Cubic elliptic functions. *Ramanujan J.*, 11(3):355–397, 2006.
- [8] A. Granville and K. Ono. Defect zero  $p$ -blocks for finite simple groups. *Trans. Amer. Math. Soc.*, 348(1):331–347, 1996.
- [9] T. Huber. A theory of theta functions to the quintic base. *Submitted to the Journal of Number Theory, August 2012*.
- [10] C. G. J. Jacobi. *Fundamenta nova theoriae functionum ellipticarum*. Bornträger, Regiomonti, 1829.
- [11] D. B. Lahiri. Some congruences for the elementary divisor functions. *Amer. Math. Monthly*, 76:395–397, 1969.
- [12] S. Ramanujan. *Notebooks. Vols. 1, 2*. Tata Institute of Fundamental Research, Bombay, 1957.
- [13] A. Sebbar. Modular subgroups, forms, curves and surfaces. *Canad. Math. Bull.*, 45(2):294–308, 2002.
- [14] L.-C. Shen. On some cubic modular identities. *Proc. Amer. Math. Soc.*, 119(1):203–208, 1993.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS - PAN AMERICAN, 1201 WEST UNIVERSITY AVENUE, EDINBURG, TEXAS 78539, USA