

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

School of Mathematical and Statistical
Sciences Faculty Publications and
Presentations

College of Sciences

6-2-2021

Domes over Curves

Alexey Glazyrin

The University of Texas Rio Grande Valley

Igor Pak

Follow this and additional works at: https://scholarworks.utrgv.edu/mss_fac



Part of the [Mathematics Commons](#)

Recommended Citation

Alexey Glazyrin, Igor Pak, Domes over Curves, *International Mathematics Research Notices*, 2021,, rnab138, <https://doi.org/10.1093/imrn/rnab138>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in School of Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

DOMES OVER CURVES

ALEXEY GLAZYRIN* AND IGOR PAK[◊]

ABSTRACT. A closed piecewise linear curve is called *integral* if it is comprised of unit intervals. Kenyon's problem asks whether for every integral curve γ in \mathbb{R}^3 , there is a *dome over* γ , i.e. whether γ is a boundary of a polyhedral surface whose faces are equilateral triangles with unit edge lengths. First, we give an algebraic necessary condition when γ is a quadrilateral, thus giving a negative solution to Kenyon's problem in full generality. We then prove that domes exist over a dense set of integral curves. Finally, we give an explicit construction of domes over all regular n -gons.

1. INTRODUCTION

The study of polyhedra with regular polygonal faces is a classical subject going back to ancient times. It was revived periodically when new tools and ideas have developed, most recently in connection to algebraic tools in rigidity theory. In this paper we study one of most basic problems in the subject – polyhedral surfaces in \mathbb{R}^3 whose faces are congruent equilateral triangles. We prove both positive and negative results on the types of boundaries these surfaces can have, suggesting a rich theory extending far beyond the current state of the art.

Formally, let $\gamma \subset \mathbb{R}^3$ be a closed piecewise linear (PL-) curve.¹ We say that γ is *integral* if it is comprised of intervals of integer length. Now, let $S \subset \mathbb{R}^3$ be a PL-surface realized in \mathbb{R}^3 with the boundary $\partial S = \gamma$, and with all facets comprised of unit equilateral triangles. In this case we say that S is a *unit triangulation* or *dome over* γ , that γ is *spanned* by S , and that γ *can be domed*. By a PL-surface we mean a realization of a pure connected finite 2-dimensional simplicial complex, with no additional restriction of embedding or immersion.

Question 1.1 (Kenyon, see §6.2). *Is every integral closed curve $\gamma \subset \mathbb{R}^3$ spanned by a unit triangulation? In other words, can every such γ be domed?*

For example, the unit square and the (unit sided) regular pentagon can be domed by a regular pyramid with triangular faces. Of course, there is no such simple construction for a regular heptagon. Perhaps surprisingly, the answer to Kenyon's question is negative in general.

A 3-dimensional *unit rhombus* is a closed curve $\rho \subset \mathbb{R}^3$ with four edges of unit length. The unit rhombi form a 2-parameter family of space quadrilaterals $\rho(a, b)$ parametrized by the *diagonals* a and b , defined as distances between pairs of opposite vertices.

Theorem 1.2. *Let $\rho(a, b) \subset \mathbb{R}^3$ be a unit rhombus with diagonals $a, b > 0$. Suppose $\rho(a, b)$ can be domed. Then there is a nonzero polynomial $P \in \mathbb{Q}[x, y]$, such that $P(a^2, b^2) = 0$.*

In other words, for $a, b > 0$ algebraically independent over $\overline{\mathbb{Q}}$, the corresponding unit rhombus cannot be domed, giving a negative answer to Kenyon's question. In fact, our tools give further examples of unit rhombi which cannot be domed, such as $\rho(\frac{1}{\pi}, \frac{1}{\pi})$, see Corollary 4.9.

The following result is a positive counterpart to the theorem. We show that the set of integral curves spanned by a unit triangulation is everywhere dense within the set of all integral curves, with respect to the topology induced by Fréchet distance, see below.

*School of Mathematical & Statistical Sciences, University of Texas Rio Grande Valley, Brownsville, TX 78520. Email: alexey.glazyrin@utrgv.edu.

[◊]Department of Mathematics, UCLA, Los Angeles, CA, 90095. Email: pak@math.ucla.edu.

¹To avoid unnecessary technical difficulties, we do not consider curves with different vertices mapped to the same point although all our results remain true for them as well.

Let $\gamma, \gamma' \subset \mathbb{R}^3$ be two integral closed curves of equal length. We assume the vertices of γ, γ' are similarly labeled $[v_1 \dots v_n]$ and $[v'_1 \dots v'_n]$, giving parametrizations of the curves. The *Fréchet distance* $|\gamma, \gamma'|_F$ in this case is given by

$$|\gamma, \gamma'|_F = \max_{1 \leq i \leq n} |v_i v'_i|,$$

where $|v_i v'_i|$ is the Euclidean distance between v_i and v'_i .

Theorem 1.3. *For every integral curve $\gamma \subset \mathbb{R}^3$ and $\varepsilon > 0$, there is an integral curve $\gamma' \subset \mathbb{R}^3$ of equal length, such that $|\gamma, \gamma'|_F < \varepsilon$ and γ' can be domed.*

The theorem above does not give a concrete characterization of domed integral curves, and such a characterization seems difficult (see §5). We conclude with one interesting special case:

Theorem 1.4. *Every regular integral n -gon in the plane can be domed.*

This gives a new infinite class of *regular polygon surfaces*, comprised of one regular n -gon and many unit triangles. See Section 3 for the proof and some previously known special cases.

Outline of the paper. We begin with a technical proof of Theorem 1.3 in Section 2. Our proof is constructive and almost completely self-contained except for the Steinitz Lemma with Bergström constant, see §2.7. In Section 3, we follow with a (much shorter) constructive proof of Theorem 1.4, which is almost completely independent of the previous section, except for the earlier analysis of rhombi which can be domed, see §2.3.

In Section 4, we prove Theorem 1.2 by extending the results of Gaifullin brothers [13]. We assume that the reader is familiar with the *theory of places*, see e.g. [21, Ch. 1] and [26, §41.7], which played a key role in the solution of the *bellows conjecture*, see [7] (see also [26, §34]). Shifting gears once again, Section 5 is independent of the rest of the paper. Here we make a number of interrelated conjectures on the integral curves which can be domed, which we then relate to the *rigidity theory* and the *Euclidean Ramsey theory*. Final remarks are given in Section 6.

In the Appendix A, we include a negative solution of the question in [13] on the dimension of the flexes of doubly periodic surfaces. This counterexample arose upon careful inspection of our proof of Theorem 1.2, and is of independent interest (cf. [29]).

Notation. Let $|vw|$ denote the Euclidean distance between $v, w \in \mathbb{R}^3$. We use $[v_1 \dots v_n]$, $v_i \in \mathbb{R}^d$, to denote a closed polygonal curve $\gamma \subset \mathbb{R}^3$ with vertices v_1, \dots, v_n . We use parentheses notation (a_1, \dots, a_n) , $a_i > 0$, to denote the edge lengths of γ , i.e. $a_i = |v_i v_{i+1}|$, and $a_n = |v_n v_1|$. Denote by $|\gamma| = a_1 + \dots + a_n \in \mathbb{N}$ the length of the integral curve γ .

Throughout the paper all curves will be integral and closed PL-curves in \mathbb{R}^3 , unless stated otherwise. Similarly, all PL-surfaces S will have unit triangles, unless stated otherwise. They are realized in \mathbb{R}^3 by the vertex coordinates and such realizations have no additional extrinsic constraints (such as being embedding or immersion, cf. §6.2).

2. INTEGRAL CURVES WHICH CAN BE DOMED ARE DENSE

2.1. Understanding domes over curves. Before proceeding to the proofs of the theorems, let us give some basic ideas of domes over curves, and how they can be built. In Figure 1 two regular pentagonal pyramids give an example of domes over a regular pentagon. It's really the same dome up to a rigid motion. Similarly, the regular pentagonal biprism on the right can also be viewed as a dome over a disconnected integral curve comprised of two pentagons. We will not consider disconnected curves until §5.5.

Now, arrange the pyramid and the biprism as in the figure and notice that they can be attached to either of them to form yet another example of a dome over a pentagon. The idea of attaching triangulated PL-surfaces will be used repeatedly through the paper, both for explicit constructions and to disprove existence of other domes.




FIGURE 1. Three parts of an icosahedron as domes.

2.2. Mapping the proof of Theorem 1.3. The basic idea is to show that “generic” curves can be simplified to curves which can then be broken into pieces, each of which can have an explicit construction of a dome. The process of simplifying the curve and the construction are sufficiently robust to allow reversing the process. In essence we first target a curve, and construct a “generic” curve that is close enough.

Formally, denote by \mathcal{M}_n the space of all integral curves of length n in \mathbb{R}^3 , modulo rigid motions, which is compact in the Fréchet topology. Let $\mathcal{D}_n \subset \mathcal{M}_n$ denote the subset of integral curves which can be domed. The goal of this section is to prove Theorem 1.3, which states that \mathcal{D}_n is dense in \mathcal{M}_n .

The proof goes through several stages of simplification of integral curves, along the following route:


$$\text{integral curves} \longrightarrow \text{generic curves} \longrightarrow \text{near planar curves} \longrightarrow \text{compact near planar curves}.$$

Compact curves are curves which fit inside a ball of radius $3/2$ and they are much simpler to analyze by induction. At each stage, the simplification of curves is made by a sequence of certain local transformations. Starting the second arrow, these transformations are called *flips* and are obtained by attaching unit rhombi which can be domed. Making these reductions rigorous is somewhat technical and will occupy much of this section. The rhombi $\rho \in \mathcal{D}_4$ will play a special role, so we consider them first.

2.3. Dense rhombi. Throughout the paper, a unit closed curve of length 4 is called a *unit rhombus*, or just a rhombus. Each unit rhombus is determined by the *diagonals* a and b ; we denote such unit rhombus by $\rho(a, b)$. Observe that $a^2 + b^2 \leq 4$, with the equality achieved on plane rhombi.

Lemma 2.1. *Fix the diagonal a , and suppose $0 < a < 2$, $a \notin \overline{\mathbb{Q}}$. Then the set of values of $b \geq 0$ for which $\rho(a, b) \in \mathcal{D}_4$ is dense in $[0, \sqrt{4 - a^2}]$.*

Proof. Consider a planar isosceles trapezoid with side lengths $(1, 1, 1, a)$. Its circumradius is $\frac{1}{\sqrt{3-a}}$. Take a line through the circumcenter orthogonal to the plane containing the trapezoid and choose two points on this line with distance 1 from all vertices of the trapezoid (see Figure 2). Connecting these two points to the vertices, we obtain six unit triangles and a unit rhombus ρ_1 spanned by them. The diagonals of ρ_1 are a and $c_1 = 2\sqrt{\frac{2-a}{3-a}}$.

FIGURE 2. Construction of unit rhombus ρ_1 in the proof of Lemma 2.1.

We glue two copies of ρ_1 , with a common diagonal of length a , via two of their sides. Define $\rho_2 = \rho(a, c_2)$ to be a rhombus obtained as a boundary of the two glued copies of ρ_1 . Similarly, define $\rho_3 = \rho(a, c_3)$ by gluing three copies of ρ_1 , etc. Clearly, every rhombus ρ_m , $m \geq 1$, can be domed by surface with $6m$ unit triangles. We have:

$$c_m = \sqrt{4 - a^2} \cdot |\sin m\alpha|, \quad \text{where } \alpha := \arcsin \frac{2}{\sqrt{(2+a)(3-a)}}.$$

Thus, set $\{c_m, m \geq 1\}$ is dense in $[0, \sqrt{4-a^2}]$, for all $a \notin \pi\mathbb{Q}$. Finally, we have $a \notin \pi\mathbb{Q}$, since otherwise $\sin \alpha = \frac{2}{\sqrt{(2+a)(3-a)}} \in \overline{\mathbb{Q}}$, a contradiction with the assumption that $a \notin \overline{\mathbb{Q}}$. \square

Let \mathcal{D}'_4 denote the set of unit rhombi that can be domed using the construction from Lemma 2.1. The integer m in the proof will be called a *multiplier* throughout this section. We can now prove Theorem 1.3 for $|\gamma| = 4$. Let

$$(2.1) \quad \mathcal{X} := \left\{ x > 0 : \arcsin \frac{2}{\sqrt{(2+x)(3-x)}} \in \pi\mathbb{Q} \right\}.$$

Lemma 2.2. *Let $\rho = \rho(a, b) \in \mathbb{R}^3$ be a unit rhombus, and let $\varepsilon > 0$. Then there is a unit rhombus $\rho' = \rho(a', b') \in \mathcal{D}_4$, such that $|\rho, \rho'|_F < \varepsilon$. Moreover, if $a \notin \mathcal{X}$, one can take $a' = a$.*

Proof. The second part follows from the above proof of Lemma 2.1. For the first part, choose $a \notin \mathcal{X}$, so that $|a - a'| < \varepsilon$. Then apply the construction as above. \square

2.4. Reachable curves. Let us introduce some definitions and notation. Consider two integral curves $\gamma = [v_1 \dots v_k \dots v_n]$ and $\gamma' = [v_1 \dots v_{k-1} v'_k v_{k+1} \dots v_n]$, such that $[v_{k-1} v_k v_{k+1} v'_k] \in \mathcal{D}'_4$. In this case we say that γ and γ' are *k-flip connected*, or just *flip connected*; write $\gamma \rightarrow_k \gamma'$. Two integral curves $\gamma = [v_1 \dots v_n]$ and $\gamma' = [v'_1 \dots v'_n]$ are called *flip equivalent*, write $\gamma \sim \gamma'$, if

$$(2.2) \quad \gamma = \gamma_0 \rightarrow_{k_1} \gamma_1 \rightarrow_{k_2} \gamma_2 \rightarrow_{k_3} \dots \rightarrow_{k_N} \gamma_N = \gamma',$$

for some integer sequence $\mathbf{k} := (k_1, \dots, k_N)$, where $1 \leq k_i \leq N$ for all $i = 1, \dots, N$. See an example in Figure 3. Clearly, if $\gamma \sim \gamma'$ and $\gamma' \sim \gamma''$, then $\gamma \sim \gamma''$.




FIGURE 3. Sequence of two flips $\gamma_0 \rightarrow_2 \gamma_1 \rightarrow_3 \gamma_2$.

We say that an integral curve $\gamma \subset \mathbb{R}^3$ of length n is *reachable*, if for all $\varepsilon > 0$, there is an integral curve $\gamma' \subset \mathbb{R}^3$ of length n , such that $|\gamma, \gamma'|_F < \varepsilon$, and $\gamma' \in \mathcal{D}_n$. In this notation, Theorem 1.3 claims that all integral curves are reachable, while Lemma 2.2 proves this for curves of length 4.

Lemma 2.3. *Let $\gamma \sim \gamma'$ are flip equivalent integral curves in \mathbb{R}^3 . Suppose γ is reachable. Then so is γ' .*

In other words, the lemma says that if $\gamma \in \mathcal{M}_n$ is a limit point of \mathcal{D}_n , then so are all flip equivalent curves $\gamma' \sim \gamma$.

Proof. Since $\gamma \sim \gamma'$, there is a *flip sequence* \mathbf{k} as in (2.2), and a sequence of multipliers $\mathbf{m} = (m_1, \dots, m_N) \in \mathbb{Z}^m$ counting how many pairs of unit triangles are added at each flip. Use positive and negative integers m_i to denote clockwise or counterclockwise direction of the flip $\gamma_{i-1} \rightarrow \gamma_i$. A flip consists of attaching sequentially m sets of six triangles. Assume the first rhombus in a sequence spanned by six unit triangles is $[v_{k-1} v_k v_{k+1} v'_k]$. We say that the direction of the flip is positive if the determinant of the matrix $(v_{k-1} - v_k; v' - v_k; v_{k+1} - v_k)$ is positive. Otherwise, we say the direction of the flip is negative. Thus, pair (\mathbf{k}, \mathbf{m}) uniquely encodes the combinatorial structure of the flip equivalence. Let

$$(2.3) \quad \Phi_{\mathbf{k}, \mathbf{m}} : \mathcal{M}_n \rightarrow \mathcal{M}_n$$

be the map defining flip sequence as above. By construction, $\Phi_{\mathbf{k}, \mathbf{m}} : \mathcal{D}_n \rightarrow \mathcal{D}_n$, and $\Phi_{\mathbf{k}, \mathbf{m}}(\gamma) = \gamma'$.

Clearly, the map $\Phi_{\mathbf{k}, \mathbf{m}}$ is a composition of $|m_1| + \dots + |m_N|$ continuous maps, and thus also continuous on \mathcal{M}_n . Since γ is reachable, there is a sequence $\{\gamma^{(t)} \rightarrow \gamma, t \in \mathbb{N}\}$ of converging curves $\gamma^{(t)} \in \mathcal{D}_n$. Thus, we have another sequence of converging curves in \mathcal{D}_n :

$$\{\Phi_{\mathbf{k}, \mathbf{m}}(\gamma^{(t)}) \rightarrow \gamma', t \in \mathbb{N}\},$$

which shows that γ' is reachable. \square

2.5. Generic curves. Let $\gamma = [v_1 \dots v_n]$ be an integral curve in \mathbb{R}^3 . Following [7, 27] (see also [26, §34]), the distances $|v_i v_{i+2}|$ are called *small diagonals*. Here and below, we have vertex indices $1 \leq i \leq n$, and $(i+2)$ is taken modulo n .

We say that an integral curve $\gamma \subset \mathbb{R}^3$ is *generic*, if for all flip equivalent curves $\gamma' \sim \gamma$, where $\gamma' = [v'_1 \dots v'_n]$, we have small diagonals $|v'_i v'_{i+2}|$ are not in $\overline{\mathbb{Q}}$. Denote by $\mathcal{G}_n \subset \mathcal{M}_n$ the set of generic integral curves of length n . By definition, if a curve is generic then its flip equivalent curve is also generic.

Lemma 2.4. *Set \mathcal{G}_n is dense in \mathcal{M}_n .*

Proof. A neighborhood of a given curve γ in \mathcal{M}_n is a semi-algebraic set of dimension $2n$. Assume a concrete small diagonal $|v'_k v'_{k+2}|$ of a curve γ' is a fixed number from $\overline{\mathbb{Q}}$. The space of such curves in the neighborhood of γ is a semi-algebraic set of dimension at most $2n-1$. For a given sequence of flips, the space of curves produced from γ' with this sequence of flips is at most $(2n-1)$ -dimensional as well. Since both $\overline{\mathbb{Q}}$ and the set of flip sequences are countable, the set of non-generic curves in the neighborhood of γ is a union of countably many semi-algebraic sets of dimension at most $2n-1$. Therefore, there is a curve from \mathcal{G}_n in an arbitrary neighborhood of γ . \square

2.6. Planar curves. An integral curve $\gamma \in \mathcal{M}_n$ is called *planar* if it lies in a plane $H \subset \mathbb{R}^3$. Denote by $\mathcal{P}_n \subset \mathcal{M}_n$ the set of planar integral curves of length n .

Lemma 2.5. *For the flip equivalence class of every $\gamma \in \mathcal{G}_n$, there is a planar curve $\xi \in \mathcal{P}_n$ that is its limit point.*

In other words, for every $\varepsilon > 0$, and every generic integral curve $\gamma \in \mathcal{G}_n$, there is a generic integral curve $\gamma' \in \mathcal{G}_n$ and a planar integral curve $\xi \in \mathcal{P}_n$, such that $\gamma \sim \gamma'$ and $|\gamma', \xi|_F < \varepsilon$. Note that the curve ξ does not have to be generic itself, or be flip equivalent to γ .

Proof. The proof is based on the same idea of using flips to obtain a near-planar curve γ' . Let $\gamma = [v_1 \dots v_n] \in \mathcal{G}_n$, and let $\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a generic linear function, i.e. such that for any curve flip equivalent to γ , values φ are different on all its vertices. Generic linear functions exist because there are only countably many curves flip equivalent to a given one. Cyclically, for all k from 1 to n , make k -flips:

$$(2.4) \quad \gamma = \gamma_0 \rightarrow_1 \gamma_1 \rightarrow_2 \gamma_2 \rightarrow_3 \dots \rightarrow_n \gamma_n \rightarrow_1 \gamma_{n+1} \rightarrow_2 \gamma_{n+2} \rightarrow_3 \dots$$

Choose integers \mathbf{m} (see the proof of Lemma 2.3), as follows. Consider a flip

$$\gamma_{jn+k-1} = [\dots w_{k-1} w_k w_{k+1} \dots] \rightarrow_k \gamma_{jn+k} = [\dots w_{k-1} w'_k w_{k+1} \dots].$$

By the proof of Lemma 2.1, we can always choose a multiplier m_{jn+k} so that

$$(2.5) \quad \frac{2}{3}\alpha + \frac{1}{3}\beta < \varphi(w'_k) < \frac{1}{3}\alpha + \frac{2}{3}\beta,$$

where

$$\alpha := \min\{\varphi(w_{k-1}), \varphi(w_{k+1})\}, \quad \beta := \max\{\varphi(w_{k-1}), \varphi(w_{k+1})\}.$$

Note that we have $\alpha \neq \beta$, since φ is generic, so there is always room to make such flip possible.

Using (2.5), it is easy to see that there is a limit $(\varphi(w_1), \dots, \varphi(w_n)) \rightarrow (h, \dots, h)$, for some $h \in \mathbb{R}$. Here the limit is when $N \rightarrow \infty$, where N is the number of flips in (2.4). Indeed, note that $\max_k \varphi(w_k)$ is non-increasing and thus converges to some $M < \infty$. Similarly, note that $\min_k \varphi(w_k)$ is non-decreasing and thus converges to some $\mu \leq M$. If $\mu = M$, then we can take $h = M = \mu$. Therefore, it remains to show that $\mu < M$ is impossible.

Since $\max_k \varphi(w_k) \rightarrow M$, there is a moment when all values of $\varphi(w_k)$ are smaller than $M + \delta$ for a given δ . On the other hand there is ℓ such that $\varphi(w_\ell) \neq \mu$. Let us consider the next n flips starting from the vertex $\ell + 1$. By the construction of the flip sequence, we have $\varphi(w'_{\ell+1}) < \frac{1}{3}\mu + \frac{2}{3}(M + \delta)$, then $\varphi(w'_{\ell+2}) < \frac{1}{9}\mu + \frac{8}{9}(M + \delta)$, etc. We conclude that all values of φ after n flips are no greater than $\frac{1}{3^n}\mu + \frac{3^n-1}{3^n}(M + \delta)$. For $\mu < M$, we can choose $\delta > 0$ such that this value is smaller than M . This contradicts the assumption $\max_k \varphi(w_k) \rightarrow M$, and implies $\mu = M$.

The limit curve ξ is integral and lies in the plane $H := \{x \in \mathbb{R}^3 : \varphi(x) = h\}$. Therefore, for $N = N(\varepsilon)$ large enough, we obtain a curve $\gamma' := \gamma_N$, such that $\gamma' \sim \gamma$, and $|\gamma', \xi|_F < \varepsilon$. \square

2.7. Packing curves. Let $u_1, \dots, u_n \in \mathbb{R}^d$ be unit vectors which satisfy $u_1 + \dots + u_n = 0$. The *Steinitz Lemma* famously states, see e.g. [3], that there is always a permutation $\sigma \in S_n$, s.t.

$$(2.6) \quad |u_{\sigma(1)} + \dots + u_{\sigma(k)}| \leq B_d \quad \text{for all } 1 \leq k \leq n,$$

where $B_d \leq 2d$ is a universal constant which depends only on the dimension d . Bergström [4] found the optimal value $B_2 = \sqrt{5/4}$, see also §6.5.

Motivated by the Steinitz Lemma, we define a similar notion for integral curves. Let $\gamma = [v_1 \dots v_n] \in \mathcal{M}_n$ be an integral curve in \mathbb{R}^3 . We say that γ is *B-packing*, if $|v_i v_i| \leq B$ for all $1 \leq i \leq n$.

Lemma 2.6. *Every generic integral curve $\gamma \in \mathcal{G}_n$ is flip equivalent to a generic integral curve $\gamma'' \in \mathcal{G}_n$ that is 3/2-packing.*

Here the constant $B = 3/2$ is chosen somewhat arbitrary. In fact, any constant $\sqrt{5/4} < B < \sqrt{3}$ will satisfy the lemma and suffice for our purposes.

Proof. Fix $\varepsilon > 0$ and $\gamma \in \mathcal{G}_n$. Let $\gamma' \sim \gamma$ and $\xi = [w_1 \dots w_n] \in \mathcal{P}_n$ be as in the proof of Lemma 2.5, so $|\gamma', \xi| < \varepsilon$. Define $u_i = \overrightarrow{w_i w_{i+1}}$, $1 \leq i < n$, and $u_n = \overrightarrow{w_n w_1}$. Clearly, u_i are unit vectors which satisfy $u_1 + \dots + u_n = 0$. By the Steinitz Lemma, there is a permutation $\sigma \in S_n$, s.t. (2.6) holds. Consider a reduced factorization of σ into adjacent transpositions $(i, i+1) \in S_n$:

$$\sigma = (k_\ell, k_\ell + 1) \cdots (k_1, k_1 + 1),$$

where $1 \leq k_1, \dots, k_\ell \leq n-1$, and $\ell = \text{inv}(\sigma)$ is the number of inversions in σ , see e.g. [32]. This reduced factorization may not be unique, of course. Denote a partial permutation $\sigma_j = (k_j, k_{j+1}) \cdots (k_1, k_1 + 1)$, and the corresponding planar curve $\xi_j = [w_{\sigma_j(1)} \dots w_{\sigma_j(n)}]$.

Define a sequence of flips as in (2.2), according to this factorization:

$$(2.7) \quad \gamma' = \gamma_0 \xrightarrow{k_1} \gamma_1 \xrightarrow{k_2} \gamma_2 \xrightarrow{k_3} \dots \xrightarrow{k_\ell} \gamma_\ell = \gamma''.$$

Since γ_j are generic, by Lemma 2.1 and induction, we can always choose the multipliers m_j so that $|\gamma_j, \xi_j| < \varepsilon$, $1 \leq j \leq \ell$.

Now let $\varepsilon \rightarrow 0$. As in the proof of Lemma 2.3, by continuity of $\Phi_{\mathbf{k}, \mathbf{m}}$ in (2.3), we have the limit planar curve $\gamma'' \rightarrow \varrho := [y_1 \dots y_n] \in \mathcal{P}_n$. By induction on the length ℓ of the factorization, we have: $\overrightarrow{y_i y_{i+1}} = u_{\sigma(i)}$, $1 \leq i < n$, and $\overrightarrow{y_n y_1} = u_{\sigma(n)}$. By the Steinitz Lemma with Bergström constant $B_2 = \sqrt{5/4}$, we conclude that for sufficiently small $\varepsilon > 0$, the integral curve γ'' is $(B_2 + \delta)$ -packing, for all $\delta > 0$. Taking $\delta < (3/2 - B_2)$, we obtain the result. \square

Remark 2.7. In notation of the proof above, in a special case of a convex centrally symmetric n -gon $\xi \in \mathcal{P}_n$, $n = 2k$, the sequence of unit vectors is $u_1, \dots, u_k, -u_1, \dots, -u_k$. Take the permutation which gives the order $u_1, \dots, u_k, -u_k, \dots, -u_1$; the corresponding limit curve $\varrho \in \mathcal{P}_n$ is then degenerate. For every reduced factorization as in the proof, the pattern of rhombi used in the flip sequence then defines a *zonotopal tilings*, see e.g. [26, Exc. 14.25].

2.8. Proof of Theorem 1.3. We note that if the statement of the theorem is true then it holds with the additional constraint that $v_1 = v'_1$, $v_2 = v'_2$, and there is a plane through v'_1, v'_2, v'_3 containing v_1, v_2, v_3 .

We prove the result by induction. First, closed integral curve of length 3 is a unit triangle, so Theorem 1.3 is trivially true in this case. The case of length 4 is resolved in Lemma 2.2. Note that by Lemma 2.4 it suffices to prove the theorem only for generic curves $\gamma = [v_1 \dots v_n] \in \mathcal{G}_n$.

Let $n = 5$, and let $\gamma \in \mathcal{G}_5$ be a generic integral pentagon. By Lemma 2.6, there is an almost planar curve $\gamma' = [w_1, \dots, w_5] \in \mathcal{G}_5$, such that $\gamma' \sim \gamma$, and $|w_i w_i| \leq 3/2$ for all $1 \leq i \leq 5$. If the circumradius of the triangle $w_1 w_3 w_4$ is less than 1, then there is a point $z \in \mathbb{R}^3$, s.t. $|w_1 z| = |w_3 z| = |w_4 z| = 1$, see Figure 4 (left). Otherwise, we make a flip for the vertex w_1 using Lemma 2.1 to construct a generic curve $\gamma'' = [w'_1, w_2, \dots, w_5]$ such that the circumradius of the triangle $w'_1 w_3 w_4$ is less than 1 and take a point z , s.t. $|w'_1 z| = |w_3 z| = |w_4 z| = 1$. Without loss of generality, we consider the first case only.

Apply now Lemma 2.2 to rhombi $\rho_1 = [w_1 w_2 w_3 z]$ and $\rho_2 = [w_1 w_5 w_4 z]$, to obtain rhombi $\rho'_1 = [w_1 w'_2 w_3 z] \in \mathcal{D}_4$ and $\rho'_2 = [w_1 w'_5 w_4 z] \in \mathcal{D}_4$, which satisfy $|w_2 w'_2|, |w_5 w'_5| < \varepsilon$. Attach unit triangle $[w_3 w_4 z]$ to rhombi ρ'_1 and ρ'_2 . This gives the desired pentagon $\eta = [w_1 w'_2 w_3 w_4 w'_5] \in \mathcal{D}_5$, s.t. $|\gamma', \eta|_F < \varepsilon$. Thus, γ' is reachable. By Lemma 2.3, then so is γ , as desired.

We also note that for each $\varepsilon > 0$ there is $\delta > 0$ such that for any angle $\psi \in [\angle w_2 w_1 w_5 - \delta, \angle w_2 w_1 w_5 + \delta]$, there is $\gamma^* = [w_1^* w_2^* w_3^* w_4^* w_5^*] \in \mathcal{D}_5$ satisfying $|\gamma', \gamma^*|_F < \varepsilon$ and $\angle w_2^* w_1^* w_5^* = \psi$. Indeed, we can just perturb the construction of η . Fixing w_3, w_4 , and z and multipliers of the rhombi ρ_1' and ρ_2' we can move w_1^* in the neighborhood of w_1 . Then the points w_2^* and w_5^* defined by w_1^* change continuously with respect to the position of w_1^* . This means $\angle w_2^* w_1^* w_5^*$ changes continuously as well. It is easy to check that values both smaller and larger than the initial $\angle w_2 w_1 w_5$ are possible so there is an interval of angles that are possible.




FIGURE 4. Base of induction $n = 5$, and step of induction for $n = 7$.

For $n \geq 6$, we employ a similar argument. Let $\gamma \in \mathcal{G}_n$ be a generic integral curve as above. By Lemma 2.6, there is an almost planar $\gamma' = [w_1 \dots w_n] \in \mathcal{G}_n$, such that $\gamma' \sim \gamma$, and $|w_1 w_i| \leq 3/2$ for all $1 \leq i \leq n$. We choose a point $z \in \mathbb{R}^3$ such that $|w_1 z| = |w_4 z| = 1$ and the circumradius of the triangle $w_2 w_3 z$ is less than 1. Then for the curve $[w_1 w_2 w_3 w_4 z]$ we can use the construction from above. We conclude that in a neighborhood of $[w_1 w_2 w_3 w_4 z]$ there are curves $\gamma^* \in \mathcal{D}_5$ whose angles $w_1^* z^* w_4^*$ cover $[\angle w_1 z w_4 - \delta, \angle w_1 z w_4 + \delta]$.

Now take $\varepsilon' > 0$, such that $|w_1 w_1'|, |z z'|, |w_4 w_4'| < \varepsilon'$ implies $\angle w_1' z' w_4' \in [\angle w_1 z w_4 - \delta, \angle w_1 z w_4 + \delta]$. The integral curve $\gamma' = [w_1 z w_4 \dots w_n]$ has length $n - 1$ so, by the induction hypothesis, there is $\gamma'' = [w_1' z' w_4' \dots w_n'] \in \mathcal{D}_{n-1}$ such that $|\gamma'', \gamma'|_F < \varepsilon'$. Since $\angle w_1' z' w_4' \in [\angle w_1 z w_4 - \delta, \angle w_1 z w_4 + \delta]$, there is a curve of length five that can be domed in the neighborhood of $[w_1 w_2 w_3 w_4 z]$ with the same angle. Gluing the domes of these two curves, we obtain a dome of the curve in the neighborhood of γ . This completes the proof of the reduction and finishes the proof of the theorem. \square

3. REGULAR POLYGONS

3.1. Classical domes. Denote by $Q_n \subset \mathbb{R}^2$ the regular n -gon with unit sides in the xy -plane with the center at the origin O . From the introduction, there is a trivial dome over Q_3 and Q_6 , and domes over Q_4, Q_5 are given by regular pyramids. Less obviously, a tiling of Q_{12} given in Figure 5 (left), gives a natural dome over Q_{12} , when square pyramids are added.



FIGURE 5. Left: Tiling giving a dome over Q_{12} . Right: Pentagonal cupola giving a dome over Q_{10} .

Similarly, recall that the regular octagon Q_8 and decagon Q_{10} are spanned by the surfaces of Johnson solids *square cupola* and *pentagonal cupola*, respectively, see Figure 5 (right) and [17] for details.² In fact, both are cuts of the Archimedean solids, see e.g. [9, p. 88]. The faces of both surfaces are regular triangles, squares or pentagons. Adding a pyramid to each face we obtain domes over Q_8 and Q_{10} .

²The image on the right is available from the [Wikimedia Commons](#), and is free to use with attribution.

3.2. Proof of Theorem 1.4. We follow notation in the proof of Lemma 2.1, and employ the symmetry of Q_n at every step.

First, attach a unit triangle to each side of Q_n at angle $\theta > 0$ to the plane. In the construction below, we make assumptions on what are distances a_i and α defined below. These will prohibit countably many values of θ in a manner similar to the proof of Theorem 1.3. We present the construction first for the purposes of exposition.

Let the angle $\theta > 0$ be very small, to be chosen at a later point. Denote by a_1 the distances between vertices of adjacent unit triangles, see Figure 6 below. Assume that $a_1 \notin \overline{\mathbb{Q}}$. Note that $a_1 > 0$ is well defined for $n \geq 7$.

Moving along the boundaries of triangles attached to Q_n , attach to their unit edges n rhombi $R_1 = \rho_{m_1}(a_1, *)$. To simplify the notation, we use $(*)$ for the second diagonal, since it is completely determined by the multiplier m_1 and a , see the proof of Lemma 2.1. Take m_1 large enough and chosen so that R_1 is nearly planar, oriented towards the z axis, and at an angle $\theta_1 > \theta$ with the plane. Such m_1 exists by Lemma 2.1 if we assume further that $a_1 \notin \overline{\mathbb{Q}}$.




FIGURE 6. Example of rhombi R_1 , R_2 , and distances a_1 , a_2 , a_3 .

Next, moving along the boundary, denote by a_2 the distances between vertices of adjacent rhombi R_1 , and observe that $a_2 \notin \overline{\mathbb{Q}}$. Now attach to the adjacent unit edges rhombi $R_2 = \rho_{m_2}(a_2, *)$, where the multiplier m_2 is large enough and chosen so that R_2 is nearly planar, at an angle $\theta_2 > \theta_1$, see Figure 7. Again, such m_2 exists by Lemma 2.1, if we assume further that $a_2 \notin \overline{\mathbb{Q}}$.

Repeat this procedure for k iterations, until the distance β to the vertical z -axis from new rhombi vertices satisfies $\beta < \sqrt{1 - \alpha^2/4}$. Here $\alpha := a_{k+1}$ denotes the distance between closest new vertices of the adjacent rhombi $R_k = \rho_{m_k}(a_k, *)$, and we assume that $\alpha \notin \overline{\mathbb{Q}}$. The above bound on β corresponds to having the projection of the nearly planar rhombus R_{k+1} cover the origin O , see Figure 7 (center).

At this stage, attach to the adjacent unit edges new unit rhombi $R = \rho_M(\alpha, *)$ in such a way that the new vertices are at distance $\delta > 0$ from the z -axis, see Figure 7 (center). By Lemma 2.1, distance $\delta > 0$ can be made as small as necessary.

Now, the construction above is uniquely determined by the angle $\theta > 0$ and the integer sequence

$$\mathbf{m} := (m_1, m_2, \dots, m_k, M).$$

Since the number of vectors \mathbf{m} is countable, the assumptions $a_i, \alpha \notin \overline{\mathbb{Q}}$ over all \mathbf{m} represent countably many inequalities on θ , so for some $\theta > 0$ the above construction is well defined.

The resulting (partial) surface S is continuously deformed with θ for every fixed \mathbf{m} (cf. the proof of Lemma 2.3). Indeed, this follows by induction: first observe that a_1 is continuously changing with θ , then so does a_2 , etc., until we conclude with α ; details are straightforward.

Now, continuously changing $\theta > 0$ and using the symmetry, we can place all the remaining free rhombi vertices onto the z axis. This completes the construction of a dome over Q_n for all $n \geq 7$, and the smaller cases $3 \leq n \leq 6$ are discussed above.

Finally, for a regular n -gon rQ_n , replace unit triangles Q_3 with their scaled version rQ_3 and proceed as above. Now triangulate every copy rQ_3 with r^2 unit triangles Q_3 , completing the construction of a dome over rQ_n . \square




FIGURE 7. Left: Nearly planar tiling of a portion of Q_n with rhombi. Middle: Vertical slice. Right: Example of Q_{12} with triangles and nearly-flat rhombi R_1 .

Remark 3.1. Also, one can ask if a version of the *arm lemma* (see e.g. [26, §22.2]), holds in this case. We believe this to be true for every fixed \mathbf{m} , on a sufficiently small interval $\theta \in (\theta_0 - \epsilon, \theta_0 + \epsilon)$, but this result is not necessary for the argument in the proof.

4. THE ALGEBRA OF SQUARED DIAGONALS

4.1. Contractible domes. As a warm-up to the proof of Theorem 1.2, we first present a short argument for the case when the spanning surface S is homeomorphic to a disc.

Proposition 4.1. *Let $\gamma \subset \mathbb{R}^3$ be a unit rhombus $\gamma = \rho(s, t)$, with diagonal lengths s and t . Suppose γ can be domed by a surface homeomorphic to a disc. Then there exists a polynomial $P \in \mathbb{Q}[x, y]$, such that $P(s^2, t^2) = 0$.*


For the proof of the proposition, we need to consider doubly periodic surfaces homeomorphic to the plane. Let K be a simplicial connected pure 2-dimensional complex with a free action of the group $G = \mathbb{Z} \oplus \mathbb{Z}$ with generators a and b . Assume that G acts as a linear bijection on each simplex of K , and that the number of orbits of triangles under the action of G is finite. Consider a mapping $\theta : K \rightarrow \mathbb{R}^3$, linear on each simplex of K , and equivariant with respect to the action of $\mathbb{Z} \oplus \mathbb{Z}$, such that a and b act by translations with vectors α and β , respectively. Then the pair (K, θ) is called a *doubly periodic triangular surface*. Sometimes, with a slight abuse of notation, we call the surface K as well.

Now, let us construct a doubly periodic surface comprised of unit triangles for every unit triangulation of a unit rhombus.

Lemma 4.2. *For a unit rhombus $\gamma = \rho(s, t)$ with diagonals s and t that can be domed, there is a doubly periodic surface of unit triangles with two orthogonal periodicity vectors of length s and t , respectively. Moreover, there is such a surface homeomorphic to the plane if the unit triangulation of the rhombus spans a surface homeomorphic to a disc.*

Proof. First we construct a doubly periodic surface whose cells are either parallel translates of γ or $-\gamma$ (see Figure 8). This surface is combinatorially equivalent to a tiling of the plane with unit squares. A chessboard coloring of such a tiling makes white squares correspond to parallel translates of γ and black squares correspond to parallel translates of $-\gamma$. The periodicity vectors of this surface are the vectors of the diagonals of γ .

Attaching a spanning unit triangulation to each translate of γ and $-\gamma$ we obtain a doubly periodic polyhedral surface comprised of unit triangles with required periodicity vectors. Clearly, if a spanning triangulation of γ is homeomorphic to a disk, the resulting doubly periodic surface is homeomorphic to the plane. \square

FIGURE 8. Doubly periodic surface from translates of γ and $-\gamma$.

At this point we consider only doubly periodic triangular surfaces (K, θ) with unit triangular faces. For a fixed complex K , let $\mathcal{G}(K)$ be the set of all possible Gram matrices formed by vectors α and β for all doubly periodic triangular surfaces (K, θ) . For a Gram matrix $G \in \mathcal{G}(K)$, we denote its entries by g_{11} , $g_{12} = g_{21}$, and g_{22} .

Theorem 4.3 (Gaifullin–Gaifullin [13]). *Let K be a simplicial pure 2-dimensional complex homeomorphic to \mathbb{R}^2 with a free action of the group $\mathbb{Z} \oplus \mathbb{Z}$. Then there is a one-dimensional real affine algebraic subvariety of \mathbb{R}^3 containing $\mathcal{G}(K)$.*

In particular, the entries of each Gram matrix G from $\mathcal{G}(K)$ satisfy a one-dimensional system of two non-trivial polynomial equations with integer coefficients:

$$\begin{cases} p(g_{11}, g_{12}, g_{22}) = 0 \\ q(g_{11}, g_{12}, g_{22}) = 0. \end{cases}$$

Remark 4.4. In fact, the result in [13] is more general, as the authors consider all polygonal doubly periodic surfaces homeomorphic to the plane with arbitrary sets of side lengths. In this setting, the coefficients of polynomials p and q are obtained from the ideal generated by squares of all side lengths of polygons in the polygonal surface

Proof of Proposition 4.1. If a unit rhombus with diagonals s and t can be spanned by unit triangles then, by Lemma 4.2, there is a doubly periodic triangular surface with orthogonal periodicity vectors of length s and t . The entries of the Gram matrix of periodicity vectors are $g_{11} = s^2$, $g_{12} = 0$, $g_{22} = t^2$. By Theorem 4.3, there are two polynomials p, q with integer coefficients vanishing on the entries of the Gram matrix. Thus, at least one of the equations $p(s^2, 0, t^2) = 0$ and $q(s^2, 0, t^2) = 0$ is non-trivial, and can be used for the polynomial P . \square

4.2. Theory of places. There is no result generalizing Theorem 4.3 for doubly periodic surfaces of non-trivial topology and, moreover, we will show in Theorem A.2 that such a generalization is not true. However, for our purposes, we do not need two polynomials p and q as in Theorem 4.3. It is sufficient to find at least one polynomial that is non-trivial whenever $g_{12} = 0$. The machinery developed in [13] is based on the proof of the bellows conjecture for orientable 2-dimensional surfaces [7], and is also the basis for our approach. We use places of fields as the main algebraic instrument of the proof.

Let F be a field and \widehat{F} be F extended by ∞ , i.e. $\widehat{F} = F \cup \{\infty\}$ with arithmetic operations extended to \widehat{F} by

$$\begin{aligned} a \pm \infty &= \infty \quad \text{and} \quad \frac{a}{\infty} = 0, \quad \text{for all } a \in F, \\ a \cdot \infty &= \frac{a}{0} = \infty \quad \text{for all } a \in \widehat{F} \setminus \{0\}. \end{aligned}$$

The expressions

$$\frac{0}{0}, \frac{\infty}{\infty}, 0 \cdot \infty \quad \text{and} \quad \infty \pm \infty \quad \text{are not defined.}$$

Let L be a field. A map $\phi : L \rightarrow \widehat{F}$ is called a *place* if $\phi(1) = 1$ and

$$\phi(a \pm b) = \phi(a) \pm \phi(b), \quad \phi(a \cdot b) = \phi(a) \cdot \phi(b) \quad \text{for all } a, b \in L,$$

whenever the right-hand side expressions are defined.

As a direct consequence of the definition, we have $\phi(0) = 0$ for all places, and $\phi(x) = \infty$ for $x \neq 0$ if and only if $\phi(-x) = \infty$ and if and only if $\phi(1/x) = 0$. It is also clear that whenever $\text{char } F = 0$, we must also have $\text{char } L = 0$. Similarly, we have $\phi(kx) = \infty$ for a non-zero $k \in \mathbb{Z}$, if and only if $\phi(x) = \infty$. We will use the following basic fact on extensions of places.

Lemma 4.5 (see e.g. [21, Ch. 1, Thm 1]). *Let L be a field containing a ring R . Let ϕ be a homomorphism from R to an algebraically closed field Ω , and suppose $\phi(1) = 1$. Then ϕ can be extended to a place $L \rightarrow \Omega \cup \{\infty\}$.*

4.3. General domes. For a doubly periodic triangular surface (K, θ) , let α and β be the periodicity vectors of the surface and Λ be the lattice generated by α and β . The number of orbits under the action of $G = \mathbb{Z} \oplus \mathbb{Z}$ is finite, so we choose a representative of each orbit. Let $(x_1, y_1, z_1), \dots, (x_N, y_N, z_N)$ be their coordinates in \mathbb{R}^3 , and let $(x_\alpha, y_\alpha, z_\alpha), (x_\beta, y_\beta, z_\beta)$ be the coordinates of the periodicity vectors. Define field \mathbb{L} as follows:

$$\mathbb{L} := \mathbb{Q}(x_1, y_1, z_1, \dots, x_N, y_N, z_N, x_\alpha, y_\alpha, z_\alpha, x_\beta, y_\beta, z_\beta).$$

Note that \mathbb{L} does not depend on the choice of representatives of orbits and the choice of the basis for the lattice Λ .

When vertices a, b on the surface (K, θ) form an edge, denote by ℓ_{ab} the squared distance between them:

$$\ell_{ab} := (x_a - x_b)^2 + (y_a - y_b)^2 + (z_a - z_b)^2.$$

Clearly, $\ell_{ab} \in \mathbb{L}$. For each surface the set of all possible ℓ_{ab} is finite. Let R be the \mathbb{Q} -subalgebra of \mathbb{L} generated by all ℓ_{ab} of the surface.

All vectors in Λ can be written as integer linear combinations of the periodicity vectors, $\lambda = k\alpha + m\beta$. In case k and m are relatively prime, the vector λ is called *primitive*. Denote by Λ^* the set of primitive vectors $\lambda \in \Lambda$. By (λ_1, λ_2) we mean the standard inner product of vectors λ_1 and λ_2 .

As the first step in the proof of Theorem 1.2, we prove the lemma on finite elements of places.

Lemma 4.6 (Main lemma). *For a doubly periodic triangular surface (K, θ) obtained by the construction in Lemma 4.2, let $\phi : \mathbb{L} \rightarrow F \cup \{\infty\}$ be a place that is finite on all ℓ_{ab} defined by the surface and let $\text{char } F = 0$. Then there is a vector $\lambda \in \Lambda$, $\lambda \neq 0$, such that ϕ is finite on (λ, λ) .*

For the proof, we use the following technical result of Connelly, Sabitov, and Walz [7, Lemma 4], see also [26, §34.3].

Theorem 4.7 (Connelly–Sabitov–Walz). *Let u be a vertex of a triangular surface in \mathbb{R}^3 and v_1, \dots, v_d , $d \geq 4$, be adjacent to it in this cyclic order, denote also $v_{d+1} = v_1$ and $v_{d+2} = v_2$. Let ϕ be a place that is defined on $\mathbb{Q}(x_u, y_u, z_u, x_{v_1}, y_{v_1}, z_{v_1}, \dots, x_{v_d}, y_{v_d}, z_{v_d})$ and is finite on all $\ell_{uv_i}, \ell_{v_i v_{i+1}}$, $1 \leq i \leq d$. Then ϕ is finite on at least one of the squared diagonal lengths $\ell_{v_i v_{i+2}}$, $1 \leq i \leq d$.*

4.4. Proof of the Main Lemma 4.6. The statement of the lemma is true if one of the edges of the surface forms a vector from Λ . We define the *complexity* as a partial ordering of doubly periodic triangular surfaces with the same periodicity lattice Λ . Surfaces with edges from Λ are called the *least complex* (an example is given in Figure 8). For surfaces without edges from Λ , the ordering is defined as follows.

A surface K_1 is said to be *less complex* than K_2 , if the Euler characteristic of K_1/Λ is greater than the Euler characteristic of K_2/Λ . The surface K_1 is less complex than K_2 if the Euler characteristics of K_1/Λ and K_2/Λ are the same, and K_1/Λ has fewer vertices than K_2/Λ . The surface K_1 is less complex than K_2 if K_1/Λ and K_2/Λ have the same Euler characteristic and the same number of vertices, but the smallest vertex degree of K_1 is less than the smallest vertex degree of K_2 . The proof will proceed by induction on complexity.

First case. Suppose the surface contains the edges ab, bc, ca , but does not contain a triangle abc . The closed curve formed by the edges ab, bc and ca , divides its neighborhood into two components. Then we define the surgery along $[abc]$ by removing vertices a, b, c , edges ab, bc, ca , and adding two copies of the triangle abc , which we call $a'b'c'$ and $a''b''c''$. We do this in such a way that $a'b'c'$ and $a''b''c''$ retain the incidences of a, b, c in the first and the second component of the neighborhood, respectively.

We make this *surgery* for all periodic images of abc under the action of Λ . If this surgery keeps the surface connected, then it increases the Euler characteristic of K/Λ . If the surgery splits the surface into two new surfaces then the Euler characteristic for each of them is not smaller than the initial Euler characteristic, for both of them there are fewer vertices than for the initial surface, and at least one of them is a connected doubly periodic triangular surface with the periodicity lattice Λ . We call the latter the *connectivity property*, see Figure 9.




FIGURE 9. Connectivity property in the *First Case* of a doubly periodic surface.

The connectivity property of one of the two new surfaces requires some explanation. Let T be the torus generated by the lattice Λ , i.e. $T = \mathbb{R}^2/\Lambda$. Then, initially, $K/\Lambda \cong T\#D$, where $\#$ denotes the connected sum, and D corresponds to the surface formed by the two rhombi γ and $-\gamma$ and the domes over them. After the transformations to the surface described in the proof of the lemma for the resulting surface K' , we obtain that K'/Λ can be always represented as $T\#D'$ for some closed surface D' . The surgery described above preserves T in one of the two disconnected components, thus making its corresponding surface connected (cf. Remark 4.8 below). Since the set of ℓ_{ab} for either surface is a subset of the initial set, we can use the inductive step.

Second case. Suppose there are no triples of vertices a, b, c as in the first case. Consider a vertex u of the surface with the smallest degree d adjacent to vertices v_1, \dots, v_d . The smallest degree must be at least 4, because the first case holds otherwise. We use Theorem 4.7 but we have to be careful with applying it as the field in Theorem 4.7 is not a subfield of the field \mathbb{L} defined earlier. The issue is that some vertices v_i and v_j may belong to the same orbit under the action of the lattice Λ .

Let R_u be a \mathbb{Q} -subalgebra of $\mathbb{K} = \mathbb{Q}(x_u, y_u, z_u, x_{v_1}, y_{v_1}, z_{v_1}, \dots, x_{v_d}, y_{v_d}, z_{v_d})$ generated by all $\ell_{uv_i}, \ell_{v_i v_{i+1}}, 1 \leq i \leq d$. There is a natural homomorphism ψ from R_u to \mathbb{L} mapping all elements of R_u to their corresponding expressions in \mathbb{L} . Note that we cannot automatically define this homomorphism on all elements of \mathbb{K} . For example, when $v_3 = v_1 + \alpha$ and $v_6 = v_4 + \alpha$, the image of $1/(x_{v_3} + x_{v_4} - x_{v_1} - x_{v_6})$ is not defined.

At this point, we use Lemma 4.5 and extend ψ to $\bar{\psi} : \mathbb{K} \rightarrow \bar{\mathbb{L}} \cup \{\infty\}$. The place ϕ can be also extended to a place $\bar{\phi} : \bar{\mathbb{L}} \rightarrow \bar{F} \cup \{\infty\}$. In order to construct this extension, we apply Lemma 4.5 to a subring of all elements of \mathbb{L} whose images under ϕ are finite. For the constructed mapping, $\bar{\phi}(x) = 0$ if $\phi(x) = 0$. Subsequently, if $\phi(x)$ is ∞ , $\bar{\phi}(x)$ must be ∞ as well and $\bar{\phi}$ extends the whole place ϕ .

Using $\bar{\phi}(\infty) = \infty$, we can define the composition $\bar{\phi} \circ \bar{\psi}$. This composition is the place from \mathbb{K} to $\bar{F} \cup \{\infty\}$. Applying Theorem 4.7 to $\bar{\phi} \circ \bar{\psi}$ we conclude that there is i such that the composition and, subsequently, ϕ is finite on $\ell_{v_i v_{i+2}}$.

For the next step, we substitute two triangles of the surface, $uv_i v_{i+1}$ and $uv_{i+1} v_{i+2}$ with $uv_i v_{i+2}$ and $v_i v_{i+1} v_{i+2}$ simultaneously deleting the edge uv_{i+1} and adding the edge $v_i v_{i+2}$. There was no edge $v_i v_{i+2}$ prior to this operation because otherwise the triangle $uv_i v_{i+2}$ would satisfy the case considered above. At the same time we make the same operations for all triangles that are the images of $uv_i v_{i+1}$ and $uv_{i+1} v_{i+2}$ under the action of Λ . As the result we obtain another surface K' such that K'/Λ is topologically the same as K/Λ and has the same number of vertices but the minimum vertex degree of K' is smaller. The place ϕ is still finite on all ℓ_{ab} for edges ab , so all conditions of the lemma still hold.

Observe that the operations in both cases decrease the complexity of the surface. Note that this cannot continue indefinitely since the Euler characteristic is at most 2, and the number of edges and vertex degrees are positive. Thus, at some point we reach the least complex surface for which the statement of the lemma is true. \square

Remark 4.8. In the proof of the First Case, the connectivity property fails for general doubly periodic surfaces. In particular, if the elements of the fundamental group of K/Λ corresponding to periodicity vectors do not commute, two new surfaces may be both disconnected unions of one-periodic pieces. An example is given in Figure 10. Here we show only the bottom half of the surface, which has connected components periodic along α . The top half is attached to the bottom along red triangles and has similar structure, but with connected component periodic along β . This observation will prove crucial in the proof of Theorem A.2 in the Appendix.




FIGURE 10. Non-example to the connectivity property in the *First Case* for general doubly periodic surfaces.

4.5. Proof of Theorem 1.2. Let R' be the \mathbb{Q} -subalgebra of \mathbb{L} obtained by adding all $(\lambda, \lambda)^{-1}$, $\lambda \in \Lambda^*$, to the subalgebra R :

$$R' = R [(\lambda, \lambda)^{-1} \mid \lambda \in \Lambda^*].$$

Let I' be the following ideal in R' :

$$I' = ((\lambda, \lambda)^{-1} \mid \lambda \in \Lambda^*) \triangleleft R'.$$

Assume that $I' \neq R'$. Then, by Krull's theorem (see e.g. [2]), there exists a maximal ideal I , such that $I' \subset I$. Let $F = R'/I$. Since R' contains \mathbb{Q} , field F must contain \mathbb{Q} as well and $\text{char } F = 0$. Let \overline{F} be an algebraic closure of F . The quotient homomorphism $R' \rightarrow F$ satisfies the conditions of Lemma 4.5 for $\Omega = \overline{F}$ so it can be extended to the place $\phi : \mathbb{L} \rightarrow \overline{F} \cup \{\infty\}$. The quotient homomorphism is equal to 0 on all $(\lambda, \lambda)^{-1}$, $\lambda \in \Lambda^*$. Therefore, the place ϕ is infinite on (λ, λ) for all $\lambda \in \Lambda^*$. This implies that the same holds for all non-zero $\lambda \in \Lambda$. On the other hand, the quotient homomorphism is finite on R' . Therefore, we get a contradiction with Lemma 4.6. We conclude that the assumption that $I' \neq R'$ is false.

From above, we have that $I' = R'$. In particular, this implies that $1 \in I'$:

$$(4.1) \quad 1 = \sum_{i=1}^M \frac{r_i}{(\lambda_{i1}, \lambda_{i1})(\lambda_{i2}, \lambda_{i2}) \cdots (\lambda_{ip_i}, \lambda_{ip_i})},$$

where all $\lambda_{ij} \in \Lambda^*$, all $r_i \in R$, and $p_i \geq 1$. After multiplying by the least common multiple of all denominators, the left hand side of (4.1) becomes

$$Z := \prod_{j=1}^N (\lambda_j, \lambda_j) = \prod_{j=1}^N (k_j \alpha + m_j \beta, k_j \alpha + m_j \beta) = \prod_{j=1}^N (k_j^2 (\alpha, \alpha) + 2k_j m_j (\alpha, \beta) + m_j^2 (\beta, \beta)),$$

where $\lambda_j = k_j \alpha + m_j \beta$.

In the same manner we can write down the products in the right hand side of (4.1) times Z . We rewrite the equation via the entries of the Gram matrix of the lattice Λ , which are equal to $g_{11} = (\alpha, \alpha) = s^2$,

$g_{22} = (\beta, \beta) = t^2$, and $g_{12} = (\alpha, \beta) = 0$. We also use the fact that polynomial functions $r_i \in R$ take only rational values on doubly periodic unit triangular surfaces, and denote by $q_i \in \mathbb{Q}$ the value of r_i on the surface (K, θ) . We then have:

$$(4.2) \quad \prod_{j=1}^N (k_j^2 s^2 + m_j^2 t^2) - \sum_{i=1}^M q_i \prod_{j=1}^{N_i} (k_{ij}^2 s^2 + m_{ij}^2 t^2) = 0,$$

where all $N_i < N$. Note that this is the only time in the proof we use the fact that we have unit triangles, and that the periodicity vectors are orthogonal.

We conclude that the polynomial $P(s^2, t^2)$ formed by the equation (4.2) has rational coefficients. Let $x \leftarrow s^2$ and $y \leftarrow t^2$. From above, the polynomial $P(x, y)$ is nonzero since

$$\deg \prod_{j=1}^N (k_j^2 x + m_j^2 y) = N,$$

and the degree of all other terms in (4.2) have degrees $N_i < N$. This completes the proof. \square

4.6. Further applications. Note that polynomials P found in the proof of Theorem 1.2 are quite special. In some cases, with a more careful analysis, one can conclude non-existence of domes for some rhombi whose diagonals are algebraically dependent over \mathbb{Q} . For example, consider the rhombi whose ratio of diagonal lengths is algebraic:

Corollary 4.9. *Let $s \notin \overline{\mathbb{Q}}$ and $t/s \in \overline{\mathbb{Q}}$. Then the unit rhombus $\rho(s, t)$ cannot be domed.*

For example, the corollary implies that $\rho(\frac{1}{\pi}, \frac{1}{\pi})$ and $\rho(\frac{e}{\sqrt{7}}, \frac{e}{\sqrt{8}})$ cannot be domed.

Proof. Suppose $\rho(s, t) \in \mathcal{D}_4$, and let $c := t/s \in \overline{\mathbb{Q}}$. Consider a polynomial $P(s^2, t^2)$, as in the proof of Theorem 1.2. Viewed as a polynomial in $x = s^2$ over $\overline{\mathbb{Q}}$ the leading degree term of P becomes

$$\prod_{j=1}^N (k_j^2 x + m_j^2 c^2 x),$$

so it still has a higher degree than all other terms. Therefore, $s \in \overline{\mathbb{Q}}$, a contradiction. \square

The following is a generalization of the previous corollary:

Corollary 4.10. *Let $s \notin \overline{\mathbb{Q}}$, and let s^2 and t^2 be algebraically dependent with an irreducible polynomial $Q(s^2, t^2) = 0$. Suppose Q^* is the highest degree component of Q . Then the unit rhombus $\rho(s, t)$ cannot be domed unless Q^* is (up to a constant) a product of linear polynomials $(k^2 x + m^2 y)$, where k, m are non-negative integers. Moreover, the rhombus can be domed only if any monomial of Q divides one of the monomials of Q^* .*

For example, the corollary implies that $\rho(\frac{1}{\pi}, \frac{1}{\pi^2})$ cannot be domed. Indeed, the irreducible polynomial $x^2 - y$ does not satisfy the last condition of Corollary 4.10 since the monomial y does not divide a highest degree monomial.

Proof. Suppose $\rho(s, t)$ can be domed. We note that (s^2, t^2) is a root of both P from the proof of Theorem 1.2 and the irreducible polynomial Q . Either P is divisible by Q , or there are polynomials $A, B \in \overline{\mathbb{Q}}[x, y]$ and a nonzero polynomial $D \in \overline{\mathbb{Q}}[x]$ such that $AP + BQ = D$ (see the argument in [12, §1.6, Prop. 2]). The latter case contradicts our assumption on s . We conclude that P is divisible by Q . Therefore, the highest degree component of P given by

$$\prod_{j=1}^N (k_j^2 x + m_j^2 y)$$

must be divisible by Q^* , as desired.

Assume now there is a monomial of Q that does not divide any monomial of Q^* . As we already know, Q^* is a product of $x^u y^v$ and linear polynomials $(k^2 x + m^2 y)$, where k, m are positive integers. If $\deg Q = n$, then Q^* contains all monomials $x^q y^r$ such that $q+r = n$ and $q \geq u$, where $r \geq v$. A monomial of Q that does not divide any monomial of Q^* must be either divisible by x^{n-v+1} or y^{n-u+1} . Without

loss of generality assume the former. Then the largest degree of x in Q is attained on a monomial not from Q^* . Since P is divisible by Q , the largest degree of x in P is attained on a monomial not from the highest total degree component of P . This clearly contradicts the composition of P in (4.2). \square

Remark 4.11. The approach in the corollaries fails in two notable cases we cover in the next section. First, by Proposition 5.2 below, an isosceles triangle Δ with side lengths $(2, 2, 1)$ can be domed if and only if a unit rhombus $\rho(\frac{1}{2}, \sqrt{3})$ can be domed. Since the argument above is not applicable for $\rho(s, t)$ for which $s^2, t^2 \in \mathbb{Q}$, we cannot conclude that Δ cannot be domed, cf. Conjecture 5.1.

The second example where the above approach is inapplicable is the case of planar rhombi $\rho(s, t)$, where $s^2 + t^2 = 4$, see §5.3. In fact, one of the product terms $k_j^2 s^2 + m_j^2 t^2 = (k_j^2 - m_j^2) s^2 + 4m_j^2$ of the leading degree term in P can be equal to 4 when $k_j = \pm 1$ and $m_j = \pm 1$.

5. BIG PICTURE

5.1. Integer-sided triangles. It may seem from the proof of Theorem 1.2, that only integral curves with non-algebraic diagonals cannot be domed. In fact, we believe that only very few integral curves with algebraic diagonals can be domed.

Conjecture 5.1. *An isosceles triangle Δ with side lengths $(2, 2, 1)$ cannot be domed.*

As with many other domes on curves problems, this conjecture turned out to be equivalent to that over a certain unit rhombus (cf. §5.5 below).

Proposition 5.2. *Let Δ be an isosceles triangle with side lengths $(2, 2, 1)$, and let $\rho_\diamond = \rho(\frac{1}{2}, \sqrt{3})$. Then Δ can be domed if and only if ρ_\diamond can be domed.*

Proof. Attach three unit triangles to Δ as in Figure 11. Observe that the boundary of the resulting surface is exactly ρ_\diamond . \square




FIGURE 11. Proof of Proposition 5.2: $\Delta \in \mathcal{D}_5$ if and only if $\rho_\diamond \in \mathcal{D}_4$.

In a contrapositive fashion, let us show that if $\Delta \in \mathcal{D}_5$, then all integral triangles can be domed.

Proposition 5.3. *Let Δ be an isosceles triangle with side lengths $(2, 2, 1)$. If Δ can be domed, then so can every integer-sided triangle.*

Proof. Whenever clear, we denote polygons with their edge length sequence. Observe that all triangles (k, k, k) and all isosceles trapezoids $(1, \ell, 1, \ell + 1)$ can be domed by a plane triangulation. To construct domes over all integer-sided triangles, we use the following rules:

- (1) for integer $k > 1$, $1 \leq \ell < \sqrt{3}k$, two copies of $(k, k, 1)$, one (k, k, ℓ) , and a $(1, \ell, 1, \ell + 1)$ trapezoid, give a triangle $(k, k, \ell + 1)$, via a pyramid over a trapezoid (see Figure 12).
- (2) for integer $k > 1$, $\ell < \sqrt{3}k$, two copies of (k, k, ℓ) , and one $(k, k, 1)$, give a triangle $(\ell, \ell, 1)$ via a tetrahedron.

We now construct all triangles $(k, k, 1)$ one by one, alternating the rules above in the following order:

$$\Delta = (2, 2, 1) \xrightarrow{(1)} (2, 2, 3) \xrightarrow{(2)} (3, 3, 1) \xrightarrow{(1)} (3, 3, 4) \xrightarrow{(2)} (4, 4, 1) \rightarrow \dots$$

Next, we construct domes over general isosceles triangles (k, k, ℓ) , for all $1 \leq \ell < k$, as follows:

$$(k, k, 1) \xrightarrow{(1)} (k, k, 2) \xrightarrow{(1)} (k, k, 3) \xrightarrow{(1)} \dots$$

Finally, we can span (a, b, c) using triangles (k, k, a) , (k, k, b) and (k, k, c) , for $k \geq \max\{a, b, c\}$ large enough. This completes the proof. \square




FIGURE 12. Construction in the first rule: $(2, 2, 1) \rightarrow^{(1)} (2, 2, 3)$.

Remark 5.4. Suppose, contrary to Conjecture 5.1, that a triangle $(2, 2, 1)$ can be domed. That would easily imply Theorem 1.4. Indeed, let ℓ be an integer greater than the radius of rQ_n . By Proposition 5.3, triangle (ℓ, ℓ, r) can also be domed. Symmetrically attach these triangles to all edges in rQ_n , to form a pyramid over rQ_n .

5.2. Flexible surfaces. Let $S \subset \mathbb{R}^3$ be a PL-surface homeomorphic to a sphere, and whose faces are unit triangles. We say that S is a *closed dome*. Such S is called *flexible*, if there is a continuous family $\{S_t, t \in [0, 1]\}$ of (intrinsically) isometric but non-congruent closed domes; closed dome S is called *rigid* otherwise.

We say that an integral curve is *degenerate* if it has two vertices that coincide. Let $\gamma \subset S$ be a closed non-degenerate integral curve along the edges of S . We say that γ is a *separating belt* if $S \setminus \gamma$ is disconnected. We say that a dome S *flexes* γ , if S is flexible with a continuous family $\{\gamma_t \subset S_t, t \in [0, 1]\}$ of non-congruent integral curves.

Conjecture 5.5. *Let S be a closed flexible dome, and $\gamma \subset S$ be a non-degenerate separating belt. Then S does not flex γ .*

Curiously, this general conjecture implies Conjecture 5.1, which at first glance might seem unrelated.

Proposition 5.6. *Conjecture 5.5 implies Conjecture 5.1.*

Proof. By contradiction, suppose Conjecture 5.1 is false. In other words, suppose triangle Δ with side lengths $(2, 2, 1)$ can be domed. By Proposition 5.3, then so can every integer-sided triangle, including triangles with sides $(3, 7, 7)$ and $(4, 7, 7)$, respectively. Four copies of each triangle can be attached to form a flexible *Bricard octahedron* (see e.g. [26, §30.4]), refuting Conjecture 5.5. \square

5.3. Planar unit rhombi. Denote by \mathcal{A} the set of all $a \geq 0$ for which the planar unit rhombus $\rho(a, \sqrt{4 - a^2})$ can be domed. It follows from Lemma 2.1 that $\mathcal{X} \subseteq \mathcal{A}$, so \mathcal{A} is infinite, where \mathcal{X} is defined in (2.1).

Conjecture 5.7. *Set \mathcal{A} is countable.*

The following result is our only evidence in favor of this conjecture.

Proposition 5.8. *Conjecture 5.5 implies Conjecture 5.7.*

Proof. By contradiction, suppose Conjecture 5.7 is false. Since there is a countable number of combinatorial types of triangulated surfaces with quadrilateral boundary, there is one type with infinitely many boundary planar rhombi. Suppose coordinates of the vertices of a rhombus are $(\pm s/2, 0)$ and $(0, \pm t/2)$. The space of surfaces of this particular combinatorial type whose boundary is a planar rhombus is an algebraic set \mathcal{M} for s, t and coordinates of all other vertices of a dome.

As an algebraic set, \mathcal{M} must have a finite number of connected components. Since there are infinitely many values of s for points of \mathcal{M} , there are two values s_1 and s_2 corresponding to two points in the same component. Then a path in this component between these two points provides a flex of a dome. Attaching two copies of a dome along the rhombus boundary, gives a nontrivial deformation of the closed dome. This refutes Conjecture 5.5. \square

Finally, by analogy with Conjecture 5.1, we believe the following claim.

Conjecture 5.9. *We have: $1/2 \notin \mathcal{A}$. In other words, the planar unit rhombus $\rho_\diamond := \rho(1/2, \sqrt{15}/2)$ cannot be domed.*

There is a nice connection between these two conjectures.

Proposition 5.10. *If the planar integral rhombus $2\rho_\diamond$ cannot be domed, then both Conjecture 5.1 and Conjecture 5.9 are true.*

Proof. Observe that $2\rho_\diamond$ can be tiled with four copies of ρ_\diamond . Similarly, $2\rho_\diamond$ can be tiled with two copies of the triangle Δ with sides $(2, 2, 1)$. This implies the result. \square

5.4. Space colorings. Denote by \mathcal{R}^3 a *unit distance graph* of \mathbb{R}^3 , i.e. a graph with vertices points in \mathbb{R}^3 and edges pairs $(x, y) \in \mathbb{R}^3 \times \mathbb{R}^3$ such that $|xy| = 1$. Questions about colorings of \mathcal{R}^3 avoiding certain subgraphs are the main subject of the *Euclidean Ramsey Theory*, see e.g. [14, 30].

Denote by $\chi : \mathcal{R}^3 \rightarrow \{1, 2, 3\}$ a *coloring* of \mathcal{R}^3 . We say that $[xyz] \subset \mathbb{R}^3$ is a *rainbow triangle* in χ , if $[xyz]$ is a unit triangle, and $\chi(x) = 1$, $\chi(y) = 2$, $\chi(z) = 3$.

Proposition 5.11. *Let $\rho = [uvw] \subset \mathbb{R}^3$ be a unit rhombus. Suppose there is a coloring $\chi : \mathcal{R}^3 \rightarrow \{1, 2, 3\}$ with no rainbow triangles, and such that $\chi(u) = \chi(v) = 1$, $\chi(w) = 2$, $\chi(x) = 3$. Then ρ cannot be domed.*

Proof. By contradiction, suppose S is a 2-dimensional triangulated surface with the boundary $\partial S = \rho$. Consider a closed 2-manifold $M := S \cup [uvx] \cup [vwx]$. By *Sperner's Lemma* for closed 2-manifolds, see [25, Cor. 3.1], the number of rainbow triangles in M is even. Note that triangle $[vwx]$ is rainbow, while triangle $[uvx]$ is not. Thus S has at least one rainbow triangle, a contradiction. \square

The idea to use the coloring to prove that some curves cannot be domed can be illustrated with the following conjecture:

Conjecture 5.12. *Let $\rho_\diamond = [uvw] \subset \mathbb{R}^3$ be a unit rhombus defined above. Then there is a coloring as in Proposition 5.11.*

5.5. Domes over multi-curves. One can generalize Kenyon's Question 1.1 to a disjoint union of integral curves $\Upsilon = \gamma_1 \cup \dots \cup \gamma_k$, and ask for a dome over Υ . A special case of this, when Υ is union of two triangles is especially important in view of the *Steinhaus problem*, see §6.5. It would be interesting if the theory of places can be applied to the following problem:

Conjecture 5.13. *There are unit triangles $\Delta_1, \Delta_2 \subset \mathbb{R}^3$, such that $\Upsilon = \Delta_1 \cup \Delta_2$ cannot be domed.*

In the spirit of Proposition 5.2 and the proof of Theorem 1.3, we conjecture that for every integral curve $\gamma \subset \mathbb{R}^3$, whether it can be domed can be reduced to a single rhombus. This is the analogue of "cobordism for domes". Formally, in the notation above, we believe the following holds:

Conjecture 5.14. *For every integral curve $\gamma \in \mathcal{M}_n$, there is a unit rhombus $\rho \in \mathcal{M}_4$, and a dome over $\gamma \cup \rho$.*

In the spirit of the proof Theorem 1.3, there is a natural way to split Conjecture 5.14 into two parts.

Conjecture 5.15. *For every integral curve $\gamma \in \mathcal{M}_n$, there is a finite set of unit rhombi $\rho_1, \dots, \rho_k \in \mathcal{M}_4$, and a dome over $\gamma \cup \rho_1 \cup \dots \cup \rho_k$.*

Conjecture 5.15 is of independent interest. If true, it reduces Conjecture 5.14 to the following claim:

Conjecture 5.16. *For every two unit rhombi $\rho_1, \rho_2 \in \mathcal{M}_4$, there is a unit rhombus $\rho_3 \in \mathcal{M}_4$ and a dome over $\rho_1 \cup \rho_2 \cup \rho_3$.*

5.6. General algebraic dependence. While much of the paper and earlier conjectures are largely concerned with reducing the problem to domes over rhombi, there is another direction one can explore. Namely, one can ask if Theorem 1.2 can be generalized to all integral curves.

Let $\gamma = [v_1 \dots v_n] \in \mathcal{M}_n$ be an integral curve. Denote by $\mathbb{L}_n = \mathbb{Q}[x_{1,3}, x_{1,4}, \dots, x_{n-2,n}]$ the ring of rational polynomials with variables corresponding to diagonals of γ . Let $\text{CM}_n \subset \mathbb{L}_n$ be the ideal spanned by all *Cayley–Menger determinants* on vertices $\{v_1, \dots, v_n\}$, see [7] and [26, §41.6]. We can now formulate the conjecture.

Conjecture 5.17. *Let $\gamma = [v_1 \dots v_n] \in \mathcal{D}_n$ be an integral curve which can be domed, where $n \geq 5$. Denote by $d_{ij} = |v_i v_j|$ the diagonals of γ , where $1 \leq i < j \leq n$. Then there is a nonzero polynomial $P \in \mathbb{L}_n$, which does not belong to the radical of CM_n and such that $P(d_{1,3}^2, d_{1,4}^2, \dots, d_{n-2,n}^2) = 0$.*

This conjecture can be viewed as a direct analogue of Sabitov’s theory of volume being algebraic over squared diagonal lengths, see §6.6. It would be interesting to see if this result can be obtained by expanding our argument in Section 4. Perhaps, Conjecture 5.14 could be used to deduce Conjecture 5.17 from Theorem 1.2.

6. FINAL REMARKS

6.1. Our choice of terminology “dome over curve γ ” owes much to the architectural style of the iconic *geodesic domes* popularized by Buckminster Fuller, and his ill-fated 1960 proposal of a *Dome over Manhattan*, see e.g. [5, pp. 321–324].

6.2. Kenyon formulated Question 1.1 in [20, Problem 2], in an undated webpage going back to at least April 2005. It is best understood in the context of *regular polygonal surfaces* (see e.g. [1, 8]). While we study only the weaker notion (realizations), both the *immersed* and the *embedded* surfaces can be considered, as they add further constraints to the domes. Note also that combinatorially, a dome is a *unit distance complex* of dimension two [18], a notion generalizing the unit distance graphs in §5.4.

6.3. One can view the proof of Theorem 1.2 as a rigidity result for 2-surfaces with unit triangular faces and a single boundary. There are several related rigidity results for surfaces with square and regular pentagonal faces, see e.g. [1, 10].

6.4. It is quite possible that Conjecture 5.12 is false while Conjecture 5.9 is true, since the former seems much stronger. This conjecture is partly motivated by our early attempts to use Monsky’s valuation approach [23, 24], to obtain a negative answer to Kenyon’s Question 1.1.

6.5. The Steinitz Lemma mentioned in §2.7 is a special case of the remarkable 1913 result by Steinitz, motivated by Riemann’s study of conditionally convergent series of real numbers. Bergström’s lower bound $B_2 \geq \sqrt{5}/4$ comes from taking unit edge vectors in the $(k, k, 1)$ triangle, while the matching upper bound is based on elementary arguments in plane geometry. For general d , the best known bound $B_d \leq d$ is due to Grinberg and Sevast’janov [15]. Bárány and others conjecture that $B_d = O(\sqrt{d})$, which would match the Bergström–type lower bound $B_d \geq \sqrt{(d+3)}/4$. We refer to an interesting survey [3] for these results and further references.

6.6. Building on his earlier work and on [7], Sabitov in [27] and [28, §14], proved that a small diagonal in a closed orientable simplicial polyhedron (of any genus), depends algebraically on the lengths of edges of the polyhedron and this dependence is generically non-trivial. Following the proof of Theorem 1.2, we can extend this result to non-orientable polyhedra.

6.7. While Theorem 1.3 is technical, it is natural in view of the existing recreational literature. Notably, in the *Scottish book*, Steinhaus introduced the *tetrahedral chains*, which are polyhedra with a chain-like partition into regular unit tetrahedra. They can be viewed as special types of domes over two triangles, see §5.5. Steinhaus’s 1957 problem asks if tetrahedral chains can be closed, and if they are dense in \mathbb{R}^3 . While the former was given a negative answer in 1959 by Świerczkowski, the latter was partially resolved only recently by Elgersma and Wagon [11]. A somewhat stronger version was later proved by Stewart [33].

Stewart’s paper is especially notable. He uses the ergodic theory of non-amenable group actions, and reproves the (previously known) fact that as a subgroup of $O(3, \mathbb{R})$, the group G of face reflections of a regular tetrahedron is isomorphic to a free product: $G \simeq \mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2$. From there, Stewart showed

that G is dense in $O(3, \mathbb{R})$. One can view this result as an advanced generalization of our Lemma 2.2. The original Steinhaus problem about the group of reflections being dense in the full group $O(3, \mathbb{R}) \rtimes \mathbb{R}^3$ of rigid motions remains open.

Acknowledgements. We are grateful to Arseniy Akopyan, Nikolay Dolbilin, Alexander Gaifullin, Oleg Karpenkov, Oleg Musin, Bruce Rothschild and Ian Stewart, for interesting comments and help with the references. We are especially thankful to Rick Kenyon for encouragement, and to anonymous referee for careful reading of the paper. The authors were partially supported by the NSF.

REFERENCES

- [1] I. M. Alevy, Regular polygon surfaces, preprint (2018), 25 pp.; [arXiv:1804.05452](https://arxiv.org/abs/1804.05452).
- [2] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley, Reading, MA, 1969.
- [3] I. Bárány, On the power of linear dependencies, in *Building bridges*, Springer, Berlin, 2008, 31–45.
- [4] V. Bergström, Zwei Sätze über ebene Vectorpolygone (in German), *Abh. Math. Sem. Univ. Hamburg* **8** (1931), 148–152.
- [5] E. Bookstein, *The Smith Tapes*, Princeton Arch. Press, New York, 2015, 416 pp.
- [6] R. Connelly, A counterexample to the rigidity conjecture for polyhedra, *IHÉS Publ. Math.* **47** (1977), 333–338.
- [7] R. Connelly, I. Sabitov and A. Walz, The bellows conjecture, *Beitr. Algebra Geom.* **38** (1997), 1–10.
- [8] H. S. M. Coxeter, *Regular polytopes* (Second ed.), Macmillan, New York, 1963.
- [9] P. R. Cromwell, *Polyhedra*, Cambridge Univ. Press, New York, 1997.
- [10] N. P. Dolbilin, M. A. Shtan'ko and M. I. Shtogrin, Nonbendability of the covering of a sphere by squares, *Dokl. Akad. Nauk* **354** (1997), no. 4, 443–445.
- [11] M. Elgersma and S. Wagon, Closing a platonic gap, *Math. Intelligencer* **37** (2015), no. 1, 54–61.
- [12] W. Fulton, *Algebraic curves. An introduction to algebraic geometry*, W. A. Benjamin, New York, 1969, 226 pp.
- [13] A. A. Gaifullin and S. A. Gaifullin, Deformations of period lattices of flexible polyhedral surfaces, *Discrete Comput. Geom.* **51** (2014), 650–665.
- [14] R. L. Graham, Euclidean Ramsey theory, in *Handbook of discrete and computational geometry* (Second ed.), CRC Press, Boca Raton, FL, 2004, 239–254.
- [15] V. S. Grinberg and S. V. Sevast'janov, Value of the Steinitz constant, *Functional Anal. Appl.* **14** (1980), 125–126.
- [16] I. Izmetiev, Classification of flexible Kokotsakis polyhedra with quadrangular base, *IMRN* **2017** (2017), 715–808.
- [17] N. W. Johnson, Convex polyhedra with regular faces, *Canadian J. Math.* **18** (1966), 169–200.
- [18] G. Kalai, Some old and new problems in combinatorial geometry I: around Borsuk's problem, in *Surveys in combinatorics*, Cambridge Univ. Press, Cambridge, 2015, 147–174.
- [19] O. N. Karpenkov, On the flexibility of Kokotsakis meshes, *Geom. Dedicata* **147** (2010), 15–28.
- [20] R. Kenyon, *Open problems*, available at <https://gauss.math.yale.edu/~rwk25/openprobs/index.html>; an April 5, 2005 version is cached by the [Wayback Machine](https://www.archive.org/web/20050405000000/https://gauss.math.yale.edu/~rwk25/openprobs/index.html).
- [21] S. Lang, *Introduction to algebraic geometry*, Addison-Wesley, Reading, MA, 1972.
- [22] J. Li, *Flexible Polyhedra: Exploring finite mechanisms of triangulated polyhedra*, Ph.D. thesis, University of Cambridge, 2018, 176 pp.; available at <https://www.repository.cam.ac.uk/handle/1810/271806>.
- [23] P. Monsky, On dividing a square into triangles, *Amer. Math. Monthly* **77** (1970), 161–164.
- [24] P. Monsky, A conjecture of Stein on plane dissections, *Math. Z.* **205** (1990), 583–592.
- [25] O. R. Musin, Extensions of Sperner and Tucker's lemma for manifolds, *J. Combin. Theory, Ser. A* **132** (2015), 172–187.
- [26] I. Pak, *Lectures on discrete and polyhedral geometry*, monograph draft (2009); available at <http://www.math.ucla.edu/~pak/book.htm>.
- [27] I. Kh. Sabitov, Algorithmic solution of the problem of the isometric realization of two-dimensional polyhedral metrics, *Izv. Math.* **66** (2002), 377–391.
- [28] I. Kh. Sabitov, Algebraic methods for the solution of polyhedra, *Russian Math. Surveys* **66** (2011), 445–505.
- [29] B. Schulze, Combinatorial rigidity of symmetric and periodic frameworks, in *Handbook of geometric constraint systems principles*, CRC Press, Boca Raton, FL, 2019, 543–565.
- [30] A. Soifer, *The mathematical coloring book*, Springer, New York, 2009.
- [31] H. Stachel, Flexible polyhedral surfaces with two flat poses, *Symmetry* **7** (2015), 774–787.
- [32] R. P. Stanley, *Enumerative combinatorics*, vol. 1 (second ed.), Cambridge Univ. Press, 2012.
- [33] I. Stewart, Tetrahedral chains and a curious semigroup, *Extracta Math.* **34** (2019), 99–122.

APPENDIX A. DOUBLY PERIODIC SURFACE WITH A THREE-DIMENSIONAL FLEX

A.1. Flex dimension. In [13], Gaifullin and Gaifullin studied the case of doubly periodic surfaces homeomorphic to the plane. In this case they proved a stronger result than Theorem 4.3, that there are two primitive vectors $\lambda, \mu \in \Lambda^*$, such that the place ϕ is finite both on (λ, λ) and (λ, μ) . They concluded the following result:

Theorem A.1 ([13, Thm 1.4]). *Every embedded doubly periodic triangular surface homeomorphic to a plane has at most one-dimensional doubly periodic flex.*

By a *doubly periodic flex* of the triangular surface S we mean a continuous rigid deformation $\{S_t, t \in [0, \delta)\}$ for some $\delta > 0$, which preserves double periodicity, i.e. invariant under the action of $G = \mathbb{Z} \oplus \mathbb{Z}$ (the action of G can also depend on t). The continuity of S is meant with respect to all dihedral angles. Here we identify deformations modulo changes of parameter t and ask for the dimension of the space of flexing at $t = 0$, i.e. when $S_0 = S$. For example, the surface in Figure 9, when triangulated along the shadow lines has only one doubly periodic flex along these lines.

Let us mention that flexible doubly periodic surfaces is an important phenomenon in Rigidity Theory, with *Kokotsakis surfaces* introduced in 1933, giving classical examples, see e.g. [16, 19]. Note that there are doubly periodic polyhedral surfaces whose flexes are not doubly periodic, see [31]. We refer to [29, §25.5] for a recent short survey on rigidity of periodic frameworks, and further references.

A.2. New construction. In [13, Question 1.5], the authors asked if Theorem A.1 can be extended to surfaces which are not homeomorphic to a plane. In this section we give a negative answer to this question by an explicit construction.

Theorem A.2. *There is a doubly periodic triangular surface whose doubly periodic flex is three-dimensional.*

Proof. First, consider a flexible polyhedron F satisfying the following conditions. Polyhedron F must have two faces f_1 and f_2 , both of them centrally symmetric, such that the distance between their centers changes during the flexing and all other faces of F are triangles. To construct such F , take, for example, one of Bricard's flexible octahedra B (see e.g. [22, §2.3] and [26, §30]), and attach two square pyramids to either two of its triangular faces. We illustrate this step in Figure 13, where B is replaced with the usual octahedron for clarity.




FIGURE 13. Illustration of the first step of the construction.

We call *the axis of F* , the line segment connecting the centers of f_1 and f_2 (see Figure 13). Let H be a flexible polyhedron that has two faces congruent to one of the triangular faces of F . We attach




FIGURE 14. Illustration of the second step of the construction.

two copies of F , which we call F_1 and F_2 , to each of these two faces of H (see Figure 14). For our construction we are interested in such H that, when flexing H , the angle σ between the axes of F_1 and F_2 changes and is never zero. Again, a suitable Bricard's octahedron satisfies this condition. We then have a three-dimensional flexing of the whole structure: flexing of F_1 changes the length of the axis of F_1 , flexing of F_2 changes the length of the axis of F_2 , flexing of H changes the angle σ between the axes.

For the next step of the construction we consider F' , the image of F under the central symmetry with respect to the center of f_2 . By \widehat{F} we denote the union of F and F' attached by f_2 (see Figure 15). From now on, we consider only flexes of \widehat{F} such that it stays centrally symmetric with respect to the center of f_2 . Note that during the flex, face f_1 and its counterpart in F' , face f'_1 , are translates of each other and the distance between their centers changes during the flex. One can think of \widehat{F} as a polyhedral version of accordion with bellows such that the sturdy parts of the accordion always stay parallel but the distance between them may change.




FIGURE 15. Illustration of the pieces of the infinite accordion.

Using infinitely many copies of \widehat{F} , we construct a periodic flexible surface S (*infinite accordion*) by attaching f_1 of one copy of \widehat{F} to f'_1 of the next copy of \widehat{F} . See Figure 16 for an illustration (cf. Figure 9). The space of periodic flexes of S is one-dimensional.

Consider a flex $\{S_t\}$ of $S = S_0$ with periodicity vector α . At one of the flexed copies of \widehat{F} we attach H along g_1 . Then attach H along g_2 to another copy of \widehat{F}' which forms its own copy S' of the infinite accordion. Assume the vector of periodicity of S' is β . Now we attach all translates $H + k\alpha$, $k \in \mathbb{Z}$, to surface S , and attach $S' + k\alpha$ to each of them. Then attach $H + k\alpha + m\beta$, $m \in \mathbb{Z}$, to all $S' + k\alpha$, and attach all translates $S + m\beta$ to all translates of H . The resulting surface \mathcal{F} is doubly periodic. It can be flexed in the following two ways:

- by changing lengths of both α and β when flexing $\{S_t\}$ and $\{S'_z\}$, respectively, and
- by changing the angle σ between the axes of S and S' , by simultaneously flexing all copies of H .

Therefore, the space of doubly periodic flexes of \mathcal{F} is three-dimensional. □




FIGURE 16. Illustration of the infinite accordion surface S in the proof, and the three other copies $S + \beta$, S' and $S' + \alpha$, of the doubly periodic surface \mathcal{F} .

Remark A.3. Note that the surface S in the proof is not necessarily embedded. One can similarly construct the analogous embedded surface, by a more careful choice of a flexible polyhedron, cf. [6]. It would be interesting to see if such constructions can have engineering applications. We refer to a recent thesis [22], which reviews several new constructions of embedded flexible polyhedra with larger flex dimensions, and discusses various applications.

Remark A.4. This surface \mathcal{F} is a counterexample to a natural generalization of the Main Lemma 4.6. Let us mention why the proof of the Main Lemma fails for \mathcal{F} . Note that when the initial polyhedra F and H are homeomorphic to a sphere, we have K/Λ is a surface of genus 2 (in the notation of the proof of the Main Lemma), where the elements of the fundamental group corresponding to α and β do not commute, i.e. stand for two different handles of the surface. In particular, the inductive step in the First Case of the proof of the Main Lemma would not work for surgeries since cutting along t would disconnect all translates of S_1 and all translates of S_2 , see Remark 4.8.