

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

Research Colloquium

Research Colloquium 2023

---

## Dementia and Stroke Risk Associated with Brain Artery Luminal Diameters

Jesus D. Melgarejo

*The University of Texas Rio Grande Valley School of Medicine*, [Jesus.Melgarejo@utrgv.edu](mailto:Jesus.Melgarejo@utrgv.edu)

Follow this and additional works at: <https://scholarworks.utrgv.edu/colloquium>



Part of the [Medicine and Health Sciences Commons](#)

---

### Recommended Citation

Melgarejo, Jesus D., "Dementia and Stroke Risk Associated with Brain Artery Luminal Diameters" (2024). *Research Colloquium*. 3.

<https://scholarworks.utrgv.edu/colloquium/2023/talks/3>

This Talk is brought to you for free and open access by the School of Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in Research Colloquium by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

# Dementia and Stroke Risk Associated with Brain Artery Luminal Diameters

## Abstract

**Importance:** It is unclear whether brain artery diameters measured on conventional T2-weighted brain MRI images relate to dementia and stroke outcomes across distinct populations. We aimed this study to evaluate the association of T2-weighted brain artery luminal diameters with dementia and stroke in three distinct population-based studies.

**Methods:** Three longitudinal population-based studies with 8420 adults >40 years old (Northern Manhattan Study [NOMAS] from the United States, and the Rotterdam Study [RS], from the Netherlands, and Three-City, from France) with brain MRI scans obtained between 1999 and 2015. The median follow-up time for clinical events ranged between 7 and 12.5 years. We tested our hypothesis in each cohort separately due to local data-sharing regulations. The exposure variable was brain carotid and basilar artery luminal diameters measured on MRI axial T2-weighted scans. Multivariable hazard ratios (HRs) and their 95% confidence intervals (CI) expressed the risk of dementia and stroke (primary outcomes) associated with the lowest (<5<sup>th</sup>) and highest (>95<sup>th</sup>) percentiles of the rank-normalized brain artery diameters compared to a reference group defined as the diameters distributed between the 5<sup>th</sup> and 95<sup>th</sup> percentiles. Secondary outcomes included total and vascular mortality, and fatal and nonfatal cardiovascular and coronary end points.

**Results:** Among the three cohorts (mean age ranged from 65 to 73 y, ≥57% women), 335 participants developed dementia and 331 strokes. Compared with the reference group, participants with arterial diameters >95<sup>th</sup> percentile had a higher risk of dementia (HR range

1.15-4.50) and any stroke (HR range 1.29-2.03). For secondary outcomes, participants with arterial diameters >95<sup>th</sup> percentile had a consistent higher risk of coronary outcomes, vascular mortality and a composite of any vascular events. The results were less supportive of a higher risk of events among participants with arterial diameters <5<sup>th</sup> percentile except for vascular mortality.

**Conclusions:**

Individuals with dilated brain arteries are at higher risk of dementia and vascular events. Our findings were consistency across distinct populations in spite of using a non-enhanced, conventional T2-weighted MRI sequence. Understanding the underlying physiopathology of the reported associations, particularly with dementia and stroke, might reveal novel vascular contributions to dementia.

**Key Words:** brain artery diameter ■ MRI ■ dementia ■ stroke ■ population-based science