

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

Physics and Astronomy Faculty Publications  
and Presentations

College of Sciences

---

2-3-2021

## Extending the frequency reach of pulsar timing array based gravitational wave search without high cadence observations

Yan Wang

Soumya Mohanty

*The University of Texas Rio Grande Valley*

Zhoujian Cao

Follow this and additional works at: [https://scholarworks.utrgv.edu/pa\\_fac](https://scholarworks.utrgv.edu/pa_fac)



Part of the [Astrophysics and Astronomy Commons](#)

---

### Recommended Citation

Yan Wang et al 2021 ApJL 907 L43 <https://doi.org/10.3847/2041-8213/abd9bd>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Physics and Astronomy Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

## Extending the frequency reach of pulsar timing array based gravitational wave search without high cadence observations

YAN WANG,<sup>1</sup> SOUMYA D. MOHANTY,<sup>2</sup> AND ZHOUIAN CAO<sup>3</sup>

<sup>1</sup>MOE Key Laboratory of Fundamental Physical Quantities Measurements, Hubei Key Laboratory of Gravitation and Quantum Physics, PGMF, Department of Astronomy and School of Physics, Huazhong University of Science and Technology, Wuhan 430074, China

<sup>2</sup>Department of Physics and Astronomy, The University of Texas Rio Grande Valley,  
One West University Boulevard, Brownsville, TX 78520, USA

<sup>3</sup>Department of Astronomy, Beijing Normal University, Beijing 100875, China

### ABSTRACT

Gravitational wave (GW) searches using pulsar timing arrays (PTAs) are assumed to be limited by the typical average observational cadence of 1/(2 weeks) for a single pulsar to GW frequencies  $\lesssim 4 \times 10^{-7}$  Hz. We show that this assumption is incorrect and that a PTA can detect signals with much higher frequencies, which are preserved in the data due to aliasing, by exploiting asynchronous observations from multiple pulsars. This allows an observation strategy that is scalable to future large-scale PTAs containing  $O(10^3)$  pulsars, enabled by the Five-hundred meter Aperture Spherical Telescope and the Square Kilometer Array, without requiring a higher per-pulsar observation cadence. We show that higher frequency GW observations, reaching up to  $4 \times 10^{-4}$  Hz with an SKA-era PTA, have significant astrophysical implications, such as (i) a three orders of magnitude better constraint than current high-cadence observations on GW strain in the [10, 400]  $\mu$ Hz band, and (ii) sensitive tests of the no-hair theorem in the mass range of supermassive black hole binaries using their inspiral, merger, and ringdown signals.

*Keywords:* Pulsar timing method (1305), Gravitational wave astronomy (675)

### 1. INTRODUCTION

The ever growing trove of gravitational wave (GW) signals from compact binary coalescences (Abbott et al. 2016, 2019) collected by the LIGO (LIGO Scientific Collaboration et al. 2015) and Virgo (Acernese et al. 2015) detectors is revealing the GW universe in the  $\sim 10$  Hz to  $10^3$  Hz band. At lower frequencies, the space-based LISA (Amaro-Seoane et al. 2017) mission will target the millihertz band from  $10^{-4}$  Hz to  $10^{-1}$  Hz while pulsar timing arrays (PTAs) are already putting meaningful constraints in the sub- $\mu$ Hz band on the stochastic GW background from an unresolved population of supermassive black hole binaries (SMBHBs) (Lentati et al. 2015; Shannon et al. 2015; Arzoumanian et al. 2016, 2018), continuous waves from resolvable SMBHBs (Zhu et al. 2014; Babak et al. 2016; Aggarwal et al. 2019), and bursts (Wang et al. 2015a; Aggarwal et al. 2020).

The numbers of millisecond pulsars currently being timed by PTA consortia are 47 (Alam et al. 2020, NANOGrav), 26 (Kerr et al. 2020, PPTA), 42 (Desvignes et al. 2016, EPTA), and 65 (Perera et al. 2019, IPTA). Next-generation radio telescopes, namely, the Five-hundred meter Aperture Spherical Telescope (FAST) (Nan et al. 2011; Hobbs et al. 2019) and the Square Kilometer Array (SKA) (Smits et al. 2009; Janssen et al. 2015) will grow the number of well-timed pulsars (noise rms  $\lesssim 100$  ns) to  $O(10^3)$ . Along with a more uniform sky coverage and standardized data spans, this will improve the sensitivity to GWs from resolvable sources by two orders of magnitude (Wang & Mohanty 2017; Wang & Mohanty 2020).

The high frequency limit of the sensitive band for PTA based GW searches is widely assumed (Lentati et al. 2015; Shannon et al. 2015; Arzoumanian et al. 2016, 2018; Zhu et al. 2014; Babak et al. 2016; Aggarwal et al. 2019) to be  $\approx 4 \times 10^{-7}$  Hz, corresponding to the Nyquist rate (Bracewell 2000) associated with the average cadence of timing observations, typically 1/(2 weeks), for individual pulsars in an array. Therefore, attempts at

extending the high frequency limit for resolvable GW sources, to frequency  $> 1 \mu\text{Hz}$ , are all based on high cadence observations of single pulsars (Yardley et al. 2010; Yi et al. 2014; Perera et al. 2018; Dolch et al. 2016). It should be noted here that the actual cadences for pulsars in current PTAs vary over a large range and  $1/(2 \text{ weeks})$  is more representative of its higher end.

In this letter, we show that the high frequency reach of PTAs for resolvable sources is not limited by the Nyquist rate,  $f_{\text{SP}}$ , of single pulsar observations and that the limiting frequency,  $f_{\text{PTA}}$ , can be much higher than assumed so far. The key here is that a higher frequency signal is preserved due to aliasing in the sequence of timing observations from each pulsar and can be unscrambled using asynchronous observations (Wong et al. 2006; Bretthorst 2001) from multiple pulsars.

The lack of synchronicity, an inherent feature of PTA data, can be turned into an observational strategy, which we call *staggered sampling*, to boost the high frequency reach of PTAs. Staggered sampling simply requires the introduction, by design, of relative time shifts between the sequences of timing observations without requiring a change in the individual observational cadence for any pulsar. Unlike high cadence observations, this approach is scalable to future large-scale PTAs with  $O(10^3)$  pulsars since it does not increase the total telescope time consumed by PTA observations.

While the sensitivity of PTA-based GW searches falls with increase in GW signal frequency, an increase in the number of pulsars enhances it. This motivates a first exploration in this letter of the astrophysical implications of high frequency searches with an SKA-era PTA, where staggered sampling could increase the frequency reach to  $\approx 4 \times 10^{-4} \text{ Hz}$  and bridge the gap in coverage of the GW spectrum between PTAs and LISA.

## 2. PRELIMINARIES

In a PTA with  $N_p$  pulsars, the timing residual of the  $I$ -th pulsar after subtracting a best-fit model (excluding GWs) of the pulse time of arrival is given by  $d^I(t) = s^I(t) + n^I(t)$ , where  $s^I(t)$  is the GW induced signal and  $n^I(t)$  is noise.

At the high signal frequencies of interest to us, the samples of  $n^I(t)$  can be assumed to be drawn from an independent and identically distributed Normal random process with zero mean and constant variance  $(\sigma^I)^2$  (i.e., white Gaussian noise). The contribution to  $n^I(t)$  from errors in fitting the timing model are negligible at higher frequencies except at very specific ones (Cutler et al. 2014; Kopeikin & Potapov 2004) such as  $1 \text{ yr}^{-1}$  and harmonics. The latter are ignored in our analysis due to the extremely narrow bands that are affected.

With  $\nu^I(t)$  and  $\nu_0^I(t)$  denoting the pulsar rotation frequencies observed at the Solar System Barycenter and at the pulsar, respectively,  $s^I(t)$  is given by (Estabrook & Wahlquist 1975; Sesana & Vecchio 2010)

$$s^I(t) = \int_0^t dt' z^I(t'), \quad (1)$$

where  $z^I(t) \equiv (\nu^I(t) - \nu_0^I(t))/\nu_0^I(t)$  is the GW induced Doppler shift. For a plane GW arriving from right ascension ( $\alpha$ ) and declination ( $\delta$ ), with polarizations  $h_{+,\times}(t; \theta)$  parametrized by source parameters  $\theta$ ,

$$z^I(t) = \sum_{A=+,\times} F_A^I(\alpha, \delta) \Delta h_A(t; \theta), \quad (2)$$

$$\Delta h_{+,\times}(t; \theta) = h_{+,\times}(t; \theta) - h_{+,\times}(t - \kappa^I; \theta), \quad (3)$$

where,  $F_{+,\times}^I(\alpha, \delta)$  are the antenna pattern functions (Lee et al. 2011) and  $\Delta h_{+,\times}$ , for the two-pulse response (Estabrook & Wahlquist 1975), contain the so-called Earth and pulsar terms that arise from the action of the GW on pulses at the time,  $t$ , of their reception and at the time,  $t - \kappa^I$ , of their emission, respectively.

For a non-evolving circular binary emitting a monochromatic signal,  $\theta$  includes the overall amplitude ( $\zeta$ ), GW frequency ( $f_{\text{gw}}$ ), inclination angle of the binary orbital angular momentum relative to the line of sight ( $\iota$ ), GW polarization angle ( $\psi$ ), and initial orbital phase ( $\varphi_0$ ) (Wang et al. 2014). The time delay  $\kappa^I$  appears as an unknown constant phase offset, called the pulsar phase parameter  $\phi_I$ , for such a source.

To search for resolvable GW sources, we use the likelihood based detection and parameter estimation method described in (Wang et al. 2015b; Wang et al. 2017; Wang & Mohanty 2020) that takes both the Earth and pulsar terms into account. The method partitions the estimation of parameters such that the pulsar phases are either maximized (Wang et al. 2015b) or, as chosen here, marginalized (Wang et al. 2017) semi-analytically, allowing the method to scale to an arbitrarily large  $N_p$ . The remaining parameters are estimated numerically by maximizing the (marginalized) likelihood using Particle Swarm Optimization (Kennedy & Eberhart 1995; Mohanty 2018; Zhu et al. 2016). The maximum value serves as the detection statistic for deciding between the null ( $H_0$ ) and alternative ( $H_1$ ) hypotheses about given data that a signal is absent or present, respectively.

## 3. STAGGERED SAMPLING

Let  $t^I = \{t_i^I\}$ ,  $i = 1, 2, \dots, N^I$ , denote the times at which the residual  $d^I(t)$  is sampled and let their spacing,  $t_{i+1}^I - t_i^I$ , be  $\Delta > 0$  on the average (e.g.,  $\Delta \geq 2 \text{ weeks}$  for IPTA pulsars (Verbiest et al. 2016)). Consider the

set  $\mathcal{T} = \cup_{I=1}^{N_p} t^I$  of sample times in ascending order from all the array pulsars and let  $x_k$ ,  $k = 1, 2, \dots, \sum_{I=1}^{N_p} N^I$ , denote an element of this set. We consider two specific schemes for staggered sampling in our analysis. In both of them, we set  $N^I = N$  to be the same for all the pulsars. This is mainly for reducing the complexity of our codes and not an essential requirement for staggered sampling.

The most straightforward scheme, called *uniform staggered sampling*, is one where  $x_{i+1} - x_i$  is a constant. This implies that the samples of  $d^I(t)$  are uniformly spaced and the sequence of samples from one pulsar has a constant time shift relative to those of others:  $t_i^I = (i-1)\Delta + \delta^I$ , with  $\delta^I = (I-1)\Delta/N_p$ .

In the second scheme, called *randomized staggered sampling*,  $x_{i+1} - x_i$  is a random variable. This is a more realistic situation given the uncertainties inherent in planning astronomical observations. However, as with current PTAs, any real observation strategy would have a target that it seeks to approximate, which is assumed here to be uniform staggered sampling. Therefore, we adopt a reasonable model for randomized staggered sampling in which  $t_i^I$  is replaced by  $t_i^I + c_i^I$ , where  $c_i^I$  is a random variable. In our analysis, we will assume that  $c_i^I$  is drawn from a truncated Cauchy probability density function (pdf) (Papoulis 1984) with a location parameter set to zero, scale factor of 1/3 day, and  $|c_i^I| \leq 7$  days. The heavy tails of this pdf allow large excursions – the 99% inter-percentile range is  $\simeq 10$  days – from the planned observation times of uniform staggered sampling.

The search for individual GW sources is carried out on staggered sampling data as described earlier – no changes are required to the detection and estimation algorithm as it works entirely in the time domain. That this leads to a higher frequency reach is validated directly in this letter using simulated data. While a rigorous analytic treatment of an arbitrary staggered sampling scheme is left to future work, the following argument indicates what the maximum detectable signal frequency should be. Take the trivial case of identical GW induced residuals,  $s^I(t) = s^J(t)$ , observed with uniform staggered sampling. Since the identical residuals lead to a common signal, pooling all data into a single time series will yield samples of the same signal but with a smaller spacing of  $\Delta/N_p$ . Hence, the maximum detectable frequency is  $f_{\text{PTA}} = N_p f_{\text{SP}}$ . Simply pooling the data does not work for the real case of a heterogeneous,  $s^I(t) \neq s^J(t)$  for  $I \neq J$ , set of GW induced residuals but one expects the same limit to hold.

Note that a higher cadence observational strategy to achieve the same high frequency limit as staggered sam-

pling, namely  $N_p f_{\text{SP}}$ , would increase the total telescope time occupied in timing observations by a factor  $N_p$  since each pulsar must be timed with a cadence of  $\Delta/N_p$ . This makes the high cadence strategy extremely costly and unviable for the large  $N_p$  of  $O(10^3)$  in an SKA-era PTA.

#### 4. DETECTION AND PARAMETER ESTIMATION


We use the following simulation setup to show that the staggered sampling schemes described above increase the frequency reach of a PTA. We consider a PTA with  $N_p = 50$  nearest pulsars chosen from the simulated catalog in Smits et al. (2009). The total observation period is set at  $T = 5$  yr with observations spaced  $\Delta = 2$  weeks apart in the case of uniform staggered sampling: this results in  $f_{\text{SP}} \approx 4 \times 10^{-7}$  Hz. ( $T$  is set lower than the typical value of 10 yr or more for PTA data to keep computational costs of the simulation in check.) We consider non-evolving sources with four angular frequencies  $\omega_{\text{gw}}$ : 64 rad/yr ( $3.23 \times 10^{-7}$  Hz), 256 rad/yr ( $1.29 \times 10^{-6}$  Hz), 1024 rad/yr ( $5.16 \times 10^{-6}$  Hz) and 4096 rad/yr ( $2.0656 \times 10^{-5}$  Hz). Note that the last three sources have frequencies  $> f_{\text{SP}}$  and the highest one is very close to the staggered sampling limit of  $f_{\text{PTA}} = N_p f_{\text{SP}} = 4098.09$  rad/yr. The sources are located at  $\alpha = 3.5$  rad and  $\delta = 0.3$  rad in equatorial coordinates. This location corresponds to the lowest degree of ill-posedness in parameter estimation for the SKA-era PTA used in Wang & Mohanty (2017). The inclination angle and the GW polarization angle are given by  $\iota = 0.5$  rad and  $\psi = 0.5$  rad, respectively. The initial orbital phase is set at  $\varphi_0 = 2.89$  rad.

Following the noise model described earlier, the standard deviation of the noise  $n^I(t)$  is set at  $\sigma^I = 100$  ns. The overall amplitude,  $\zeta$ , of the GW signal, which depends on the distance to the source, its chirp mass, and GW frequency, is determined by the specified network signal-to-noise ratio (SNR)  $\rho$ . Here  $\rho^2 = \sum_{I=1}^{N_p} (\rho^I)^2$ , where

$$(\rho^I)^2 = \frac{1}{(\sigma^I)^2 \Delta} \int_0^T (s^I(t))^2 dt \quad (4)$$

is the squared SNR of the GW induced timing residual for the  $I$ -th pulsar.

Fig. 1 shows the distributions of the detection statistic under the  $H_0$  and  $H_1$  hypotheses for both the uniform and the randomized staggered sampling strategies. From these, we estimate the detection probability of a  $\rho = 10$  signal to be  $\gtrsim 90\%$  at a false alarm probability of  $\simeq 1/500$ . The latter corresponds to setting the detection threshold at the largest value of the detection statistic obtained from  $H_0$  data realizations. Within the precision of our simulation, these numbers are fairly in-




**Figure 1.** Distributions of the detection statistic under the  $H_0$  (signal absent) and  $H_1$  (signal present) hypotheses.  $H_0$ : histograms obtained from 500 data realizations are shown for uniform (blue) and randomized (green) staggered sampling.  $H_1$ : the estimated mean (marker) and  $\pm 1\sigma$  deviation (error bar) of the detection statistic, obtained from 200 data realizations, are shown for different signal angular frequencies,  $\omega_{\text{gw}} = 64$  (circle), 256 (square), 1024 (triangle) and 4096 (diamond) rad/yr, and uniform (open markers) or randomized (filled markers) staggered sampling. In all cases, the signal SNR is  $\rho = 10$ . The vertical offset of an error bar or marker is for visual clarity only.

dependent of the staggered sampling strategy and the GW signal frequency. While a two-sample Kolmogorov-Smirnov test on the  $H_0$  distributions does show their apparent relative shift to be statistically significant, this has no noticeable effect on detection probability at the above SNR.

Given that a higher frequency signal has a larger number of cycles in a given observation period, it is natural to ask if it can be detected over shorter observation periods using staggered sampling. We verified this by repeating the above simulations with  $T = 1$  yr and  $\omega_{\text{gw}} = 512$  rad/yr and 1024 rad/yr keeping all else fixed. The detection probabilities had insignificant changes, suggesting that the performance of a staggered sampling based search depends primarily on the SNR of a signal.

Fig. 2 shows the estimated sky locations of the source for uniform staggered sampling ( $T = 5$  yr) and a moderately strong SNR of  $\rho = 20$ . Within the precision of our simulations, the error in localizing a GW source does not show a clear trend with the frequency of the signal. Resolving a trend, if it exists, would require a computationally much more expensive simulation that we leave for future work. The typical localization error in Fig. 2 of  $O(100)$  deg<sup>2</sup> makes searches for optical counter-



**Figure 2.** Distribution of sky location in equatorial coordinates  $(\alpha, \delta)$  for signals with different angular frequencies using uniform staggered sampling. Each panel shows estimated sky locations (dots) from 200 data realizations, each containing an SNR  $\rho = 20$  signal with an angular frequency,  $\omega_{\text{gw}}$ , as noted in the panel. The true location of the GW source is marked by a triangle and the mean of the estimated locations is marked by a cross. The solid contour lines are obtained using 2D Kernel Density Estimation (Botev et al. 2010) and show regions with areas  $\Delta\Omega_{68\%}$  and  $\Delta\Omega_{95\%}$  in which the probabilities of getting estimated locations are 68% and 95%, respectively. For ascending  $\omega_{\text{gw}}$ ,  $\Delta\Omega_{68\%}$  ( $\Delta\Omega_{95\%}$ ) is 103(262), 75(186), 77(305), and 115(306) deg<sup>2</sup>.

parts of GW sources feasible with the Rubin observatory (Ivezić et al. 2019) across the entire range,  $[T^{-1}, f_{\text{PTA}}]$ , of GW frequencies (Liu et al. 2015; Wang & Mohanty 2017).

## 5. ASTROPHYSICAL IMPLICATIONS

For an SKA-era PTA with  $N_p = 10^3$  pulsars and per-pulsar observational cadence of 1/(2 weeks), our results show that staggered sampling can increase  $f_{\text{PTA}} = N_p f_{\text{SP}}$  to as high as  $4 \times 10^{-4}$  Hz. To estimate the achievable sensitivity, we use the same simulated PTA as in Wang & Mohanty (2017), comprised of millisecond pulsars within 3 kpc taken from the synthetic catalog in Smits et al. (2009). From an analysis similar to Fig. 1, we find that the detection probability of a monochromatic signal for the SKA-era PTA is  $\simeq 60\%$  at  $\rho = 10$  for a false alarm probability of  $\simeq 1/50$ . We adopt this as the fiducial value for the minimum detectable SNR averaged over the sky angles  $\alpha$  and  $\delta$ . (The resulting geometrical factor is  $\simeq 1$  for this PTA.) Non-detection of a signal at this SNR will result in a sky-averaged upper

limit on monochromatic GW strain amplitude,

$$h = 8.89 \times 10^{-15} \left( \frac{f_{\text{gw}}}{10^{-6} \text{ Hz}} \right) \left( \frac{T}{5 \text{ yr}} \right)^{-\frac{1}{2}} \left( \frac{\sigma}{100 \text{ ns}} \right), \quad (5)$$


that is about three orders of magnitude lower in the  $[10, 400] \mu\text{Hz}$  band than the current one from high cadence observation of millisecond pulsar J1713+0747 (Dolch et al. 2016).

In the extended frequency range, not only would the inspiral phase of an SMBHB signal be observable but also the merger and ringdown phases. To quantify the sensitivity to each of these phases, we use the luminosity distance,  $D_L$ , for a sky-averaged SNR = 10. (The inclination and polarization angles are also averaged over in the case of inspirals). Since searches for the pulsar and Earth term can be decoupled for a strongly evolving signal (Finn & Lommen 2010), we make the conservative choice of using the SNR of only the Earth term.

The inspiral signal is calculated in the Newtonian approximation (Peters & Mathews 1963) over an observation period  $\min\{20 \text{ yr}, \tau\}$ , where  $\tau$  is the lifetime for the signal frequency to evolve from an initial value  $f_i$  to  $f_{\text{ISCO}}$ , the frequency at the innermost stable circular orbit (ISCO). The merger and ringdown phases are obtained from waveforms computed in the spin aligned effective one body numerical relativity for eccentric binary (SEOBNRE) (Cao & Han 2017; Liu et al. 2020) formalism: the part of the waveform between the instantaneous frequency exceeding  $f_{\text{ISCO}}$  and the instantaneous amplitude attaining its maximum value is defined as the merger, with the subsequent phase being the ringdown. For the latter, only the dominant  $l = 2, |m| = 2$  mode, with its corresponding frequency  $f_{2,2}$ , is used. We consider only circular binaries with zero spin and equal mass components for which the defining parameters are only the chirp mass  $\mathcal{M}_c = 0.435 M$  ( $M$  is the total mass) and, for the inspiral, the chosen  $\tau$ .

Fig. 3 shows  $D_L$  for all the different phases as a function of  $\mathcal{M}_c$  and  $\tau$  along with their characteristic frequencies. We see that for  $\mathcal{M}_c \geq 4.5 \times 10^9 M_\odot$ , the inspiral signal always stays below  $f_{\text{SP}}$  irrespective of  $\tau$ . On the other hand, the inspiral signal for  $\mathcal{M}_c < 4.5 \times 10^9 M_\odot$  would cross  $f_{\text{SP}}$  even if  $f_i < f_{\text{SP}}$ . Table 1 summarizes the distance reach, with the distance ( $D_L = 20$  Mpc) to the Virgo cluster as a baseline, for different signal phases that require the extended frequency range ( $\geq f_{\text{SP}}$ ) of staggered sampling to be observable. Further applications of Fig. 3 are considered below.

The observation of ringdown signals by an SKA-era PTA with staggered sampling could extend the test of the no-hair theorem to the extremely large mass range of SMBHB remnants. For this we consider the test in



**Figure 3.** Luminosity distance,  $D_L$ , (left  $y$ -axis and solid lines) and GW frequency (right  $y$ -axis and broken lines) as a function of chirp mass  $\mathcal{M}_c$  for a maximum observation duration of 20 yr. The gray shaded area covers  $f \in [10^{-9}, f_{\text{SP}} = 4 \times 10^{-7}]$  Hz. Lines in gray scale represent inspirals with different lifetimes as indicated in the legend. Light and dark blue lines represent merger and ringdown, respectively. The frequencies shown are  $f_i$ ,  $f_{\text{ISCO}}$ , and  $f_{2,2}$ .

Isi et al. (2019) that achieves an  $\approx 10\%$  level as defined by the fractional difference in the estimated values of a particular combination of mass and spin parameters measured from the late ringdown and the (SNR  $\approx 14$ ) post peak-amplitude waveforms of GW150914.

For the above SNR of the post peak-amplitude (our ringdown) waveform, staggered sampling will allow the ringdown from a  $\mathcal{M}_c \lesssim 2 \times 10^{10} M_\odot$  system, for which  $f_{2,2} \gtrsim f_{\text{SP}}$ , to be detected out to  $D_L \lesssim 1.32$  Gpc. Given that the corresponding inspiral signal would be extremely loud, SNR = 620 for  $\tau > 2$  yr, the source would be localized well in advance of the ringdown. This would allow a subset of favorably located pulsars to be targeted for significantly better timing over the duration of the ringdown. Assuming the timing residual noise is reduced from 100 nsec to  $\approx 20$  nsec (Feng et al. 2020), the observed ringdown SNR would increase to  $\approx 70$ , leading to a test of the no-hair theorem at the  $\approx 2\%$  level.

Considering a lower mass system such as  $\mathcal{M}_c = 5 \times 10^8 M_\odot$ , Fig. 3 shows that a  $\tau = 5$  yr inspiral signal, with  $f_i \gtrsim f_{\text{SP}}$ , will be detectable out to  $D_L \approx 100$  Mpc. While the corresponding ringdown signal ( $f_{2,2} = 0.02$  mHz) would be too weak for the SKA-era PTA, it would be extremely loud, with SNR  $\approx 220$ , for a concurrently operating LISA. Compared to the fiducial ringdown SNR above, the relative measurement accuracy of all the ringdown parameters – inversely

Signal	$D_L > 20$ Mpc		$D_L > 100$ Mpc		$D_L > 500$ Mpc	
Inspiral	$[2.2, 45] \times 10^8 M_\odot$	[2, 20] yr	$[6, 45] \times 10^8 M_\odot$	[2, 20] yr	$[15, 45] \times 10^8 M_\odot$	[2, 20] yr
Merger	$[4, 45] \times 10^8 M_\odot$	[3.7, 42] d	$[20, 45] \times 10^8 M_\odot$	[19, 42] d	$[37, 45] \times 10^8 M_\odot$	[35, 42] d
Ringdown	$[3, 20] \times 10^9 M_\odot$	[8, 54] d	$[6, 20] \times 10^9 M_\odot$	[16, 54] d	$[13, 20] \times 10^9 M_\odot$	[32, 54] d

**Table 1.** The range of chirp mass,  $\mathcal{M}_c$ , and lifetime (or duration) of different signal phases that lead to a given minimum luminosity distance,  $D_L$ , under staggered sampling. In all cases above,  $f_{\text{ISCO}} \geq f_{\text{SP}}$  for inspiral and merger while  $f_{2,2} \geq f_{\text{SP}}$  for ringdown signals. The minimum  $D_L$  in each case corresponds to the respective lowest lifetime (or duration). The duration for a ringdown is taken as 3 times its exponential damping time scale.

proportional to SNR from Fisher information analysis (Berti et al. 2006) – would improve by a factor of  $\approx 16$ . In combination with the PTA detected inspiral, this would again lead to a stringent test of the no-hair theorem.

In recent work (D’Orazio & Loeb 2020), a scheme for measuring the Hubble constant,  $H_0$ , has been proposed that uses only GW observations by an SKA-era PTA without requiring an electromagnetic counterpart. It relies on measuring both  $D_L$  and the co-moving distance  $D_c = D_L/(1+z)$ ,  $z$  is the cosmological redshift, of a GW source through the effect of GW wavefront curvature on pulsar timing residuals. The governing condition for the measurability of this effect is (D’Orazio & Loeb 2020)  $\gamma = \pi f_{\text{gw}} L^2 / D_c \gtrsim 0.1$ , where  $L$  is the Earth-pulsar distance. For the assumed GW frequency  $f_{\text{gw}} = 10^{-7}$  Hz  $< f_{\text{SP}}$  in D’Orazio & Loeb (2020), achieving a precision of  $\delta H_0 / H_0 \lesssim 30\%$  in this scheme puts a rather stringent observational requirement on the error,  $\delta L$ , in  $L$  of  $\delta L / L \sim 1\%$  at  $L > 10$  kpc. However, the feasibility of this scheme is improved if staggered sampling is used to reach higher  $f_{\text{gw}}$ . For example,  $L$  reduces to 3 kpc for the same relative error if  $f_{\text{gw}} = 10^{-6}$  Hz. This could happen if  $f_{\text{gw}} = f_{\text{ISCO}}$  for an  $\mathcal{M}_c = 2 \times 10^9 M_\odot$  system, which would be detectable out to  $D_L = 2$  Gpc (4 Gpc) for  $\tau = 5$  yr (20 yr), yielding  $\gamma = 0.21$  (0.125).

## 6. DISCUSSION

The impact of the extended frequency reach from staggered sampling on the detectability of a wider range of signals than considered here needs further study. Among

these are higher signal harmonics (Peters & Mathews 1963) from unequal mass SMBHBs that, orbital evolution studies indicate (Sesana 2010), could be driven to high eccentricities ( $\sim 0.3$ ) by interactions with the stellar environment. Independent evidence comes from observations (Dey et al. 2018) of the SMBHB candidate OJ 287 that suggest a binary mass ratio of  $\simeq 122$  and eccentricity 0.657. Besides SMBHBs, oscillation of a network of cosmic strings (Burke-Spolaor et al. 2019), superradiance from axion clouds around isolated black holes (Cardoso et al. 2017), near-zone waves induced by turbulence of solar convection (Bennett & Melatos 2014), and solar oscillation modes (Cutler & Lindblom 1996; Polnarev et al. 2009) could be potential targets for a staggered sampling based search.

## ACKNOWLEDGMENTS

Y. W. gratefully acknowledges support from the National Natural Science Foundation of China (NSFC) under Grants No. 11973024 and No. 11690021, and Guangdong Major Project of Basic and Applied Basic Research (Grant No. 2019B030302001). The contribution of S. D. M. to this paper is supported by NSF Grant No. PHY-1505861. Z. C. gratefully acknowledges support from NSFC under Grant No. 11690023. We thank Wen-Fan Feng for discussions on LISA. We acknowledge the Texas Advanced Computing Center (TACC) at the University of Texas at Austin (www.tacc.utexas.edu) for providing high performance computing resources. We thank the anonymous referee for helpful comments and suggestions.

## REFERENCES

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016, Physical Review Letters, 116, 061102, doi: [10.1103/PhysRevLett.116.061102](https://doi.org/10.1103/PhysRevLett.116.061102)
- . 2019, Physical Review X, 9, 031040, doi: [10.1103/PhysRevX.9.031040](https://doi.org/10.1103/PhysRevX.9.031040)
- Acernese, F., Agathos, M., Agatsuma, K., et al. 2015, Classical and Quantum Gravity, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001)
- Aggarwal, K., Arzoumanian, Z., Baker, P. T., et al. 2019, ApJ, 880, 116, doi: [10.3847/1538-4357/ab2236](https://doi.org/10.3847/1538-4357/ab2236)
- . 2020, ApJ, 889, 38, doi: [10.3847/1538-4357/ab6083](https://doi.org/10.3847/1538-4357/ab6083)

- Alam, M. F., Arzoumanian, Z., Baker, P. T., et al. 2020, *The Astrophysical Journal Supplement Series*, 252, 5, doi: [10.3847/1538-4365/abc6a1](https://doi.org/10.3847/1538-4365/abc6a1)
- Amaro-Seoane, P., Audley, H., Babak, S., et al. 2017, arXiv e-prints, arXiv:1702.00786. <https://arxiv.org/abs/1702.00786>
- Arzoumanian, Z., Brazier, A., Burke-Spolaor, S., et al. 2016, *ApJ*, 821, 13, doi: [10.3847/0004-637X/821/1/13](https://doi.org/10.3847/0004-637X/821/1/13)
- Arzoumanian, Z., Baker, P. T., Brazier, A., et al. 2018, *ApJ*, 859, 47, doi: [10.3847/1538-4357/aabd3b](https://doi.org/10.3847/1538-4357/aabd3b)
- Babak, S., Petiteau, A., Sesana, A., et al. 2016, *MNRAS*, 455, 1665, doi: [10.1093/mnras/stv2092](https://doi.org/10.1093/mnras/stv2092)
- Bennett, M. F., & Melatos, A. 2014, *The Astrophysical Journal*, 792, 55, doi: [10.1088/0004-637x/792/1/55](https://doi.org/10.1088/0004-637x/792/1/55)
- Berti, E., Cardoso, V., & Will, C. M. 2006, *Phys. Rev. D*, 73, 064030, doi: [10.1103/PhysRevD.73.064030](https://doi.org/10.1103/PhysRevD.73.064030)
- Botev, Z. I., Grotowski, J. F., & Kroese, D. P. 2010, *Ann. Statist.*, 38, 2916, doi: [10.1214/10-AOS799](https://doi.org/10.1214/10-AOS799)
- Bracewell, R. N. 2000, *The Fourier transform and its applications* (Boston, The United States of America: McGraw Hill)
- Bretthorst, G. L. 2001, *AIP Conference Proceedings*, 567, 1, doi: [10.1063/1.1381847](https://doi.org/10.1063/1.1381847)
- Burke-Spolaor, S., Taylor, S. R., Charisi, M., et al. 2019, *A&A Rv*, 27, 5, doi: [10.1007/s00159-019-0115-7](https://doi.org/10.1007/s00159-019-0115-7)
- Cao, Z., & Han, W.-B. 2017, *Phys. Rev. D*, 96, 044028, doi: [10.1103/PhysRevD.96.044028](https://doi.org/10.1103/PhysRevD.96.044028)
- Cardoso, V., Pani, P., & Yu, T.-T. 2017, *Phys. Rev. D*, 95, 124056, doi: [10.1103/PhysRevD.95.124056](https://doi.org/10.1103/PhysRevD.95.124056)
- Cutler, C., Burke-Spolaor, S., Vallisneri, M., Lazio, J., & Majid, W. 2014, *PhRvD*, 89, 042003, doi: [10.1103/PhysRevD.89.042003](https://doi.org/10.1103/PhysRevD.89.042003)
- Cutler, C., & Lindblom, L. 1996, *PhRvD*, 54, 1287, doi: [10.1103/PhysRevD.54.1287](https://doi.org/10.1103/PhysRevD.54.1287)
- Desvignes, G., Caballero, R. N., Lentati, L., et al. 2016, *MNRAS*, 458, 3341, doi: [10.1093/mnras/stw483](https://doi.org/10.1093/mnras/stw483)
- Dey, L., Valtonen, M. J., Gopakumar, A., et al. 2018, *ApJ*, 866, 11, doi: [10.3847/1538-4357/aadd95](https://doi.org/10.3847/1538-4357/aadd95)
- Dolch, T., Ellis, J. A., Chatterjee, S., et al. 2016, *Journal of Physics: Conference Series*, 716, 012014, doi: [10.1088/1742-6596/716/1/012014](https://doi.org/10.1088/1742-6596/716/1/012014)
- D’Orazio, D. J., & Loeb, A. 2020, arXiv e-prints, arXiv:2009.06084. <https://arxiv.org/abs/2009.06084>
- Estabrook, F. B., & Wahlquist, H. D. 1975, *General Relativity and Gravitation*, 6, 439, doi: [10.1007/BF00762449](https://doi.org/10.1007/BF00762449)
- Feng, Y., Li, D., Zheng, Z., & Tsai, C.-W. 2020, *Phys. Rev. D*, 102, 023014, doi: [10.1103/PhysRevD.102.023014](https://doi.org/10.1103/PhysRevD.102.023014)
- Finn, L. S., & Lommen, A. N. 2010, *ApJ*, 718, 1400, doi: [10.1088/0004-637X/718/2/1400](https://doi.org/10.1088/0004-637X/718/2/1400)
- Hobbs, G., Dai, S., Manchester, R. N., et al. 2019, *Research in Astronomy and Astrophysics*, 19, 020, doi: [10.1088/1674-4527/19/2/20](https://doi.org/10.1088/1674-4527/19/2/20)
- Isi, M., Giesler, M., Farr, W. M., Scheel, M. A., & Teukolsky, S. A. 2019, *Phys. Rev. Lett.*, 123, 111102, doi: [10.1103/PhysRevLett.123.111102](https://doi.org/10.1103/PhysRevLett.123.111102)
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., et al. 2019, *ApJ*, 873, 111, doi: [10.3847/1538-4357/ab042c](https://doi.org/10.3847/1538-4357/ab042c)
- Janssen, G., Hobbs, G., McLaughlin, M., et al. 2015, in *Advancing Astrophysics with the Square Kilometre Array (AASKA14)*, 37. <https://arxiv.org/abs/1501.00127>
- Kennedy, J., & Eberhart, R. C. 1995, in *Proceedings of the IEEE International Conference on Neural Networks: Perth, WA, Australia, Vol. 4 (IEEE)*, 1942
- Kerr, M., Reardon, D. J., Hobbs, G., et al. 2020, *PASA*, 37, e020, doi: [10.1017/pasa.2020.11](https://doi.org/10.1017/pasa.2020.11)
- Kopeikin, S. M., & Potapov, V. A. 2004, *MNRAS*, 355, 395, doi: [10.1111/j.1365-2966.2004.08331.x](https://doi.org/10.1111/j.1365-2966.2004.08331.x)
- Lee, K. J., Wex, N., Kramer, M., et al. 2011, *MNRAS*, 414, 3251, doi: [10.1111/j.1365-2966.2011.18622.x](https://doi.org/10.1111/j.1365-2966.2011.18622.x)
- Lentati, L., Taylor, S. R., Mingarelli, C. M. F., et al. 2015, *Monthly Notices of the Royal Astronomical Society*, 453, 2576, doi: [10.1093/mnras/stv1538](https://doi.org/10.1093/mnras/stv1538)
- LIGO Scientific Collaboration, Aasi, J., Abbott, B. P., et al. 2015, *Classical and Quantum Gravity*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](https://doi.org/10.1088/0264-9381/32/7/074001)
- Liu, T., Gezari, S., Heinis, S., et al. 2015, *The Astrophysical Journal Letters*, 803, L16. <http://stacks.iop.org/2041-8205/803/i=2/a=L16>
- Liu, X., Cao, Z., & Shao, L. 2020, *Phys. Rev. D*, 101, 044049, doi: [10.1103/PhysRevD.101.044049](https://doi.org/10.1103/PhysRevD.101.044049)
- Mohanty, S. D. 2018, *Swarm Intelligence Methods for Statistical Regression* (Chapman and Hall/CRC)
- Nan, R., Li, D., Jin, C., et al. 2011, *International Journal of Modern Physics D*, 20, 989, doi: [10.1142/S0218271811019335](https://doi.org/10.1142/S0218271811019335)
- Papoulis, A. 1984, *Probability, random variables and stochastic processes*
- Perera, B. B. P., Stappers, B. W., Babak, S., et al. 2018, *MNRAS*, 478, 218, doi: [10.1093/mnras/sty1116](https://doi.org/10.1093/mnras/sty1116)
- Perera, B. B. P., DeCesar, M. E., Demorest, P. B., et al. 2019, *MNRAS*, 490, 4666, doi: [10.1093/mnras/stz2857](https://doi.org/10.1093/mnras/stz2857)
- Peters, P. C., & Mathews, J. 1963, *Physical Review*, 131, 435, doi: [10.1103/PhysRev.131.435](https://doi.org/10.1103/PhysRev.131.435)
- Polnarev, A. G., Roxburgh, I. W., & Baskaran, D. 2009, *Phys. Rev. D*, 79, 082001, doi: [10.1103/PhysRevD.79.082001](https://doi.org/10.1103/PhysRevD.79.082001)
- Sesana, A. 2010, *ApJ*, 719, 851, doi: [10.1088/0004-637X/719/1/851](https://doi.org/10.1088/0004-637X/719/1/851)

- Sesana, A., & Vecchio, A. 2010, *PhRvD*, 81, 104008, doi: [10.1103/PhysRevD.81.104008](https://doi.org/10.1103/PhysRevD.81.104008)
- Shannon, R. M., Ravi, V., Lentati, L. T., et al. 2015, *Science*, 349, 1522, doi: [10.1126/science.aab1910](https://doi.org/10.1126/science.aab1910)
- Smits, R., Kramer, M., Stappers, B., et al. 2009, *Astronomy and Astrophysics*, 493, 1161, doi: [10.1051/0004-6361:200810383](https://doi.org/10.1051/0004-6361:200810383)
- Verbiest, J. P. W., Lentati, L., Hobbs, G., et al. 2016, *MNRAS*, 458, 1267, doi: [10.1093/mnras/stw347](https://doi.org/10.1093/mnras/stw347)
- Wang, J. B., Hobbs, G., Coles, W., et al. 2015a, *MNRAS*, 446, 1657, doi: [10.1093/mnras/stu2137](https://doi.org/10.1093/mnras/stu2137)
- Wang, Y., & Mohanty, S. D. 2017, *Phys. Rev. Lett.*, 118, 151104, doi: [10.1103/PhysRevLett.118.151104](https://doi.org/10.1103/PhysRevLett.118.151104)
- Wang, Y., & Mohanty, S. D. 2020, *Phys. Rev. Lett.*, 124, 169901, doi: [10.1103/PhysRevLett.124.169901](https://doi.org/10.1103/PhysRevLett.124.169901)
- Wang, Y., & Mohanty, S. D. 2020, RAAPTR (Repository of Analysis Algorithms for Pulsar Timing Residuals), [github.com/yanwang2012/RAAPTR](https://github.com/yanwang2012/RAAPTR)
- Wang, Y., Mohanty, S. D., & Jenet, F. A. 2014, *ApJ*, 795, 96, doi: [10.1088/0004-637X/795/1/96](https://doi.org/10.1088/0004-637X/795/1/96)
- . 2015b, *ApJ*, 815, 125, doi: [10.1088/0004-637X/815/2/125](https://doi.org/10.1088/0004-637X/815/2/125)
- Wang, Y., Mohanty, S. D., & Qian, Y.-Q. 2017, *Journal of Physics: Conference Series*, 840, 012058, doi: [10.1088/1742-6596/840/1/012058](https://doi.org/10.1088/1742-6596/840/1/012058)
- Wong, J. L., Megerian, S., & Potkonjak, M. 2006, in *SENSORS, 2006 IEEE*, 777–780, doi: [10.1109/ICSENS.2007.355584](https://doi.org/10.1109/ICSENS.2007.355584)
- Yardley, D. R. B., Hobbs, G. B., Jenet, F. A., et al. 2010, *MNRAS*, 407, 669, doi: [10.1111/j.1365-2966.2010.16949.x](https://doi.org/10.1111/j.1365-2966.2010.16949.x)
- Yi, S., Stappers, B. W., Sanidas, S. A., et al. 2014, *MNRAS*, 445, 1245, doi: [10.1093/mnras/stu1826](https://doi.org/10.1093/mnras/stu1826)
- Zhu, X.-J., Wen, L., Xiong, J., et al. 2016, *MNRAS*, 461, 1317, doi: [10.1093/mnras/stw1446](https://doi.org/10.1093/mnras/stw1446)
- Zhu, X.-J., Hobbs, G., Wen, L., et al. 2014, *MNRAS*, 444, 3709, doi: [10.1093/mnras/stu1717](https://doi.org/10.1093/mnras/stu1717)